



ORIGINAL ARTICLE

Two factors that can increase the length of hospital stay of patients with dementia

Mary Duah-Owusu White^{a,*}, Michael Vassallo^b, Fiona Kelly^c, Samuel Nyman^d^a Bournemouth University, United Kingdom^b University Hospitals Dorset NHS Foundation Trust, United Kingdom^c Queen Margaret University, United Kingdom^d Bournemouth University Clinical Research Unit, Bournemouth University, Bournemouth, Dorset, United Kingdom

ARTICLE INFO

Article history:

Received 22 February 2022

Accepted 20 October 2022

Available online 18 November 2022

Keywords:

Dementia

Hospital

Length of stay

Falls

Discharge

ABSTRACT

Objectives: Patients with dementia are at greater risk of a long hospital stay and this is associated with adverse outcomes. The aim of this service evaluation was to identify variables most predictive of increased length of hospital stay amongst patients with dementia.

Methods/Design: We conducted a retrospective analysis on a cross-sectional hospital dataset for the period January–December 2016. Excluding length of stay less than 24 h and readmissions, the sample comprised of 1133 patients who had a dementia diagnosis on record.

Results: The highest incidence rate ratio for length of stay in the dementia sample was: (a) discharge to a care home (IRR: 2.443, 95% CI 1.778–3.357), (b) falls without harm (IRR: 2.486, 95% CI 2.029–3.045).

Conclusions: Based on this dataset, we conclude that improvements made to falls prevention strategies in hospitals and discharge planning procedures can help to reduce the length of stay for patients with dementia.

© 2022 SEGG. Published by Elsevier España, S.L.U. All rights reserved.

Dos factores que pueden aumentar la duración de la estancia hospitalaria de los pacientes con demencia

RESUMEN

Objetivos: Los pacientes con demencia tienen mayor riesgo de una estancia hospitalaria prolongada y esto se asocia con resultados adversos. El objetivo de esta evaluación del servicio fue identificar las variables predictivas de una mayor duración de la estancia hospitalaria de los pacientes con demencia.

Métodos/diseño: Realizamos un análisis retrospectivo de un conjunto de datos hospitalarios transversales en el período enero-diciembre de 2016. Excluyendo la estancia hospitalaria inferior a 24 h y los reingresos, la muestra estaba compuesta por 1133 pacientes que tenían un diagnóstico de demencia registrado.

Resultados: Los pacientes con demencia con estancia más prolongada presentaban mayor tasa de incidencia de: a) alta a una residencia (TIR: 2.443, IC 95% 1.778–3.357) b) caídas sin daño (TIR: 2.486, IC 95% 2.029–3.045)

Conclusiones: En base a estos resultados, concluimos que las mejoras realizadas en las estrategias de prevención de caídas en los hospitales y los procedimientos de planificación del alta pueden ayudar a reducir la duración de la estancia de los pacientes con demencia.

© 2022 SEGG. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Demencia

Hospital

Duración de la estancia

Caídas

Alta

Introduction

Dementia is a condition that affects the brain and subsequently results in memory and cognitive difficulties.¹ In the UK, 885,000 people have dementia² and are more likely to be admitted into hospital compared with their peers without dementia.³

* Corresponding author.

E-mail address: mduahowusuwhite@bournemouth.ac.uk (M. Duah-Owusu White).

Also, patients with dementia experience longer hospital stays than patients without dementia.⁴ A likely consequence of this increased length of stay is the possibility of acquiring nosocomial infections⁵ which may require the use of antibiotics and subsequently result in the development of antimicrobial resistance.⁶ This situation can further increase patient length of stay. In addition to having extended hospital stays, researchers have reported that people with dementia or cognitive impairment experience adverse outcomes in hospitals.^{7–10} Some of the adverse outcomes include malnourishment,⁸ the occurrence of delirium,⁷ complications after surgical procedures⁹ and the development of pressure sores.¹⁰ Patients who stay in hospital for a longer period of time are also likely to experience functional decline.¹¹ Other researchers have suggested that longer hospital stays can make patients vulnerable to harmful medication reactions.¹² Extended hospital stays also reduces the availability of beds for those requiring urgent admission⁵ at an increased cost to the National Health Service (NHS).¹³

Therefore, it is necessary to investigate the factors that influence length of stay of patients with dementia in hospital to identify appropriate strategies needed to improve patient outcomes and minimise the time they are in hospital. One way of improving patient outcomes is to generate knowledge from the evaluation of services so as to enhance bedside practice and organisational culture.¹⁴ The aim of this service evaluation is to investigate the impact of demographic (e.g. age), administrative (e.g. admission day) and clinical factors (e.g. early warning signal) on the hospital stay of patients with dementia. Most of the variables investigated in this study have been previously reviewed. However, most of the previous studies have been limited by small sample sizes that reduced their statistical power and generalisability (e.g. a study on falls and length of stay for 622 patients with a range of medical conditions¹⁵). We explored factors not previously investigated with a large sample including the patient's admission method (i.e. whether the patient attended A&E themselves or were referred by a doctor or transferred from another hospital) and also the various categories of pressure sores.

Materials and methods

This study was a cross-sectional retrospective analysis of a hospital dataset. The dataset contained information on 35,792 patients who were admitted and discharged between 01/01/2016 and 31/12/2016 at a hospital in the south west of the UK. This hospital had a Care Quality Commission rating of 'requires improvement' in October/November 2015 (i.e. prior to the collection of the data). Findings from the report indicated that the hospital needed to improve hospital care for older patients.¹⁶ The integrity of the dataset (i.e. compliance with NHS regulations) has been discussed in detail elsewhere.¹⁷ This process involved the use of trained clinical coders who followed the UK coding rules for entering aspects of the data which are sent to the Secondary Uses Service of the NHS Digital Department. Patient data was excluded if they had been readmitted because findings from a recent study indicated that such patients are likely to have more co-morbidities and receiving a variety of medication.¹⁸ Similarly, people who were admitted for a short period (i.e. less than 24 h) were excluded as they are likely to be healthier than their peers who have been on the ward for a longer period of time.¹⁹ Our sample comprised of patients with dementia ($n = 1133$). Dementia was defined as the existence of a dementia coding applied to a spell. The following codes were used for dementia: Alzheimer's disease, Multi-infarct dementia or vascular dementia, dementia due to other causes such as Picks disease, dementia with an unnamed aetiology, Alzheimer's Disease with an onset in older people and other types of Alzheimer's

disease.¹ Patients with dementia may have been lost because we know that when it is not the principal diagnosis, it is sometimes not recorded/coded.

Statistical analysis

Length of stay was defined as the time the patient was admitted until their discharge. The following variables were evaluated for their impact on the patient's length of stay (Table 1).

These variables were selected because they are routinely collected in hospitals. The information available in the dataset were recorded by nursing and medical staff as well as clinical coders at the hospital.

Descriptive, bivariate and multivariate analysis were conducted using SPSS 19. The following non-parametric tests were used as the dependent variable (length of stay) was skewed and continuous: Mann–Whitney, Spearman and Kruskal–Wallis Test.²⁰ Negative Bionomial Regression was used to analyse the data because findings from a previous study indicated that it was the most appropriate multivariate test for a dependent variable that is skewed and continuous.²¹ Missing data was handled using the exclude pairwise option in SPSS.

Ethics

Ethics approval was obtained from the University (Ethics ID: 23681) as the study was classified as a service evaluation. Before the NHS trust provided the dataset, it was first anonymised, encrypted and password-protected. The information was transferred onto a password-protected university laptop for analysis.

Results

Descriptive statistics

The length of stay for dementia sample was high at an average of 344.9 h. Patients with dementia were on average 85.5 years old. Some patients with dementia were admitted from a care home (36.7%). Some patients with dementia were discharged to a care home (40.4%). Patients with dementia experienced falls without harm (9.7%). Other patients with dementia experienced minor harm during a fall (4%). Some patients with dementia also experienced falls with moderate and major harm (0.4%). Some patients with dementia had pre-admission pressure sores of category 1 (8.4%), 2 (9.1%), 3 (2.3%) and 4 (0.9%). Some patients with dementia had post-admission pressure sores of category 1 (5.1%), 2 (7.1%), 3 (0.9%) and 4 (0.4%) (Table 2).

Bivariate statistics

In the dementia sample, the following variables were all significantly associated with length of stay (age, discharge method, admission from a care home, discharge to a care home, falls without harm, falls with minor harm, MUST scores, Category 1 and 3 pre-admission pressure sores, Category 1, 2, 3, 4 and un-stageable post-admission pressure sores) (Table 3).

Multivariate statistics

For the multivariate analysis only variables that were significant in the bivariate analysis were entered in the Negative Bionomial Regression Model. In the dementia sample, the highest incidence rate ratio for length of stay was: (a) discharge to a care home (IRR: 2.443, 95% CI 1.778–3.357), (b) falls without harm (IRR: 2.486, 95% CI 2.029–3.045). Findings from the multivariate analysis indicate

Table 1
Variables evaluated for their impact on the patient's length of stay.

Age	Gender	Admission method (e.g. referral from a doctor or hospital)	Discharge method (e.g. based on clinical advice or a court (e.g. mental health tribunal))	Admission from a care home
Discharge to a care home	Admission day	Admission is a bank holiday	First Early Warning Signal (based on physiological measurements such as blood pressure and respiration)	Falls without harm
Falls with minor harm	Falls with moderate and major harm	MUST (Malnutrition Universal Screening Tool which is based on clinical factors such as the patient's weight and height) scores on admission	Category 1, 2, 3, 4 and unstageable pre and post-admission pressure	Waterlow scores on admission (this assesses the patient's risk of developing pressure sores by for example, looking at the patient's ability to mobilise and eat).

that some clinical variables (Medium MUST scores, Category 1, 3 pre-admission pressure sores and Category 3, 4 as well as unstageable post-admission pressure sores) were not associated with length of stay in the dementia sample (Table 4).

Discussion

The aim of this study was to investigate factors associated with the length of hospital stay of patients with dementia. Findings from this study indicated that two modifiable factors associated with an increased length of hospital stay amongst patients with dementia were discharge to care homes and experiencing an inpatient fall. Our analysis was novel in including admission method, which was not associated with length of hospital stay. In addition, we were able for the first time to demarcate in a large sample the level of harm sustained by an inpatient fall and also categories of pressure sores.

This study showed that the decision to discharge a patient to a care home was significantly associated with an increased length of stay. Previous literature has shown that the decision to discharge an older patient to a care home could be triggered by the needs of the patient (e.g. requiring additional nursing care) or the patient being unsafe in their own home (e.g. possibility of having a fall with an extended wait for a medical response).²² In addition to the decision making processes required for the safe discharge of patients, there is a serious shortage of care home beds due to the high demand for community services and residential facilities.²³ Appropriate national strategies are therefore needed to ease the pressures on care homes and community services. Furthermore, prolonged hospital stay could potentially be as a result of the appearance of new conditions (due to functional and cognitive decline, delirium, malnutrition) that can lead to institutionalisation. On the other hand, the difficulty of obtaining a place in a nursing home can also lead to a prolonged hospital stay. Based on the evidence available, we believe that prolonged hospital stay is due to difficulty in obtaining a place in a nursing home.

This study indicated that falls with and without harm were significantly associated with an increased hospital stay. The findings of this research is consistent with that of previous researchers who have found an association between falls and the hospital stay of patients with and without dementia.^{15,24} In addition to increasing the length of hospital stay, falls can cause physical injuries,²⁵ increase hospital costs²⁶ and have adverse psychological consequences.²⁷ The inadequacy of current fall prevention measures for people with dementia is therefore concerning.²⁸ There is an urgent need to develop innovative falls prevention interventions for patients with dementia in the hospital setting. One strategy would be to adjust the hospital stay to what is strictly necessary and to then develop care resources that are alternatives to conventional hospitalisation such as hospital at home, primary care, and home-based physiotherapy.

The authors found that some types of pressure sores were associated with the length of stay in the dementia sample. This finding is consistent to previous research which has found an association between pressure sores and hospital length of stay.²⁹ Also, it has been reported that the development of pressure sores could lead to: an increase use of a hospital's financial resources,³⁰ cause the patient pain, have negative psychological consequences (e.g. make patient feel sad) and also limit the patient's ability to engage in social activities.³¹ The incidence of post-admission pressure sores needs to be minimised. The presence of pressure ulcers in people with advanced dementia usually indicates a shorter survival period,³² so the increase in hospital stay may be more related to the functional grade of dementia (GDS 7, FAST >7c) than to the ulcers themselves. With advanced dementia, social activities may not be valued when compared to interaction with family members and usual caregivers.

The main strength of this study was its large sample size. Although a previous study used a large dementia dataset, the authors did not investigate the impact of the various types of pressure sores on the hospital length of stay of patients with dementia.²⁴ They also did not look at the relationship between the various categories of falls and length of hospital stay.²⁴ Another strength of the study is the assessment of the administrative variables (i.e. if they are referred by their doctor or not, if they were admitted during public holiday). It seems to be something interesting and to be considered in future studies. In the current study, although the authors could not validate the dementia diagnosis, the dataset used for this service evaluation complied with NHS quality assessments. Staff documentation of variables such as falls and MUST scores can be prone to errors which could not be eliminated in the analysis of this retrospective study. Also, data regarding the occurrence of delirium was not readily available in the dataset. In future studies, other strategies such as prevention of delirium should be investigated to ascertain whether they play a role in the length of stay of patients with dementia, as well as prevention of falls or discharge planning procedures.

To conclude, the analysis was conducted in a hospital which had a Care Quality Commission rating of 'requires improvement' prior to the collection of data. The findings may therefore not be transferable to hospitals which do not have a similar Care Quality Commission rating. Based on this dataset, we conclude that some of the factors that can be modified in order to reduce the hospital stay of patients with dementia are discharge planning and the prevention of falls. Other alternatives to conventional hospitalisation (hospital at home, primary care, physiotherapy) need to be considered.

Implications for clinical practice

Our research indicates that nursing/care staff can reduce the length of hospital stay of patients with dementia by focusing their

Table 2
Descriptive statistics for inpatients with dementia.

Variable	n = 1133(%)
<i>Discharge age</i>	Mean = 85.5
<i>Gender</i>	
Male	460 (41%)
Female	673 (59%)
<i>Admission methods</i>	
Attendance at Emergency unit	706 (62%)
Referral from a medical practitioner	400 (35%)
Referral from a consultant	2 (0.2%)
Other (e.g. hospital)	25 (2%)
<i>Discharge methods</i>	
Based on clinical advice	984 (87%)
Self-discharge	2 (0.2%)
Discharged by institution (e.g. court)	–
Death	147 (13%)
<i>Admissions from a care home</i>	
Yes	416 (37%)
No	717 (63%)
<i>Discharges to a care home</i>	
Yes	458 (40%)
No	675 (60%)
<i>Admission day</i>	
Friday	161 (14%)
Monday	155 (13%)
Saturday	166 (15%)
Sunday	192 (17%)
Thursday	170 (15%)
Tuesday	155 (14%)
Wednesday	134 (12%)
<i>Admission is a bank holiday</i>	
Yes	28 (2%)
No	1105 (98%)
<i>First Early Warning Signal</i>	
High (>6)	61 (5%)
Medium (5–6)	106 (10%)
Low (0–4)	808 (71%)
Missing	158 (14%)
<i>Falls without harm</i>	
Yes	110 (10%)
No	1,023 (90%)
<i>Falls with minor harm</i>	
Yes	45 (4%)
No	1088 (96%)
<i>Falls major and moderate</i>	
Yes	5 (0.4%)
No	1128 (99.6%)
<i>MUST scores</i>	
High (>1)	261 (23%)
Medium (1)	44 (4%)
Low (0)	816 (72%)
Missing	12 (1%)
<i>Category 1 preadmission pressure sores</i>	
Yes	95 (8%)
No	1038 (92%)
<i>Category 2 preadmission pressure sores</i>	
Yes	103 (9%)
No	1030 (91%)
<i>Category 3 preadmission pressure sores</i>	
Yes	26 (2%)
No	1107 (98%)
<i>Category 4 preadmission pressure sores</i>	
Yes	10 (1%)
No	1123 (99%)
<i>Un-stageable preadmission pressure sores</i>	
Yes	3 (0.3%)
No	1130 (99.7%)

Table 2 (Continued)

Variable	n = 1133(%)
<i>Category 1 Postadmission pressure sores</i>	
Yes	58 (5%)
No	1075 (95%)
<i>Category 2 Postadmission pressure sores</i>	
Yes	80 (7%)
No	1053 (93%)
<i>Category 3 Postadmission pressure sores</i>	
Yes	10 (0.9%)
No	1123 (99.1%)
<i>Category 4 Postadmission pressure sores</i>	
Yes	5 (0.4%)
No	1128 (99.6%)
<i>Un-stageable Postadmission pressure sores</i>	
Yes	4 (0.4%)
No	1129 (99.6%)
<i>Waterlow scores</i>	
High (>15)	878 (77%)
Medium (11–15)	179 (16%)
Low (0–10)	66 (6%)
Missing	10 (1%)

Table 3
Bivariate statistics for inpatients with dementia (N = 1133).

Variable	U or χ or rho value/p value/Z value
Age	.072/.015
Gender	153,033.5/.745/–.325
Admission methods	5.250/.154
Discharge methods	11.866/.003
Admissions from a care home	111,724.5/.000/–7.047
Discharge to a care home	133,349.5/.000/–3.927
Admission Day	4.856/.562
Admission is a bank holiday	15,275.5/.909/–.114
First Early Warning Signal	.562/.755
Falls without harm	24,117.5/.000/–9.858
Falls with minor harm	9828.5/.000/–6.812
Falls major and moderate	2143.0/.354/–.927
MUST scores	6.045/.049
Category 1 Preadmission pressure sores/	42,454.0/.025/–2.244
Category 2 Preadmission pressure sores/	47,821.0/.099/–1.650
Category 3 Preadmission pressure sores	10,376.0/.015/–2.435
Category 4 Preadmission pressure sores	4523.0/.289/–1.060
Unstageable Preadmission pressure sores	1214.5/.396/–.849
Category 1 Postadmission pressure sores	16,831.5/.000/–5.909
Category 2 Postadmission pressure sores	25,481.5/.000/–5.897
Category 3 Postadmission pressure sores	2598.0/.003/–2.929
Category 4 Postadmission pressure sores	1252.5/.032/–2.147
Unstageable Postadmission pressure sores	848.5/.031/–2.158
Waterlow scores	.172/.917

Table 4
Negative binomial regression for inpatients with dementia with length of stay as the dependent variable (N = 1133).

Variable	IRR/p value	95% confidence interval	
		Lower	Upper
<i>Discharge age</i>	1.000/.992	.992	1.008
<i>Ref: clinical advice</i>			
Death as a discharge cause	1.228/.028	1.022	1.476
Self-discharge	.541/.390	.134	2.193
Admissions from a care home	.252/.000	.182	.349
Discharges to a care home	2.443/.000	1.778	3.357
Falls without harm	2.486/.000	2.029	3.045
Falls with minor harm	1.750/.000	1.284	2.384

Table 4 (Continued)

Variable	IRR/p value	95% confidence interval	
		Lower	Upper
Ref: Low MUST score			
High MUST scores	1.163/.037	1.009	1.340
Medium MUST scores	.945/.721	.694	1.287
Category 1 Preadmission pressure sores/	1.189/.113	.960	1.474
Category 3 Preadmission pressure sores	.978/.916	.652	1.467
Category 1 Postadmission pressure sores/	1.787/.000	1.363	2.342
Category 2 Postadmission pressure sores/	1.621/.000	1.275	2.060
Category 3 Postadmission pressure sores/	.966/.915	.508	1.837
Category 4 Postadmission pressure sores/	1.641/.279	.669	4.028
Unstageable Postadmission pressure sores	1.814/.242	.669	4.916

efforts on minimising in-patient falls and improving the discharge planning process. Hospital/Trust policies need to be improved by incorporating findings from robust patient focused research into their formation and application.

Funding

The research was funded by Bournemouth University and the University Hospitals Dorset NHS Foundation Trust via a match-funded PhD studentship. The findings of the study do not represent the views of the University or the Hospital.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

We would like to thank Dave Moore, Anton Parker, Leeanne Dove and Camilla Axtell for facilitating the receipt of the hospital dataset.

References

- World Health Organisation. International Statistical Classification of Diseases and Related Health Problems 10th Revision, 2010; Available from: <http://apps.who.int/classifications/icd10/browse/2010/en>.
- Wittenberg R, Hu B, Jagger C, Kingston A, Knapp M, Comas-Herrera A, et al. Projections of Older People Living with Dementia and Costs of Dementia Care in the United Kingdom 2019–2040. CPEC Working Paper 5. London School of Economics and Political Science; 2019.
- Shepherd H, Livingston G, Chan J, Sommerlad A. Hospitalisation rates and predictors in people with dementia: a systematic review and meta-analysis. *BMC Med*. 2019;17:130.
- Möller T, Stocker H, Wei W, Perna L, Brenner H. Length of hospital stay and dementia: a systematic review of observational studies. *Int J Geriatr Psychiatry*. 2019;34:8–21.
- Toh HJ, Lim ZY, Yap P, Tang T. Factors associated with prolonged length of stay in older patients. *Singap Med J*. 2017;58:134–8.
- Hyunyoung B, Cho M, Kim S, Hwang H, Song M, Yoo S. Analysis of length of hospital stay using electronic health records: a statistical and data mining approach. *PLOS ONE*. 2018;13, e0195901.
- Travers C, Byrne GJ, Pachana NA, Klein K, Gray LC. Prospective observational study of dementia in older patients admitted to acute hospitals. *Aust J Ageing*. 2014;33:55–8.
- Fogg C, Meredith P, Bridges J, Gould GP, Griffiths P. The relationship between cognitive impairment, mortality and discharge characteristics in a large cohort of older adults with unscheduled admissions to an acute hospital: a retrospective observational study. *Age Ageing*. 2017;46:794–801.
- Hu C-J, Liao CC, Chang CC, Wu CH, Chen TL. Postoperative adverse outcomes in surgical patients with dementia: a retrospective cohort study. *World J Surg*. 2012;36:2051–8.
- Bail K, Berry H, Grealish L, Draper B, Karmel R, Gibson D, et al. Potentially preventable complications of urinary tract infections, pressure areas, pneumonia, and delirium in hospitalised dementia patients: retrospective cohort study. *BMJ Open*. 2013;3:e002770.
- van Vliet M, Huisman M, Deeg DJH. Decreasing hospital length of stay: effects on daily functioning in older adults. *J Am Geriatr Soc*. 2017;65:1214–21.
- Morimoto T, Sakuma M, Matsui K, Kuramoto N, Toshiro J, Murakami J, et al. Incidence of adverse drug events and medication errors in Japan: the JADE study. *J Gen Intern Med*. 2011;26:148–53.
- Annear MJ, Tierney LT, Vickers JC, Palmer AJ. Counting the cost of dementia-related hospital admissions: a regional investigation. *Aust J Ageing*. 2016;35:E32–5.
- Moule P, Armoogum J, Douglass E, Taylor J. Evaluation and its importance for nursing practice. *Nurs Stand*. 2017;31:55.
- Dunne TJ, Gaboury I, Ashe MC. Falls in hospital increase length of stay regardless of degree of harm. *J Eval Clin Pract*. 2014;20:396–400.
- Care Quality Commission, The Royal Bournemouth and Christchurch Hospitals NHS Foundation Trust Quality Report. 2016.
- Duah-Owusu White M. Improving the short-term management of patients with dementia admitted to hospital. Bournemouth University; 2021.
- Glans M, Kragh Ekstam A, Jakobsson U, Bondesson Å, Midlöv P. Risk factors for hospital readmission in older adults within 30 days of discharge – a comparative retrospective study. *BMC Geriatr*. 2020;20:467.
- Trentino KM, Swain SG, Burrows SA, Sprivilis PC, Daly FF. Measuring the incidence of hospital-acquired complications and their effect on length of stay using CHADx. *Med J Aust*. 2013;199:543–7.
- Field A. Discovering statistics using SPSS. Sage publications; 2005.
- Carter EM, Potts HW. Predicting length of stay from an electronic patient record system: a primary total knee replacement example. *BMC Med Inform Decis Mak*. 2014;14:26.
- Rhynas SJ, Garrido AG, Burton JK, Logan G, MacArthur J. New care home admission following hospitalisation: How do older people, families and professionals make decisions about discharge destination? A case study narrative analysis. *Int J Older People Nurs*. 2018;13:e12192.
- Kable A, Chenoweth L, Pond D, Hullick C. Health professional perspectives on systems failures in transitional care for patients with dementia and their carers: a qualitative descriptive study. *BMC Health Serv Res*. 2015;15:567.
- Kasteridis P, Goddard MK, Jacobs R, Santos R, Mason A. The impact of primary care quality on inpatient length of stay for people with dementia: an analysis by discharge destination. In: CHE Research Paper No. 113, Centre for Health Economics, University of York, York. 2015.
- Terroso M, Rosa N, Torres Marques A, Simoes R. Physical consequences of falls in the elderly: a literature review from 1995 to 2010. *Eur Rev Aging Phys Activ*. 2014;11:51–9.
- Morello RT, Barker AL, Watts JJ, Haines T, Zavarsek SS, Hill KD, et al. The extra resource burden of in-hospital falls: a cost of falls study. *Med J Aust*. 2015;203, 367.e1–e8.
- Cox C, Vassallo M. Fear of falling assessments in older people with dementia. *Rev Clin Gerontol*. 2015;25:98–106.
- Peek K, Bryant J, Carey M, Dodd N, Freund M, Lawson S, et al. Reducing falls among people living with dementia: a systematic review. *Dementia*. 2020;19:1621–40.
- Theisen S, Drabik A, Stock S. Pressure ulcers in older hospitalised patients and its impact on length of stay: a retrospective observational study. *J Clin Nurs*. 2012;21:380–7.
- Dealey C, Posnett J, Walker A. The cost of pressure ulcers in the United Kingdom. *J Wound Care*. 2012;21:261–4.
- Gorecki C, Closs SJ, Nixon J, Briggs M. Patient-reported pressure ulcer pain: a mixed-methods systematic review. *J Pain Symptom Manage*. 2011;42:443–59.
- Jaul E, Meiron O, Menczel J. The effect of pressure ulcers on the survival in patients with advanced dementia and comorbidities. *Exp Aging Res*. 2016;42.