



Revista Iberoamericana de Micología

www.elsevier.es/reviberoammicol



Original article

Molecular phylogenetic diversity of the emerging mucoralean fungus *Apophysomyces*: Proposal of three new species

Eduardo Alvarez^a, Alberto M. Stchigel^a, Josep Cano^{a,*}, Deanna A. Sutton^b, Annette W. Fothergill^b, Jagdish Chander^c, Valentina Salas^a, Michael G. Rinaldi^b and Josep Guarro^a

^a Mycology Unit, Medical School and IISPV, Universitat Rovira i Virgili, Reus, Spain

^b Fungus Testing Laboratory, University of Texas Health Science Center, San Antonio, TX, USA

^c Government Medical College and Hospital, Chandigarh, India

ARTICLE INFO

Article history:

Received 13 November 2009

Accepted 12 January 2010

Available online 4 de marzo de 2010

Keywords:

Apophysomyces elegans

Apophysomyces ossiformis

Apophysomyces trapeziformis

Apophysomyces variabilis

Phylogeny

Taxonomy

Zygomycetes

ABSTRACT

Background: *Apophysomyces* is a monotypic genus belonging to the order Mucorales. The species *Apophysomyces elegans* has been reported to cause severe infections in immunocompromised and immunocompetent people. In a previous study of Álvarez et al.³ [J Clin Microbiol 2009;47:1650–6], we demonstrated a high variability among the 5.8S rRNA gene sequences of clinical strains of *A. elegans*.

Material and methods: We performed a polyphasic study based on the analysis of the sequences of the histone 3 gene, the internal transcribed spacer region of the rDNA gene, and domains D1 and D2 of the 28S rRNA gene, as well as by evaluation of some relevant morphological and physiological characteristics of a set of clinical and environmental strains of *A. elegans*.

Results and conclusions: We have demonstrated that *A. elegans* is a complex of species. We propose as new species *Apophysomyces ossiformis*, characterised by bone-shaped sporangiospores, *Apophysomyces trapeziformis*, with trapezoid-shaped sporangiospores, and *Apophysomyces variabilis*, with variable-shaped sporangiospores. These species failed to assimilate esculin, whereas *A. elegans* was able to assimilate that glycoside. Amphotericin B and posaconazole are the most active *in vitro* drugs against *Apophysomyces*.

© 2009 Revista Iberoamericana de Micología. Published by Elsevier España, S.L. All rights reserved.

Diversidad filogenética del hongo mucoral emergente *Apophysomyces*: propuesta de tres nuevas especies

RESUMEN

Palabras clave:

Apophysomyces elegans

Apophysomyces ossiformis

Apophysomyces trapeziformis

Apophysomyces variabilis

Filogenia

Taxonomía

Zigomicetos

Antecedentes: *Apophysomyces* es un género monoespecífico perteneciente al orden Mucorales. La especie *Apophysomyces elegans*, ha sido reportada como causante de infecciones severas en pacientes inmunocomprometidos e inmunocompetentes. En un estudio previo (Álvarez et al., J Clin Microbiol. 2009;47:1650–6), demostramos la elevada variabilidad dentro de las secuencias del gen 5.8S del ARNr en un grupo de cepas clínicas de *A. elegans*.

Material y métodos: Hemos realizado un estudio polifásico basado en el análisis de las secuencias del gen de la histona 3, la región de los espaciadores internos del ADNr y los dominios D1 y D2 del gen 28S del ARNr, así como la evaluación de caracteres morfológicos y fisiológicos relevantes de un grupo de cepas clínicas y ambientales de *A. elegans*.

Resultados y conclusiones: Hemos demostrado que *A. elegans* es un complejo de especies. Proponemos como nuevas especies para la ciencia *Apophysomyces ossiformis*, caracterizada por sus esporangiosporas con forma de hueso; *Apophysomyces trapeziformis*, con esporangiosporas trapezoidales; y *Apophysomyces variabilis*, con esporangiosporas de formas variables. Las nuevas especies no asimilan la esculina, en tanto que *A. elegans* fue capaz de asimilar dicho glicósido. La anfotericina B y el posaconazol fueron los antifúngicos más activos frente a *Apophysomyces*.

© 2009 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

The genus *Apophysomyces*, belonging to the subphylum Mucoromycotina (oldest phylum Zygomycota),¹¹ was erected by Misra et al.¹⁸ in 1979 to accommodate the only species of the genus,

Apophysomyces elegans, which was isolated from soil samples in northern India. This fungus was characterized by pyriform sporangia, conspicuous funnel- and/or bell-shaped apophyses, and subhyaline, thin-, and smooth-walled sporangiospores that are mostly oblong with rounded ends. It is a thermotolerant fungus that grows rapidly between 26 and 42 °C.^{6,18} *A. elegans* is not only isolated from soil, decaying vegetation, and as an environmental

* Corresponding author.

E-mail address: josep.cano@urv.cat (J. Cano).

contaminant,^{17,18,24} but it is also able to cause severe human infections.⁸ Unlike other members of Mucorales, this fungus primarily infects immunocompetent hosts.¹⁷ The infection typically follows traumatic implantation of the agent, but may also result from inhalation of spores into the sinus.^{5,6,12,15,17,21} This disease is more common in tropical and subtropical climates. Cases have been reported in Australia,^{6,20} India,^{5,13,14,26} the United States,^{3,15} Sri Lanka,⁴ Thailand,²⁴ and in Central and South America.^{21,22}

The genetic population structure of *A. elegans* remains largely unknown and may be due in part to the lack of preservation of strains for study. In a recent survey on the spectrum of species of Mucorales from clinical sources in the United States, we demonstrated a high intraspecific 5.8S rRNA gene sequence diversity in *A. elegans*.³ Additionally, in a typing study of *A. elegans* using microsatellites markers, it was demonstrated that, in a set of clinical strains, mainly from India, different banding patterns exist.⁵ These data suggest that more than one phylogenetic species may be present within the morphospecies *A. elegans*.

To determine possible cryptic species in *A. elegans*, we performed a polyphasic study on a diverse panel of strains, based on a multilocus sequence analysis of three loci (the histone 3 gene (H3), internal transcribed spacer region of the rDNA (ITS), and domains D1 and D2 of the 28S rRNA gene) and the evaluation of different morphological and physiological characters.

Materials and methods

Fungal strains

A total of 16 strains from different origins were included in the study (Table 1). The fungi were cultured on potato dextrose agar (PDA, Pronadisa, Madrid, Spain) and incubated at $35 \pm 1^\circ\text{C}$ for 2–5 days.

DNA extraction, amplification, and sequencing

DNA was extracted and purified directly from fungal colonies following a slightly modified Fast DNA kit protocol (Bio101, Vista,

CA, USA), consisting of a homogenization step repeated three times with a FastPrep FP120 instrument (Thermo Savant, Holbrook, NY, USA). DNA was quantified by the GeneQuant pro (Amersham Pharmacia Biotech, Cambridge, England). The internal transcribed spacer (ITS) region of the nuclear rDNA was amplified with the primer pair ITS5 and ITS4, the D1–D2 domains of the 28S rRNA gene were amplified with the primer pair NL1–NL4, and histone 3 (H3) gene was amplified with the primer pair H3-1a–H3-1b.¹⁰

The PCR mix (25 μl) included 10 mM Tris–HCl (pH 8.3), 50 mM KCl, and 2.5 mM MgCl_2 (10 \times Perkin-Elmer buffer II plus MgCl_2 solution Roche Molecular Systems, Branchburg, NJ, USA), 100 μM of each dNTP (Promega, Madison, WI, USA), 1 μM of each primer, and 1.5 U of AmpliTaq DNA polymerase (Roche). The amplification program for the three DNA fragments included an initial denaturation at 94°C for 5 min, followed by 35 cycles of denaturation at 95°C for 30 s, annealing for 1 min at 54°C , and extension for 1 min at 72°C . The products were purified with an Illustra GFXTM PCR DNA and Gel Band Purification Kit (General Electric Healthcare, Buckinghamshire, UK) and stored at -20°C until they were used in sequencing. PCR products were sequenced using the same primers employed for amplification and following the Taq DyeDeoxy Terminator Cycle Sequencing Kit protocol (Applied Biosystems, Gouda, Netherlands). Reactions were run on a 310 DNA sequencer (Applied Biosystems). Consensus sequences were obtained using the Autoassembler program (Perkin-Elmer-Applied Biosystems) and Seqman software (Laser-gene, Madison, WI).

Phylogenetic analyses

The sequences were aligned using Clustal X (version 1.8) computer program, followed by manual adjustments with a text editor. Most-parsimonious tree (MPT) analyses were performed using PAUP* version 4.0b10. One hundred heuristic searches were conducted with random sequence addition and tree bisection–reconnection branch-swapping algorithms, collapsing zero-length branches, and saving all minimal-length trees (MulTrees). *Saksenaia vasiformis* (FMR 10131) was chosen as the outgroup. Support for internal branches was assessed using a heuristic parsimony search of 1000 bootstrapped data sets. The combined data set of the ITS, D1–D2, and H3 was tested for incongruence with the

Table 1

Origin of *Apophysomyces* strains included in the study

Isolate	Source	GenBank accession no.		
		ITS1–5.8S–ITS2	D1/D2 domains of 28S rDNA	H3
CBS ^a 476.78	Soil, Deoria, India	FN556440	FN554249	FN555155
CBS 477.78	Soil, Gorakhpur, India	FN556437	FN554250	FN555154
CBS 658.93	Osteomyelitis, Netherlands Antilles	FN556436	FN554258	FN555161
GMCH ^b 480/07	Cutaneous infection, India	FN556442	FN554253	FN555163
GMCH 211/09	Cutaneous infection, India	FN556443	FN554254	FN555164
IMI ^c 338332	Ankle aspirate, Australia	FN556438	FN554257	FN555159
IMI 338333	Daly river, Australia	FN556439	FN554256	FN555160
UTHSC ^d 03-3644	Dolphin, Florida, USA	FN556431	FN554259	FN555158
UTHSC 04-838	Cellulitis wound leg, Minnesota, USA	FN556432	FN554252	FN555157
UTHSC 04-891	Sinus, Minnesota, USA	FN556433	FN554264	FN555165
UTHSC 06-2356	Dolphin, Texas, USA	FN556427	FN554262	FN555167
UTHSC 06-4222	Dolphin, Bahamas	FN556428	FN554255	FN555162
UTHSC 07-204	Facial cellulitis, Arizona, USA	FN556435	FN554251	FN555156
UTHSC 08-1425	Abdominal tissue, Phoenix, USA	FN556429	FN554261	FN555168
UTHSC 08-2146	Skin biopsy, Colorado, USA	FN556430	FN554260	FN555169
UTHSC R-3841	Necrotic face tissue, Georgia, USA	FN556434	FN554263	FN555166

^a CBS, Centraalbureau voor Schimmelcultures, Utrecht, Netherlands.

^b GMCH, Government Medical College and Hospital, Chandigarh, India.

^c IMI, International Mycological Institute, CABI-Bioscience, Egham, UK.

^d UTHSC, Fungus Testing Laboratory, University of Texas Health Science Center, San Antonio, Texas, USA.

partition homogeneity test (PHT) as implemented in PAUP*. The Kishino–Hasegawa test was performed to determine whether trees differed significantly. Gaps were treated as missing data.

Morphological studies

The strains were subcultured on PDA, Czapek agar (CZA; Difco, Becton Dickinson, France), malt extract agar (MEA; 10 g of malt extract, 20 g of agar, and 1000 ml of distilled water), and starch agar (SA; 5 g of soluble starch, 15 g of agar, and 1000 ml of distilled water), and incubated at 37 and 42 °C. The microscopic features were determined on the sixth day in wet mounts on water and on lactic acid, which were examined under a light microscope. The strains were identified using schemes based on morphological characters.^{8,18}

Physiological studies

Growth rates at 4, 15, 24, 30, 35, 37, 42, and 50 °C were determined on PDA, MEA, CZA, and SA for each of the strains included in the study. The Petri dishes were inoculated in the center, incubated in darkness, and the colony diameters (in millimeters) were measured daily.

Carbon source assimilation profiles were determined with the commercial kit API 50CH (bioMérieux, Marcy, l'Etoile, France), following protocols described previously but with minor modifications.²³ To obtain sufficient sporulation, all strains were cultured for 6 days on CZA at 42 °C. A final concentration of 5×10^5 CFU/ml was prepared in 20 ml of yeast nitrogen base (7.7 g/l; Difco), containing 0.5 g/l L-chloramphenicol (Sigma-Aldrich) and 0.1% Bacto agar (Difco), and each well of the strips was inoculated with 300 µl of medium. The viability of the conidia was verified by plating 100 µl of serial dilutions of each inoculum onto PDA and incubating at 42 °C for 6 days. The inoculated API 50 CH strips were incubated for 48–72 h at 37 °C in darkness. After incubation, the strips were read visually and growth or lack of growth was noted. Weak growth was considered as a positive result. For nitrogen source assimilations, we used the same inoculum described above, but the yeast nitrogen base broth was replaced by carbon nitrogen base broth (Difco), and testing was performed in sterile, disposable, multiwell microplates. The medium with the nitrogen sources was dispensed into the wells in 150 µl volumes with a multichannel pipette and each well was inoculated with 50 µl of the conidial suspension. The microplates were incubated at 37 °C in darkness for 48 and 72 h. We also determined growth of the strains on NaCl (2%, 5%, 7%, 10%), MgCl₂ 2%, and cycloheximide 0.1%.^{9,27} All tests were performed in duplicate.

The production of urease was determined after incubation on Christensen's urea agar slants at 37 °C for 8 days.¹⁶

Mating tests

Sixteen *Apophysomyces* strains were grown on CZA plates at 37 °C in the dark, and then paired in all combinations, including self-crosses, on CZA. Each strain was streaked onto one half of a CZA plate opposite to the streak of another strain, allowing for a central zone of contact as the strains grew. Plates were incubated at 37 °C and examined macroscopically each week for up to 6 months for the presence of zygosporangia. All tests were performed in duplicate.

Antifungal susceptibility testing

The *in vitro* activity of seven antifungal agents against the 16 strains of *Apophysomyces* was evaluated according to Clinical and

Laboratory Standards Institute guidelines (M38-A2).¹⁹ The drugs tested were amphotericin B (USP, Rockville, MD, USA), anidulafungin (Pfizer Inc., New York, NY, USA), caspofungin (Merck & Co., Inc., Rahway, NJ, USA), itraconazole (Jansen Pharmaceutica, Beerse, Belgium), posaconazole (Schering-Plough Ltd., Hertfordshire UK), ravuconazole (Bristol-Myers Squibb Company, New Brunswick, NJ, USA), and voriconazole (Pfizer Inc., New York, NY, USA).

Nucleotide sequence accession numbers

All the sequences obtained in this study were deposited in GenBank database and assigned the accession numbers listed in Table 1.

Results

Phylogeny

With the primers used, we were able to amplify and sequence 684–820, 582–683, and 345–382 bp of the ITS, D1–D2, and H3 loci, respectively. Of the 1630 nucleotides sequenced, 48 characters were parsimony informative in the different strains, the lowest number was eight for the H3 gene, and the highest was 27 for the ITS region. Sequences of the three region genes were analyzed phylogenetically as separate (data not shown) and combined datasets. The partition homogeneity test demonstrated that the three loci sequence data sets were congruent ($P=0.05$) and could therefore be combined. A total of 36 MPT was produced from a heuristic search, using the combined dataset from the three loci (Fig. 1). The trees had a consistency index of 0.969, a retention index of 0.956, and a homoplasy index of 0.031. Clustering was similar to that observed in the particular trees of the different genes analyzed. Most nodes in the combined analysis showed increased clade support as measured by bootstrapping (six nodes with $\geq 70\%$). Analyses of the combined partitions support the recognition of four well supported clades (Fig. 1), each of which could be considered separate phylogenetic species. Clade 1 (bootstrap support (bs) 73%) was composed of two strains from India, two from Australia, and one strain each from the United States, Netherlands Antilles, and Bahamas. Within clade 2 (bs 100%) were included the type strain of *A. elegans* and one strain from Indian soil. In clade 3 five strains (bs 96%) were located, four of them of clinical origin from the United States, and one from a dolphin. Finally, clade 4 (bs 98%) consisted of two clinical strains, both from the United States.

Physiology

Carbon assimilation profiles of the different strains on API 50CH strips are shown in Table 2. Assimilation patterns of all the strains were positive for 20 tests. Twenty-seven carbon sources were not assimilated by any strain. The profiles of assimilation of two carbon sources, esculin and D-lyxose, were species- and strain-dependent, respectively. Esculin was weakly assimilated only by the strains nested in clade 2. The assimilation of D-lyxose was highly variable among the strains of the different clades. By contrast, the variability in the assimilations of nitrogen sources, and tolerance to NaCl, MgCl₂, and cycloheximide was nule among the species (Table 3). All the strains were positive for 11 nitrogen sources. Nitrite was not assimilated by any of the strains. All strains were able to grow at 2% NaCl and at 2% MgCl₂, but failed to grow at 5% NaCl and at 0.1% cycloheximide.




Fig. 1. One of the 36 most parsimonious trees obtained from heuristic searches based on analysis produced from the combined data set. Sequence of *Saksenaea vasiformis* was chosen as outgroup. Bootstrap support values above 70% are indicated at the nodes. T: Type strain.

Morphology

In general, all the strains examined displayed the typical features of the genus *Apophysomyces* described by Misra et al.¹⁸ However, a more detailed microscopic study of these fungi showed important and consistent differences, mainly in the morphology of sporangiophores and sporangiospores, which correlated with the different phylogenetic species. The strains included in clade 1 showed some morphological diversity. The sporangiospores ranged from broadly clavate to ellipsoidal, were flattened on one side, and measured $5\text{--}14 \times 3\text{--}6 \mu\text{m}$. The strains included in clade 2, which comprises the type strain of *A. elegans*, showed ovoid, subspherical, broadly ellipsoidal to barrel shaped sporangiospores, although more irregularly shaped spores were also present, measured $6\text{--}12 \times 5\text{--}8 \mu\text{m}$, and were the largest for the different species in the complex. The sporangiospores of the strains included in clade 3 were trapezoidal and smaller ($5\text{--}8.5 \times 3\text{--}5 \mu\text{m}$) while those of strains in clade 4 were thick-walled and clearly biconcave (bone-shaped) in side view, measuring $6\text{--}8 \times 3\text{--}5.5 \mu\text{m}$. In addition to differences in spore morphology, the strains in clade 2 also showed two types of sporangiophores: (i) large (up to $540 \mu\text{m}$), bearing vase- or bell-shaped apophyses and (ii) shorter (up to $400 \mu\text{m}$), bearing funnel-shaped apophyses. The sporangiophores in strains of clades 1, 3, and 4 are similar to the short ones of clade 2.

Mating test

Zygospore formation was not observed after 6 months of incubation in all the mating tests assayed.

Based on the described morphological and physiological differences, which correlated with the molecular data, we concluded that clades 1, 3, and 4 represent three species of *Apophysomyces*, different from *A. elegans* (clade 2), which are proposed here as new species.

Apophysomyces variabilis Alvarez, Stchigel, Cano, D.A. Sutton et Guarro, sp. nov. (Figs. 2A, B; 3H, I).

Coloniae in CZA ad 37°C rapide crescentes, albae, sed sparsis, inmersis pro parte maxima compositae. Sporangiophora erecta, plerumque simplicia, $100\text{--}400 \mu\text{m}$ longa, $2\text{--}3.5 \mu\text{m}$ lata, brunnea, cum sporangio apophysati. Apophyses plerumque infundibuliformes, $15\text{--}20 \times 15\text{--}20 \mu\text{m}^2$. Sporangiosporae variabiles in forma et magnitudine, trapezoides, ellipsoideae, subtriangulares vel claviformis, $5\text{--}14 \times 3\text{--}6 \mu\text{m}^2$. Holotypus, CBS H-658.93, ex osteomyelitis (cultura viva FMR 10381, CBS 658.93).

Etymology: the epithet refers to the variable morphology of the sporangiospores.

Table 2
Carbon assimilation profiles for *Apophysomyces* species obtained with API 50 CH strips

[illegible]

AMD (amidon)	+	+	+	+	+	+	—	+	+	+	+	+	+	+	+	+
GLYG (glycogen)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
XLT (xylitol)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
GEN (gentiobiose)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
TUR (D-turanose)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
LYX (D-lyxose)	—	—	—	—	+	+	—	—	—	—	+	—	—	—	+	+
TAG (D-tagatose)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
DFUC (D-fucose)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
LFUC (L-fucose)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
DARL (D-arabitol)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
LARL (L-arabitol)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
GNT (potassium gluconate)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
2KG (potassium 2-keto-gluconate)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
5KG (potassium 5-keto-gluconate)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

Table 3
Nitrogen assimilation and tolerance to chemical compounds for the *Apophysomyces* species included in this study

Nitrogen source and other tests	Species															
	<i>A. variabilis</i>							<i>A. elegans</i>		<i>A. trapeziformis</i>				<i>A. ossiformis</i>		
	CBS 658.93	UTHSC 06-4222	GMCH 211/09	GMCH 480/07	IMI 338332	IMI 338333	UTHSC 03-3644	CBS 476.78	CBS 477.78	UTHSC 04-891	UTHSC 06-2356	UTHSC 08-1425	UTHSC 08-2146	UTHSC R-3841	UTHSC 04-838	UTHSC 07-204
Cadaverine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Creatine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Creatinine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
L-lysine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Nitrate	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Nitrite	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
L-tryptophan	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
L-proline	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
L-leucine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
L-ornithine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
L-cysteine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Arginine	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
2% NaCl	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
5% NaCl	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
7% NaCl	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10% NaCl	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
2% MgCl ₂	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Cycloheximide 0.1%	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—




Fig. 2. Morphology of the apophyses and sporangiospores of *Apophysomyces*: (A, B) *A. variabilis* CBS 658.93 (A, apophyses funnel-shaped; B, sporangiospores). (C, D) *A. ossiformis* UTHSC 04-838 (C, apophyses funnel-shaped; D, bone-like shaped sporangiospores). (E) *A. trapeziformis* UTHSC 08-1425 (E, trapezoid-shaped sporangiospores). (F, G) *A. elegans* CBS 476.78 (F, apophyses bell- and funnel-shaped; (G) sub-spherical to broadly ellipsoidal sporangiospores). All bars = 10 µm.

Colonies attaining a diameter of 90 mm after 4 days of incubation at 37 °C on CZA, whitish, with scarce aerial mycelium; hyphae branched, hyaline, smooth-walled, 3–5.5 µm in diameter; reverse concolorous. Sporangioophores erect, generally arising singly, at first hyaline, soon becoming light greyish brown, generally straight, slightly tapered towards the apex, unbranched, 100–400 µm long, 2–3.5 µm wide, and smooth-walled. Sporangia apophysate, terminal, pyriform, multisporous, white at first, becoming light greyish brown when mature, and 15–50 µm in diameter. Apophyses short, funnel-shaped, and 15–20 ×

15–20 µm. Sporangiospores variable in shape, trapezoid, ellipsoid, subtriangular or claviform, hyaline to light brown in mass, smooth- and thin-walled, and 5–14 × 3–6 µm. Not able to assimilate esculin.

Colonies on SA, PDA, and MEA showed similar features than on CZA, but they were more floccose, white, and with less sporulation. The optimum growth temperature was 35–42 °C and the minimum temperature of growth was 15 °C. The fungus did not grow at 50 °C.

Apophysomyces ossiformis Alvarez, Stchigel, Cano, D.A. Sutton et Guarro sp. nov. (Figs. 2C, D; 3D, E).

Coloniae in CZA ad 37 °C rapide crescentes, albae, sed sparsis, immersis pro parte maxima compositae. Sporangiophora erecta, plerumque simplicia, 100–400 μm longa, 2–3.5 μm lata, brunnea, cum sporangio apophysati. Apophyses plerumque infundibuliformes, 15–20 \times 15–20 μm . Sporangiosporae ossiformis, 6–8 \times 3–5.5 μm . Holotypus, CBS H-20328, ex cellulitis cruris vulnus hominis (cultura viva FMR 9913, UTHSC 04-838).

Etymology: the epithet refers to the bone-like shape of the sporangiospores.

Colonies attaining a diameter of 90 mm after 4 days of incubation at 37 °C on CZA, whitish, with scarce aerial mycelium, branched, hyaline, smooth-walled, and 3–5.5 μm in diameter; reverse concolorous. Sporangiophores erect, generally arising singly, at first hyaline soon becoming light greyish brown, generally straight, slightly tapered towards the apex, unbranched, 100–400 μm long, 2–3.5 μm wide, and smooth-walled. Sporangia apophysate, terminal, pyriform, multisporous, white at first,




Fig. 3. (A)–(C) *Apophysomyces elegans* (A, vessel-shaped sporangiophore; B, funnel-shaped sporangiophore; C, sporangiospores). (D, E) *Apophysomyces ossiformis* (D, sporangiophore; E, sporangiospore, frontal and side views). (F, G) *Apophysomyces trapeziformis* (F, sporangiophore; G, sporangiospore). (H, I) *Apophysomyces variabilis* (H, sporangiophore; I, sporangiospores). Bars: 10 μm .

becoming light greyish brown when mature, 15–50 µm in diameter. Apophyses short, funnel-shaped, 15–20 × 15–20 µm. Sporangiospores mostly bone-like shaped, hyaline to light brown in mass, smooth- and thick-walled, and 6–8 × 3–5.5 µm. Not able to assimilate esculin.

Colonies on SA, PDA, and MEA showed similar features than on CZA, but they were more floccose, white, and with less sporulation. The optimum growth temperature was 35–42 °C and the minimum 15 °C. The fungus did not grow at 50 °C.

Apophysomyces trapeziformis Alvarez, Cano, Stchigel, D.A. Sutton et Guarro sp. nov. (Figs. 2E; 3F, G).

Coloniae in CZA ad 37 °C rapide crescentes, albae, sed sparsis, immersis pro parte maxima compositae. Sporangio-phora erecta, plerumque simplicia, usque ad 400 µm longa, 2–3.5 µm lata, brunnei, cum sporangio apophysati. Apophyses plerumque infundibuliformes, 15–20 × 15–20 µm. Sporangiosporae trapezoids vel ellipsoideae, 5–8.5 × 3–5 µm.

Holotypus, CBS H-20329, ex abscessus abdominis humanus (cultura viva FMR 10456, UTHSC 08-1425).

Etymology: the epithet refers to the trapezoid shape of the sporangiospores in side view.

Colonies attaining a diameter of 90 mm after 4 days of incubation at 37 °C on CZA, whitish, with scarce aerial mycelium, branched, hyaline, smooth-walled, 3–5.5 µm in diameter; reverse concolorous. Sporangio-phores erect, generally arising singly, at first hyaline soon becoming light greyish brown, generally straight, slightly tapered towards the apex, unbranched, up to 400 µm long, 2–3.5 µm wide, and smooth-walled. Sporangia apophysate, terminal, pyriform, multispored, white at first, becoming light greyish brown when mature, and 15–50 µm in diameter. Apophyses short, funnel-shaped, and 15–20 × 15–20 µm. Sporangiospores mostly trapezoid-shaped in side view, more or less cylindrical in front view, flattened at one side and broadly convex on opposite side, hyaline to light brown in mass, smooth- and thin-walled, and 5–8.5 × 3–5 µm. Not able to assimilate esculin.

Colonies on SA, PDA, and MEA showed similar features than on CZA, but they were more floccose, white, and with less sporulation. The optimum growth temperature was 35–42 °C and the minimum 15 °C. The fungus did not grow at 50 °C.

Based on our morphologic and physiologic studies, the type species of *Apophysomyces* is redefined as follows:

A. elegans Misra, Srivastava, and Lata (Figs. 2F,G; 3A–C).

Colonies attaining a diameter of 90 mm after 4 days at 37 °C on CZA, whitish at first, becoming brownish grey, with scarce aerial mycelium; reverse concolorous. Sporangio-phores generally arising singly, emerging from aerial hyphae, straight or curved, mainly unbranched or some times branched at the apex, light greyish brown, with two types of morphology, i.e. (i) those that were large (up to 540 µm), bearing vase- or bell-shaped apophyses (15–46 × 11–40 µm) and (ii) those that were shorter (up to 400 µm) and bore funnel-shaped apophyses (15–20 × 15–20 µm), of 4–7.5 µm wide, and smooth-walled. Sporangia produced terminally, pyriform, with distinct apophyses, and 20–60 µm in diameter. Sporangiospores ovoid, subspherical, broadly ellipsoidal to barrel-shaped, frequently irregularly shaped, sub-hyaline, smooth- and thin-walled, and 6–12 × 5–8 µm. The strains analyzed were able to assimilate esculin.

Similar colonies features as described on CZA were observed on AS, PDA, and MEA, with the exception of lesser production of mycelium in CZA. The optimum growth temperature was 35–42 °C and the minimum 15 °C. The fungus did not grow at 50 °C.

Table 4

In vitro antifungal susceptibility data for *Apophysomyces* species

Species (# of strains tested)	Antifungal agent	MIC or MEC (µg/mL) 24 h	
		GM	Range
<i>A. variabilis</i> (7)	Amphotericin B	1.0	1
	Posaconazole	1.1	1–2
	Voriconazole	29.0	16– > 16
	Itraconazole	1.0	0.5–2
	Ravuconazole	1.0	0.5–2
	Anidulafungin	7.2	4–8
	Caspofungin	19.5	4– > 16
<i>A. elegans</i> (2)	Amphotericin B	0.5	0.5
	Posaconazole	0.5	0.5
	Voriconazole	8.0	8
	Itraconazole	1.0	0.5–2
	Ravuconazole	1.0	0.5–2
	Anidulafungin	8.0	8
	Caspofungin	32.0	> 16
<i>A. trapeziformis</i> (5)	Amphotericin B	0.8	0.5–1
	Posaconazole	0.8	0.5–1
	Voriconazole	16.0	8– > 16
	Itraconazole	0.9	0.5–2
	Ravuconazole	0.9	0.5–2
	Anidulafungin	6.1	4–8
	Caspofungin	27.9	16– > 16
<i>A. ossiformis</i> (2)	Amphotericin B	1.4	1–2
	Posaconazole	0.7	0.5–1
	Voriconazole	5.7	4–8
	Itraconazole	1.4	1–2
	Ravuconazole	2.8	2–4
	Anidulafungin	5.7	4–8
	Caspofungin	32.0	< 16

Antifungal susceptibility tests

The results of antifungal susceptibility testing for *Apophysomyces* strains are shown in Table 4. Amphotericin B and posaconazole were the most active antifungal agents. Itraconazole and ravuconazole were more active than voriconazole, and caspofungin and anidulafungin were inactive against all strains.

Discussion

Apophysomyces has been traditionally considered a monotypic genus. However, on the basis of genetic, physiological, and morphological data, we have demonstrated here that the genus constitutes a complex of species. DNA sequences from three different loci were analyzed to infer phylogenetic relationships and species boundaries within strains morphologically identified as *A. elegans*. The informations provided by the three loci evaluated were similar, and proved to be useful markers for species level differentiation in *Apophysomyces*. Although our study included strains from very diverse origins, the number of isolates we could obtain was small, and we anticipate even greater diversity as more strains become available. Given this limitation we were, however, able to recognize at least four phylogenetically, morphologically, and physiologically different species. The shape and size of the sporangiospores, the type of the sporangio-phore, and the shape of the apophyses were the most useful characters for this purpose.

As carbon assimilation profiles can be useful for differentiation of human pathogenic mucoralean genera,²³ we tested the assimilation of numerous carbon sources (Table 2). Low interspecific variability within *Apophysomyces* was noted, with the only exception of esculin assimilation, which was positive for *A. elegans*, and negative for the

other species in the complex. In this study, *Apophysomyces* strains also showed negative results for D-galactose, amygdalyn, arbutin, salicin, and gentiobiose assimilation, while in the study of Schwarz et al.²³ these same tests were positive for the members of six other pathogenic genera, i.e. *Cunninghamella*, *Lichtheimia* (*Absidia*), *Mucor*, *Rhizopus*, *Rhizomucor*, and *Syncephalastrum*. In contrast, carbon sources such as L-sorbose, L-rhamnose, dulcitol, inositol, erythritol, D-arabinose, methyl- β -D-xylopyranoside, methyl-D-mannopyranoside, methyl-D-glucopyranoside, D-tagatose, D-fucose, L-fucose, and inulin were all negative for both the *Apophysomyces* strains and the other six mentioned genera.²³ Other carbon sources such as glycerol were assimilated by *Apophysomyces*, *Rhizopus*, and *Cunninghamella* but not by *Lichtheimia*, *Rhizomucor*, *Mucor*, and *Syncephalastrum*; D-ribose was assimilated by *Apophysomyces*, *Rhizopus*, and *Mucor* and not by *Cunninghamella*, *Lichtheimia*, *Rhizomucor*, and *Syncephalastrum*; L-xylose was assimilated only by *Mucor* and some strains of *Rhizopus*, and D-lactose was assimilated only by *Lichtheimia*, *Rhizomucor*, and *Syncephalastrum*, although it was species dependent in *Rhizomucor*. The nitrogen assimilation profiles and tolerance to various chemical agents for the *Apophysomyces* strains tested in this study were non-discriminatory (Table 3).

From a clinical point of view it is also worth mentioning that none of the clinical strains included in this study belonged to clade 2, which contains the type strain, *A. elegans*, and which has previously been considered a pathogenic species. Clade 2 included only two environmental strains isolated from Indian soils.

The *in vitro* activity of the antifungal drugs tested appeared to corroborate data from previous studies.^{1,2,7,25}

Acknowledgments

This work was supported by the Spanish Ministerio de Ciencia y Tecnología Grants CGL 2007-65669/BOS and CGL 2009-08698/BOS.

References

- Alastruey-Izquierdo A, Castelli MV, Cuesta I, Monzón A, Cuenca-Estrella M, Rodríguez-Tudela JL. Activity of posaconazole and other antifungal agents against mucorales strains identified by sequencing of internal transcribed spacers. *Antimicrob Agents Chemother*. 2009;53:1686–9.
- Almyroudis NG, Sutton DA, Fothergill AW, Rinaldi MG, Kusne S. In vitro susceptibilities of 217 clinical isolates of zygomycetes to conventional and new antifungal agents. *Antimicrob Agents Chemother*. 2007;51:2587–90.
- Álvarez E, Sutton DA, Cano J, Fothergill AW, Stchigel A, Rinaldi MG, et al. Spectrum of zygomycete species identified in clinically significant specimens in the United States. *J Clin Microbiol*. 2009;47:1650–6.
- Andresen D, Donaldson A, Choo L, Knox A, Klaassen M, Ursic C, et al. Multifocal cutaneous mucormycosis complicating polymicrobial wound infections in a tsunami survivor from Sri Lanka. *Lancet*. 2005;365:876–8.
- Chakrabarti A, Ghosh A, Prasad GS, David JK, Gupta S, Das A, et al. *Apophysomyces elegans*: an emerging zygomycete in India. *J Clin Microbiol*. 2003;41:783–8.
- Cooter RD, Lim IS, Ellis DH, Leitch IOW. Burn wound zygomycosis caused by *Apophysomyces elegans*. *J Clin Microbiol*. 1990;28:2151–3.
- Dannaoui E, Meletiadis J, Mouton JW, Meis JF, Verweij PE. In vitro susceptibilities of zygomycetes to conventional and new antifungals. *J Antimicrob Chemother*. 2003;51:45–52.
- de Hoog GS, Guarro J, Gené J, Figueras MJ. Atlas of clinical fungi, 2nd ed. Centraalbureau voor Schimmelfcultures: Utrecht, The Netherlands; 2000.
- de Hoog GS, Marvin-Sikkema FD, Lahpoor GA, Gottschall JC, Prins RA, Guého E. Ecology and physiology of the emerging opportunistic fungi *Pseudallescheria boydii* and *Scedosporium prolificans*. *Mycoses*. 1994;37:71–8.
- Glass NL, Donaldson GC. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol*. 1995;61:1323–30.
- Hibbett DS, Binder M, Bischoff JF, Blackwell M, Cannon PF, Eriksson OE, et al. A higher level phylogenetic classification of the fungi. *Mycol Res*. 2007;111:509–47.
- Kimura M, Smith MB, McGinnis MR. Zygomycosis due to *Apophysomyces elegans*: report of 2 cases and review of the literature. *Arch Pathol Lab Med*. 1999;123:386–90.
- Kindo AJ, Shams NR, Kumar K, Kannan S, Vidya S, Kumar AR, et al. Fatal cellulitis caused by *Apophysomyces elegans*. *Indian J Med Microbiol*. 2007;25:285–7.
- Lakshmi V, Suda Rani T, Sharma S, Mohan VS, Sundaram C, Rao RR, et al. Zygomycotic necrotizing fasciitis caused by *Apophysomyces elegans*. *J Clin Microbiol*. 1993;31:1368–9.
- Liang KP, Tleyjeh IM, Wilson WR, Roberts GD, Temesgen Z. Rhino-orbitocerebral mucormycosis caused by *Apophysomyces elegans*. *J Clin Microbiol*. 2006;44:892–8.
- Marimon R, Gené J, Cano J, Trilles L, Dos Santos Lazéra M, Guarro J. Molecular phylogeny of *Sporothrix schenckii*. *J Clin Microbiol*. 2006;44:3251–6.
- Meis JF, Chakrabarti A. Changing epidemiology of an emerging infection: zygomycosis. *Clin Microbiol Infect*. 2009;15(Suppl. 5):10–4.
- Misra PC, Srivastava KJ, Lata K. *Apophysomyces*, a new genus of the Mucorales. *Mycotaxon*. 1979;8:377–82.
- National Committee for Clinical Laboratory Standards. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi: approved standard—second edition. Document M38-A2. National Committee for Clinical Laboratory Standards: Wayne, Pa; 2008.
- Page R, Gardam DJ, Heath CH. Severe cutaneous mucormycosis (zygomycosis) due to *Apophysomyces elegans*. *Aust N Z J Surg*. 2001;71:184–6.
- Ruiz CE, Arango M, Correa AL, López LS, Restrepo A. Necrotizing fasciitis in an immunocompetent patient caused by *Apophysomyces elegans*. *Biomedica*. 2004;24:239–51.
- Saravia-Flores M, Guaran DM, Argueta V. Invasive cutaneous infection caused by *Apophysomyces elegans* associated with a spider bite. *Mycoses*. 2010. doi:10.1111/j.1439-0507.2009.01698.x.
- Schwarz P, Lortholary O, Dromer F, Dannaoui E. Carbon assimilation profiles as a tool for identification of zygomycetes. *J Clin Microbiol*. 2007;45:1433–9.
- Snell BJ, Tavakoli K. Necrotizing fasciitis caused by *Apophysomyces elegans* complicating soft-tissue and pelvic injuries in a tsunami survivor from Thailand. *Plast Reconstr Surg*. 2007;119:448–9.
- Sun QN, Fothergill AW, McCarthy DI, Rinaldi MG, Graybill JR. In vitro activities of posaconazole, itraconazole, voriconazole, amphotericin B, and fluconazole against 37 clinical isolates of zygomycetes. *Antimicrob Agents Chemother*. 2002;46:1581–2.
- Suryanarayan Rao S, Panda NK, Pragache G, Chakrabarti A, Saravanan K. Sinoorbital mucormycosis due to *Apophysomyces elegans* in immunocompetent individuals—an increasing trend. *Am J Otolaryngol*. 2006;27:366–9.
- Yarrow D. Methods for the isolation, maintenance and identification of yeasts. In: Kurtzman CP, Fell JW, editors. *The yeasts, a taxonomic study*, 4th ed. Amsterdam, The Netherlands: Elsevier; 1998. p. 95–7.