



Note

Characterization, enzymatic activity and biofilm formation of *Candida* species isolated from goat milk

Carolina Segundo Zaragoza^{a,*}, Itzel López Ortiz^a, David Alejandro Contreras Caro del Castillo^b, Yesmín María Domínguez Hernández^a, Juan Antonio Rodríguez García^a

^a Laboratorio de Micología Veterinaria, Unidad de Servicios de Diagnóstico y Constatación, Centro de Enseñanza, Investigación y Extensión en Producción Animal en Altiplano, Facultad de Medicina Veterinaria y Zootecnia, UNAM, Mexico

^b Departamento de Medicina y Zootecnia de Rumiantes, Facultad de Medicina Veterinaria y Zootecnia, UNAM, Mexico

ARTICLE INFO

Article history:

Received 21 October 2020

Accepted 16 June 2021

Available online 10 September 2021

Keywords:

Yeast

Mastitis

Goat

Milk

Mexico

ABSTRACT

Background: Data regarding yeast microbiota in goat milk is scarce.

Aims: To isolate and identify species of the genus *Candida* in milk samples from clinically healthy goats, and evaluate their enzymatic activity and biofilm formation.

Methods: 1092 milk samples from clinically healthy goats were collected and processed. The yeast isolates were identified by phenotypic, methods and their enzymatic activity (phospholipase, hemolysin and protease) and biofilm formation evaluated.

Results: We obtained 221 *Candida* isolates belonging to six species: *Candida kefyr* (35.7%), *Candida guilliermondii* (33%), *Candida famata* (23.5%), *Candida glabrata* (5.9%), *Candida albicans* (1.35%) and *Candida parapsilosis sensu lato* (0.45%). Protease activity was detected in all *Candida* species while hemolysin activity was only present in *C. kefyr*, *C. guilliermondii*, *C. famata* and *C. albicans*. Only *C. albicans* showed phospholipase activity. With the exception of *C. parapsilosis sensu lato*, all *Candida* species formed biofilm, with 60.19% of the isolates being poor producers, 9.93% moderate producers, and 1.35% strong producers.

Conclusions: The milk of clinically healthy goats contains several species of the genus *Candida* that could play a role as opportunistic pathogens in mastitis.

© 2021 The Authors. Published by Elsevier España, S.L.U. on behalf of Asociación Española de Micología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Caracterización, actividad enzimática y formación de biopelículas de especies de *Candida* aisladas de leche de cabra

RESUMEN

Antecedentes: El conocimiento de la microbiota levaduriforme presente en la leche de cabra es escaso.

Objetivos: Aislar e identificar especies del género *Candida* en muestras de leche de cabras clínicamente sanas, evaluar su actividad enzimática y su capacidad de formar biopelículas.

Métodos: Se recogieron y procesaron 1092 muestras de leche de cabras clínicamente sanas. Las levaduras aisladas fueron identificadas mediante métodos fenotípicos, evaluándose posteriormente su actividad enzimática (producción de fosfolipasas, hemolisinas y proteasas) y la formación de biopelículas.

Resultados: Se obtuvieron 221 aislamientos de *Candida* de seis especies: *Candida kefyr* (35,7%), *Candida guilliermondii* (33%), *Candida famata* (23,5%), *Candida glabrata* (5,9%), *Candida albicans* (1,35%) y *Candida parapsilosis sensu lato* (0,45%). En todas las especies de *Candida* se detectó actividad proteolítica, y únicamente *C. kefyr*, *C. guilliermondii*, *C. famata* y *C. albicans* presentaron actividad hemolítica. Por su parte, *C. albicans* fue la única especie con actividad fosfolipasa. Con excepción de *C. parapsilosis sensu lato*, todas las especies de *Candida* formaron biopelícula, con el 60,19% de los aislamientos poco formadores de biopelícula, el 9,93% moderadamente formadores y el 1,35% altamente formadores.

Palabras clave:

Levadura

Mastitis

Cabra

Leche

México

* Corresponding author.

E-mail address: c.segund@yahoo.com.mx (C. Segundo Zaragoza).

Conclusiones: La leche de cabras clínicamente sanas presenta diversas especies del género *Candida* que podrían actuar como patógenos oportunistas de mastitis.

© 2021 Los Autores. Publicado por Elsevier España, S.L.U. en nombre de Asociación Española de Micología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The mammary glands in ruminants are known to harbor diverse bacteria; little is known, however, about their fungal microbiota. In dairy cattle, studies report the presence of different species of the genus *Candida*.^{7,18,21,25,34,37} A variety of enzymatic activities, mainly proteases, hemolysins and phospholipases, allow different species of *Candida* to adhere, colonize and invade host tissues.^{8,10} In addition, biofilm production enables *Candida* infection and resistance to antimicrobials.^{13,33} The objective of this study was to isolate and identify *Candida* species in milk samples from clinically healthy goats, and to evaluate their enzymatic activity and biofilm formation.

We collected 1092 milk samples during a six month-period from a farm with Alpina goats clinically healthy ($n = 100$) in Querétaro, Mexico. The samples were centrifuged (4000 rpm, 10 min), the pellets resuspended into 2 ml of yeast extract peptone dextrose (YEPD) broth (casein peptone 2%, dextrose 2%, yeast extract 1%), and incubated 48 h at 37 °C. Yeasts were isolated by plating 50 µl from each suspension on Sabouraud dextrose agar (SDA) supplemented with 50 mg/l chloramphenicol. Conventional phenotypic methods were used for yeast identification: Gram staining, germ tube production, pseudohyphae formation, sensitivity to 0.1% cycloheximide, film formation in liquid culture, urease production, carbohydrates assimilation and fermentation, and development in Biggy medium and CHROMagar *Candida*.^{2,5,6,14,24,27,28,30,32,40} Each yeast isolate was evaluated for enzymatic activity and biofilm production. Phospholipase and protease activity were performed according to Kantarcioglu and Yücel,²⁰ and hemolysin activity according to Luo et al.²⁶ Briefly, 6 mm diameter sterile filter discs were embedded with 10 µl yeast suspension (1×10^6 – 5×10^6 CFU/ml). These were placed onto SDA agar supplemented with 4% egg yolk, 0.2% bovine serum albumine or 7% sheep blood for phospholipase, protease and hemolysin activity evaluation, respectively. After incubation (37 °C, 7 days), the yeast colony and the corresponding clear halo diameters were measured. The enzymatic activity was determined through the enzyme activity index according to Williamson's formula.⁴¹ *Candida* biofilm formation was evaluated using the Gokce et al. safranin method.¹⁵ For each isolate a 1×10^8 CFU/ml yeast suspension was prepared in yeast nitrogen base medium (YNBG) with 8% glucose. From this stock suspension a 1:100 dilution was made, and 96-well polystyrene flat-bottom microplates were filled (200 µl/well) and incubated 48 h at 37 °C. Afterwards, the plates were washed three times with sterile phosphate-buffered saline (PBS, pH 7.4). The biofilms formed were fixed with methanol for 15 min (200 µl/well); methanol was discarded and the microplates were allowed to dry for 10 min at 37 °C. Thereafter, 1% safranin solution was added (200 µl/well), and after 20 min at room temperature a washing with PBS was done. Immediately after, 95% ethanol was added (200 µl/well, 20 min at room temperature). Finally, the optical density (OD) was determined at 490 nm using a Biotek ELISA reader. Categorization of biofilm production was established through OD intervals as described by Stepanovic et al.^{16,39} Descriptive statistics were used to analyze the results.

Candida was the solely yeast genus found in the goat milk samples analyzed. We obtained 221 *Candida* isolates, further identified

as *Candida kefyr* 35.74%, *Candida guilliermondii* 33%, *Candida famata* 23.53%, *Candida glabrata* 5.9%, *Candida albicans* 1.35%, and *Candida parapsilosis* (*sensu lato*)⁴⁴ 0.45% (Table 1). Concerning enzymatic activity, 98.65% of the isolates showed protease activity, 55.2% hemolytic activity, and 1.36% phospholipase activity. Protease activity varied among the species and within the isolates of a given species. *C. kefyr*, *C. guilliermondii*, and *C. famata* showed the whole range of protease activity, having more than 87% of them medium and high activity. All *C. glabrata* presented protease activity, circa 70% of them showing high activity. *C. albicans* and *C. parapsilosis* isolates exhibited only high protease activity. Hemolysin activity was observed in *C. albicans* (100%), *C. guilliermondii* (71.23%), *C. famata* (67.30%) and *C. kefyr* (40.50%). Phospholipase activity was just detected in *C. albicans* (Table 2). Regarding biofilm formation, four categories were determined: no biofilm formation, low biofilm formation, moderate biofilm formation, and high biofilm formation. Biofilm was produced in various degrees by 71.49% of all *Candida* isolates but *C. parapsilosis* (*sensu lato*) (Table 3).

Yeast identification was performed by conventional methods, which are the routine diagnostic methods used in our laboratory. Although known to be less sensitive and more time consuming than molecular methodologies, they are considered the gold *Candida* identification standard.^{1,43} Predominance of non-*C. albicans* *Candida* species in milk from clinically healthy goats, and from other ruminants with and without mastitis have been reported.^{9,17,19,21,22,25} Our findings support this since, out of the six *Candida* species found, *C. albicans* accounted only for 1.35% of the isolates. Little is known about the association of *Candida* spp. enzymatic production and biofilm formation with the yeasts virulence. Both proteases and hemolysins were detected in *C. albicans*, *C. famata*, *C. kefyr*, *C. guilliermondii* and *C. parapsilosis*, whereas phospholipases were only detected in *C. albicans*. These features are in agreement with previous studies.^{3,4,12,15,26,29,36} *In vitro* production of these virulence factors are strain-dependent and may differ according to the anatomical site infected or the yeasts involvement in a pathological process.^{4,33} Biofilm formation has been described in *C. albicans*, *C. tropicalis*, *C. parapsilosis* and *C. glabrata*, and correlation between yeast virulence and high biofilm production has been demonstrated in human beings.^{11,13,15,23,31,33,42} In animals, a research studying milk from buffaloes with mastitis have shown high biofilm production in *Candida zeylanoides*, *Candida rugosa* and *Candida kefyr*.³⁵ In this study, *C. albicans*, *C. kefyr*, *C. guilliermondii*, and *C. famata* isolates mostly showed a low production of biofilms. This might be related to the samples origin, clinically healthy animals, where formation of high density biofilms might not be crucial to yeasts persistence. Nevertheless, all these microorganisms could be opportunistic pathogens in mastitis.^{35,38}

More research on *Candida* presence, both in milk of clinically healthy and diseased goats, and their virulence factors is needed to determine their relationship with the development of fungal mastitis in this animal species. Our study is the first of its kind in Mexico, demonstrating the presence of different species of the genus *Candida* in goat milk from clinically healthy animals; the isolates exhibited various virulence factors under laboratory conditions.

Table 1
Identification of *Candida* isolates from the milk of clinically healthy goats.

Candida species	Pseudohyphae formation		Yeast development			Urease	Germ tube formation	Assimilation and fermentation																Yeast development 37 °C	
			SDA 37 °C	YNB + cycloheximide (0.1%)	Sabouraud broth			glu		sac		lac		mal		xyl		raf		tre		mel		Chromagar Candida	BIGGY
								A	F	A	F	A	F	A	F	A	F	A	F	A	F	A	F		
C. kefyr (n = 79)	+	V	+	–	–	–	–	+	+	+	+	V	V	–	–	V	–	+	+	–	–	–	–	Matte light pink	Light brown
C. guilliermondii (n = 73)	V	V	+	–	–	–	–	+	+	+	+	–	–	+	–	+	–	+	+	+	V	+	–	Bright pink	Reddish brown
C. famata (n = 52)	V	V	–	–	–	–	–	+	V	+	V	V	–	+	V	–	–	+	V	+	V	V	V	Matte pale pink	Reddish brown
C. glabrata (n = 13)	–	+	–	–	–	–	–	+	+	–	–	–	–	–	–	–	–	V	–	V	–	–	–	Matte pale pink	Light brown
C. albicans (n = 3)	+	+	+	–	–	–	+	+	+	V	–	–	–	+	+	+	–	–	–	V	V	–	–	Emerald green	Brown and silver appearance on the surface
C. parapsilosis (sensu lato) (n = 1)	+	+	–	–	–	–	–	+	+	+	–	–	–	+	–	+	–	–	–	+	–	–	–	Matte pale pink	Reddish brown

A: assimilation; F: fermentation; SDA: Sabouraud dextrose agar; YNB: yeast nitrogen base; glu: glucose; sac: saccharose; lac: lactose; mal: maltose; xyl: xylose; raf: raffinose; tre: trehalose; mel: melezitose; V: variable strains.

Table 2Enzymatic activity of *Candida* isolates from the milk of clinically healthy goats.

Candida species	Number of isolates (n)	Scores of enzymatic activities (Pz)											
		Protease				Phospholipase				Hemolysin			
		1	2	3	4	1	2	3	4	1	2	3	4
<i>C. kefyr</i>	79	1	9	20	49	79	–	–	–	47	23	9	–
<i>C. guilliermondii</i>	73	1	5	7	60	73	–	–	–	21	46	6	–
<i>C. famata</i>	52	1	–	9	42	52	–	–	–	17	31	4	–
<i>C. glabrata</i>	13	–	–	4	9	13	–	–	–	13	–	–	–
<i>C. albicans</i>	3	–	–	–	3	–	2	1	–	–	1	2	–
<i>C. parapsilosis</i>	1	–	–	–	1	1	–	–	–	1	–	–	–
Total	221	3	14	40	164	218	2	1	–	99	101	21	–

Enzyme activity: 1 = null (Pz = 1); 2 = low (Pz = 0.61–0.99); 3 = medium (Pz = 0.41–0.60); 4 = high (Pz ≤ 0.40).

$$Pz = \frac{\text{Diameter of the colony}}{\text{Diameter of the halo}}$$
Table 3Biofilm formation of *Candida* isolates from the milk of clinically healthy goats.

Candida species	Number of isolates (n)	No biofilm		Low biofilm formation		Moderate biofilm formation		High biofilm formation	
		OD ≤ 0.120 ^a	%	OD (0.121–0.240)	%	OD (0.241–0.480)	%	OD (≥ 0.481)	%
<i>C. kefyr</i>	79	21	9.5	50	22.62	7	3.16	1	0.45
<i>C. guilliermondii</i>	73	21	9.5	44	19.9	8	3.61	–	–
<i>C. famata</i>	52	18	8.14	26	11.8	6	2.71	2	0.9
<i>C. glabrata</i>	13	2	0.9	10	4.52	1	0.45	–	–
<i>C. albicans</i>	3	–	–	3	1.35	–	–	–	–
<i>C. parapsilosis</i>	1	1	0.45	–	–	–	–	–	–
Total	221	63	28.49	133	60.19	22	9.93	3	1.35

OD: optical density.

^a Cut-off point (mean OD value of negative controls + 2 standard deviation).

Financing

Program for Support to Projects for the Innovation and Improvement of Teaching (PAPIME) grant PE206819. DGAPA-UNAM. México.

Conflict of interests

The authors declare that they have no conflict of interest.

Acknowledgments

The authors are grateful to Dr. Cristina Escalante Ochoa for her valuable review of this manuscript.

References

- Arvanitis M, Anagnostou T, Fuchs BB, Caliendo AM, Mylonakis E. Molecular and nonmolecular diagnostic methods for invasive fungal infections. *Clin Microbiol Rev.* 2014;27(July):490–526. <http://dx.doi.org/10.1128/CMR.00091-13>.
- Barnett J, Payne R, Yarrow D. Yeast: characteristics and identification. Cambridge University Press; 1983.
- Bezerra de Melo RE, Menezes RP, Amante PMP, Dos Santos PR. Enzymatic and hemolytic in different *Candida* species. *Rev Iberoam Micol.* 2015;32:79–82. <http://dx.doi.org/10.1016/j.riam.2013.11.003>.
- Borst A, Fluit AC. High levels of hydrolytic enzymes secreted by *Candida albicans* isolates involved in respiratory infections. *J Med Microbiol.* 2003;52:971–4. <http://dx.doi.org/10.1099/jmm.0.05228-0>.
- Campbell MC, Stewart JL. The medical mycology handbook. University of Michigan; 1980.
- Campos PR, Franco GR, Rosa CA, Hahn RC, Hamdan JS. Phenotypic and genotypic identification of *Candida* spp. isolated from hospitalized patients. *Rev Iberoam Micol.* 2004;21:24–8.
- Costa EO, Gandra CR, Pires MF, Coutinho SD, Castillo W, Teixeira CM. Survey of bovine mycotic in dairy herds in the State of São Paulo, Brazil. *Mycopathologia.* 1993;124:7–13. <http://dx.doi.org/10.1007/BF01103051>.
- Da Costa KRC, Ferreira JC, Komesu MC, Candido RC. *Candida albicans* and *Candida tropicalis* in oral candidosis: quantitative analysis, exoenzyme activity, and antifungal drug sensitivity. *Mycopathologia.* 2009;167:73–9. <http://dx.doi.org/10.1007/s11046-008-9154-8>.
- Delavenne E, Mounier J, Asmani K, Jany JL, Barbier G, Le Blay G. Fungal diversity in cow, goat and ewe milk. *Int J Food Microbiol.* 2011;151:247–51. <http://dx.doi.org/10.1016/j.ijfoodmicro.2011.08.029>.
- De la Calle RN, Santa VC, Cardona CN. Factores de virulencia para la infección de tejidos queratinizados por *Candida albicans* y hongos dermatofitos. *Rev CES Med.* 2012;26:43–55. <http://www.scielo.org.co/pdf/cesm/v26n1/v26n1a05.pdf>.
- Del Pozo JL, Cantón E. Candidiasis asociada a biopelículas. *Rev Iberoam Micol.* 2016;33:176–83. <http://dx.doi.org/10.1016/j.riam.2015.06.004>.
- Fatahinia M, Halvaezadeh M, Rezaei-Matehkolaei A. Comparison of enzymatic activities in different *Candida* species isolated from women with vulvovaginitis. *J Mycol Méd.* 2017;27:188–94. <http://dx.doi.org/10.1016/j.mycmed.2017.01.009>.
- Flemming HC, Wingender J. The biofilm matrix. *Nat Rev Microbiol.* 2010;8:623–33. <http://dx.doi.org/10.1038/nrmicro2415>.
- Freydere AM, Guinet R, Boirin P. Yeast identification in the clinical microbiology laboratory: phenotypic methods. *Med Mycol.* 2001;39:9–33. <http://dx.doi.org/10.1080/mmy.39.1.9.33>.
- Gökce G, Cerikcioglu N, Yagci A. Acid proteinase, phospholipase, and biofilm production of *Candida* species isolates from blood cultures. *Mycopathologia.* 2007;164:265–9. <http://dx.doi.org/10.1007/s11046-007-9053-4>.
- Gómez J, Gómez-Lus ML, Bas P, Ramos C, Cafini F, Maestre JR, et al. ¿Es la cuantificación del biofilm un elemento diferenciador en la patogenia de bacilos gramnegativos? *Rev Esp Quimioter.* 2013;26:97–102.
- Hassan AA, Hassan MA, Rasha MH, Sayed EA, Darwish AS. Prevalence of yeast infections in small ruminants with particular references to their treatment by some natural herbal extracts. *Bull Environ Pharmacol Life Sci.* 2012;1:12–22.
- Hayashi T, Sugita T, Hata E, Katsuda K, Zhang E, Kiku Y, et al. Molecular-based identification of yeasts isolated from bovine clinical mastitis in Japan. *J Vet Med Sci.* 2013;75:387–90. <http://dx.doi.org/10.1292/jvms.12-0362>.
- Ilhan Z, Ekin IH, Koltas S, Gulaydin O, Ozturk C, Borum AE. Occurrence of fungal agents in mastitis in dairy goats. *J Anim Plant Sci.* 2016;29:4691–700.
- Kantarcioglu AS, Yücel A. Phospholipase and protease activities in clinical *Candida* isolates with reference to the sources of strains. *Mycoses.* 2002;45(5–6):160–5. <http://dx.doi.org/10.1046/j.1439-0507.2002.00727.x>.
- Krukowski H, Tietze M, Majewski T, Różański P. Survey of yeast mastitis in dairy herds of small-type farms in the Lublin region, Poland. *Mycopathologia.* 2001;150:5–7. <http://dx.doi.org/10.1023/A:1011047829414>.
- Ksouri S, Djebir S, Hadeif Y, Benakhla A. Survey of bovine mycotic mastitis in different mammary gland statuses in two north-eastern regions of Algeria. *Mycopathologia.* 2015;179:327–31. <http://dx.doi.org/10.1007/s11046-014-9845-2>.
- Kuhn DM, Chandra J, Mukherjee PK, Ghannoum MA. Comparison of biofilms formed by *Candida albicans* and *Candida parapsilosis* on bioprosthetic surfaces. *Infect Immun.* 2002;70. <http://dx.doi.org/10.1128/iai.70.2.878-888.2002>.

24. Kurtzman FC, Fell SJ, Boekhout ST. The yeast. In: A taxonomic study. 5th ed. Elsevier Science; 2011.
25. Lagneau PE, Lebtahi K, Swinne D. Isolation of yeast from bovine milk in Belgium. Mycopathologia. 1996;135:99–102, <http://dx.doi.org/10.1007/BF00436458>.
26. Luo G, Samaranayake LP, Yau JY. *Candida* species exhibit differential *in vitro* hemolytic activities. J Clin Microbiol. 2001;39:2971–4, <http://dx.doi.org/10.1128/JCM.39.8.2971-2974.2001>.
27. MacFaddin JF. Biochemical tests for identification of medical bacteria. 3rd ed. Jones and Bartlett Learning; 2000.
28. Nickerson WJ. Reduction of inorganic substances by yeasts: I. Extracellular reduction of sulfite by species of *Candida*. J Infect Dis. 1953;93:43–56, <http://dx.doi.org/10.1093/infdis/93.1.43>.
29. Panizo MM, Reviákina V, Flores Y, Montes W, González G. Actividad de fosfolipasas y proteasas en aislados clínicos de *Candida* spp. Rev Soc Venezol Microbiol [electronic journal]. 2005;25:64–71. Available in: <http://ve.scielo.org/scielo.php?script=sci.arttext&pid=S1315-25562005000200006&lng=es&tlng=es> [accessed 18.08.20].
30. Pontón J. Diagnóstico microbiológico de las micosis. Rev Iberoam Micol. 2002;19:25–9.
31. Ramage G, Vadevalle K, Wickes LB, López-Ribbot J. Characteristics of biofilm formation by *Candida albicans*. Rev Iberoam Micol. 2001;18:163–70.
32. Ruiz AJ, García MP, Puerto JL, Marín P, Saldaña A, Moya P. Evaluación de un nuevo medio CHROMagar *Candida* para la identificación presuntiva de levaduras. Rev Diagn Biol [electronic journal]. 2003;52:19–22. Available in: <http://scielo.isciii.es/scielo.php?script=sci.arttext&pid=S0034-79732003000100002&lng=es&nrm=iso/> [accessed 19.08.20].
33. Sacristán B, Blanco MT, Galán-Ladero MA, Blanco J, Pérez-Giraldo C, Gómez-García AC. Aspartyl proteinase, phospholipase, hemolytic activities and biofilm production of *Candida albicans* isolated from bronchial aspirates of ICU patients. Med Mycol. 2011;49:94–7, <http://dx.doi.org/10.3109/13693786.2010.482947>.
34. Segundo ZC, Cervantes ORA, Ducoing WAE, De la Peña MA, Villa TL. Yeasts isolation from bovine mammary glands under different mastitis status in the Mexican High Plateau. Rev Iberoam Micol. 2011;28:79–82, <http://dx.doi.org/10.1016/j.riam.2011.01.002>.
35. Seker E, Zenc E. *In vitro* biofilm activity of *Candida* species isolated from Anatolian buffaloes with mastitis in Western Turkey. Veterinaski Archiv. 2011;81:723–30.
36. Shirkahani S, Sepahvand A, Mirzaee M, Anbari K. Phospholipase and proteinase activities of *Candida* spp. Isolates from vulvovaginitis in Iran. J Mycol Méd. 2016;26:255–60, <http://dx.doi.org/10.1016/j.mycmed.2016.05.001>.
37. Spanemberg A, Wunder JREA, Brayer PDI, Argenta J, Cavallini SEM, Laerte FPV. Diversity of yeasts from bovine mastitis in Southern Brazil. Rev Iberoam Micol. 2008;25:154–6, [http://dx.doi.org/10.1016/S1130-1406\(08\)70036-6](http://dx.doi.org/10.1016/S1130-1406(08)70036-6).
38. Spanemberg A, Ramos JP, Leoncini O, Alves SH, Valente P. High frequency of potentially pathogenic yeast species in goat's raw milk and creamed cheese in Southern Brazil. Scient Veter. 2009;37:133–41.
39. Stepanovic S, Vukovic D, Dakic I, Savic B, Svabic VM. A modified microtiter-plate test for quantification of staphylococcal biofilm formation. J Microbiol Methods. 2000;40:175–9, [http://dx.doi.org/10.1016/S0167-7012\(00\)00122-6](http://dx.doi.org/10.1016/S0167-7012(00)00122-6).
40. St-Germain G, Summerbell R. Identifying fungi. In: A clinical laboratory handbook. 2nd ed. Star Publishing Company, Inc.; 2011.
41. Williamson MI, Samaranayake LP, MacFarlane TW. Phospholipase activity as a criterion for biotyping *Candida albicans*. J Med Vet Mycol. 1986;24:415–7, <http://dx.doi.org/10.1080/02681218680000631>.
42. Yigit N, Aktas E, Dagistan S, Ayyildiz A. Investigating biofilm production, coagulase and hemolytic activity in *Candida* species isolated from denture stomatitis patients. Eurasian J Med. 2011;43:27–32, <http://dx.doi.org/10.5152/eajm.2011.06>.
43. Yis R, Doluca M. Identification of *Candida* species by restriction enzyme analysis. Turk J Med Sci. 2018;48:1058–106, <http://dx.doi.org/10.3906/sag-1802-11>.
44. Ziccardi M, Souza LO, Gandra RM, Galdino AC, Baptista AR, Nunes AP, et al. *Candida parapsilosis* (*sensu lato*) isolated from hospitals located in the Southeast of Brazil: Species distribution, antifungal susceptibility and virulence attributes. Int J Med Microbiol. 2015;305:848–59, <http://dx.doi.org/10.1016/j.ijmm.2015.08.00>.