



REVISTA IBEROAMERICANA DE PSICOLOGÍA Y SALUD

Revista Oficial de la Federación Iberoamericana de Asociaciones de Psicología (FIAP)
[Official Journal of the Latin-American Federation of Psychological Associations]

www.elsevier.es/rips



The effect of recording interval length on behavioral assessment using the forced swimming test



Paloma Álvarez-Suárez^a, María Banqueri^a, Marina Vilella^a,
Marta Méndez^{a,b,*}, Jorge L. Arias^{a,b}

^a Laboratory of Neurosciences, Faculty of Psychology, University of Oviedo, Oviedo, Spain

^b Institute of Neuroscience of the Principality of Asturias, University of Oviedo, Oviedo, Spain

Received 8 April 2015; accepted 28 April 2015

Available online 30 May 2015

KEYWORDS

Forced swimming test;
Interval length;
Recording method;
Rats

Abstract The forced swimming test is a method used in the assessment of depressive-like behavior in rodents. Changes in the original forced swimming test procedure developed by Porsolt et al. and their influence in the results is a controversial issue and has been discussed in many studies. Animal's behavior is usually recorded by partial interval recording, dividing the total recording time into equal intervals and manually recording the predominant behavior. Despite the influence of the recording method in the subsequent results, this issue has not been further studied nor normalized. The aim of this study was to assess whether the representativeness of the data is influenced by the recording interval length, by recording behaviors (immobility, swim and climbing) in the same subjects at 3, 5 and 10 s recording intervals. We used a non-pathological sample of male and female adult Wistar rats. Our results show no differences in the use of these three recording intervals in the registration method of the forced swimming test, for the main three behaviors measured.

© 2015 Sociedad Universitaria de Investigación en Psicología y Salud. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Test de natación forzada;
Longitud del intervalo;
Método de registro;
Ratas

El efecto de la longitud del intervalo de registro en la evaluación conductual mediante el test de natación forzada

Resumen El test de natación forzada es un método utilizado para evaluar el comportamiento depresivo en roedores. Las modificaciones del procedimiento del test de natación forzada desarrollado por Porsolt y su influencia en sus resultados es un tema que suscita controversia y ha sido analizado en numerosos estudios. El comportamiento del animal se analiza generalmente mediante el registro de intervalos parciales, dividiendo el tiempo total de grabación en intervalos iguales y registrando manualmente la conducta predominante durante ese tiempo. A pesar

* Corresponding author.

E-mail address: mendezlmarta@uniovi.es (M. Méndez).

de la influencia del método de registro en los resultados posteriores, esta cuestión no se ha analizado ni normalizado. El objetivo de este estudio fue comprobar si la representatividad de los datos se ve afectada por la longitud del intervalo de observación, registrando a 3, 5 y 10 seg la conducta (inmovilidad, natación y escalada) de los mismos sujetos. Se utilizó una muestra no patológica de ratas Wistar macho y hembra adultas. Los resultados mostraron que no existen diferencias significativas entre estos intervalos analizados en la prueba de natación forzada para las 3 principales conductas registradas.

© 2015 Sociedad Universitaria de Investigación en Psicología y Salud. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The forced swimming test (FST) is a method used in the assessment of depressive-like behavior in rodents that was developed by [Porsolt, Bertin, and Jalfre \(1977\)](#). It is based on learned helplessness, as the animal is repeatedly exposed to an aversive stimulus, water, which it cannot escape. When the animal is introduced in an inescapable cylinder filled with water, it learns that there is nothing it can do to escape and, therefore, it reduces the time of regular escape behaviors (swimming, climbing and diving) in subsequent trials and spends more time making no movements. In the animal model of depression, the total time of immobilization is higher than in control condition and it has been seen that antidepressants reduce the differences in total time of immobility between control and experimental groups ([Porsolt, LePichon, & Jalfre, 1977](#)).

An animal model of a particular psychological condition must meet certain criteria to be applicable to humans: it should resemble the human condition in its etiology, biochemistry, symptoms and treatment ([McKinney & Bunney, 1969](#)). As reviewed in the text of [Belzung and Lemoine \(2011\)](#), most authors have focused on these external validity criteria. In this review, five basic criteria of validity of an animal model, which differ slightly from the classical criteria, are proposed. These criteria are homological validity, pathogenic validity, mechanistic validity, face validity and predictive validity ([Belzung & Lemoine, 2011](#)). According to many of these criteria, major depressive disorder has been reproduced in animals in order to study the underlying neurobiological mechanisms. This model has been developed by replicating aspects of the depressive syndrome that are not intrinsically human features, i.e. anhedonia, helplessness and behavioral despair ([Krishnan & Nestler, 2011](#)).

The validity of the FST has been questioned many times. Several authors have compared the results of FST with other depression-measuring tests, such as sucrose preference test ([Grillo et al., 2011; Hong et al., 2012; Karson, Demirtas, Bayramgürler, Balci, & Utkan, 2013](#)), and have found consistent results. Nevertheless, there is controversy about the validity of FST to measure behavioral 'despair', as habituation has been proposed as an explanation to immobility ([Hawkins, Hichs, Phillips, & Moore, 1978](#)) due to a process of familiarization ([Borsini, Volterra, & Meli, 1986](#)). There are multiple variations on the original methodology of the FST that lead to differences in the results ([Borsini & Meli,](#)

1988). While most of the researchers record behaviors manually ([Grillo et al., 2011; Kawai, Ishibashi, Kudo, Kawashima, & Mitsumoto, 2012; Sirianni, Olausson, Chiu, Taylor, & Saltzman, 2010; Ulloa, Díaz-Valderrama, Herrera-Pérez, León-Olea, & Martínez-Mota, 2014](#)), others use automated devices ([El-Alfy et al., 2010; Uz, Dimitrijevic, Imbesi, Manev, & Manev, 2008](#)). Therefore, behavioral results used to be based on subjective recording.

Animal's behavior is usually recorded by partial interval recording (PIR), consisting on dividing the total recording time into equal intervals and manually recording the predominant behavior in each interval ([Borsini & Meli, 1988](#)). Some authors have also tried to automate the recording process by developing software that measures different mobility parameters, such as Ethovision 3.0 by Noldus ([Hédou, Pryce, Di Iorio, Heidbreder, & Feldon, 2001](#)) or CVA software by ProTrack ([Gersner, Gordon-Kiwkowitz, & Zangen, 2009](#)), although the preferential method is still the trained observer. In [Fiske and Delmolino \(2012\)](#), advantages and disadvantages of different recording methods are mentioned. Using PIR, some authors conclude that the smaller the interval, the lower the absolute and relative errors ([Wirth, Slaven, & Taylor, 2014](#)). Also, it has been observed that the length of the recording interval inversely affects the representativeness of data ([Repp, Roberts, Slack, Repp, & Berkler, 1976](#)). Finally, regarding recording methods, it has been shown that PIR method is a more sensitive recording method than momentary time sampling (MTS) ([Harrop & Daniels, 1986](#)).

We have found in the literature (summarized in [Table 1](#)) that many studies using FST do not include information in their methodological section about the recording method. Most of the few authors who describe the recording method usually record behavior at intervals of 5 s. There are exceptions such as [Su, Hato-Yamada, Araki, and Yoshimura \(2013\)](#), who quantify the duration and frequency of the behavior, and [Grillo et al. \(2011\)](#), who recorded at intervals of 3 s. Despite the influence of the recording method in the subsequent results, this issue has not been further studied nor normalized.

In the present study, we assessed the relevance of using a longer or shorter interval in the recording method of the FST. To do this, we compared the behavioral results of the FST in the same sample of animals at 3, 5 and 10 s recording intervals. We expected to find more accurate results in the shortest interval and a masking effect in the widest one.

Table 1 Summary of the studies that used forced swimming test.

Authors	Recording method	Interval size (s)	Software
Citó et al. (2015)	Unspecified		
Martínez, Brunelli, and Zimmerberg (2015)	Unspecified		
Ulloa et al., 2014	PIR	5	
Kołaczkowski et al. (2014)	Unspecified		
Karson et al. (2013)	Unspecified		
Yang, Hu, Zhou, Zhang, and Yang (2013)	Unspecified		
Su et al. (2013)	Duration and frequency		
Kawai et al. (2012)	PIR	5	
Hong et al. (2012)	PIR	5	
Grillo et al. (2011)	PIR	3	
El-Alfy et al. (2010)	Automated		SMART II Video Tracking
Sirianni et al. (2010)	PIR	5	
Uz et al. (2008)	Automated		MotorMonitor 4.11
Kuœmider, Solich, Pasach, and Dziedzicka-Wasylewska (2007)	PIR	5	
Deak et al. (2005)	PIR	5	
Holmes, Yang, Murphy, and Crawley (2001)	PIR	5	
Lucki, Dalvi, and Mayorga (2001)	Unspecified		
Hédou et al. (2001)	Automated		Ethovision 3.0

Method

Animals

A total of 10 (4 male/6 female) adult Wistar rats (*Rattus norvegicus*) between 80 and 150 g were used. The animals were obtained from the University of Oviedo central vivarium (Oviedo, Asturias, Spain). They were housed under standard conditions (12-h light/dark cycle with lights on from 08:00 to 20:00 h), at constant room temperature of $23 \pm 2^\circ\text{C}$ with ad libitum access to food and water. All experimental procedures carried out with animals were approved by a local veterinary committee from the University of Oviedo vivarium and subsequent handling strictly followed the European Communities Council Directive 2010/63/UE. All efforts were made to minimize the number of animals used and their suffering.

Apparatus

The FST was performed in a Plexiglas cylindrical bin (20 cm diameter, 50 cm high) filled with water ($23\text{--}27^\circ\text{C}$ water temperature) to a depth of 30 cm. Essays were recorded using an automated video-tracking system (Noldus MPEG-4 Recorder V1.1.6). Then, they were burn in a DVD and analyzed by two experimented researchers.

Procedure

Rats were placed individually into the Porsolt bin. We used 10 min in the first day for the habituation as previously described (Kawai et al., 2012). Next day, test was performed and the animals were placed in the Porsolt bin for 5 min, then

carefully dried and put it again in their house-cage. Between every subject the water was removed and filled again in order to avoid any smell trail. Essays were analyzed by two experimented researchers. They analyzed animal's behavior (immobility, swimming and climbing) for three different intervals: 3, 5 and 10 s. Immobility behavior was considered when the rats was floating passively, making small movements to keep its nose above the water surface. Swimming was measured when the rats made horizontal movements more than those necessary to merely keep the head above the water. Climbing was defined as when the rats was in active vertical motion; trying to escape.

Data analysis

All data were analyzed by SigmaStat 3.2 software (Systat Software, Chicago, USA) and were expressed as mean $\pm$ SEM. The results were considered statistically significant when $p < .05$.

For each behavior a one factor ANOVA with three levels were performed (dependent variable: swimming, climbing or immobility; independent variable: 3 s interval, 5 s interval and 10 s interval).

For each interval a one factor ANOVA with three levels were performed (dependent variable: 3 s interval, 5 s interval or 10 s interval; independent variable: swimming, climbing and immobility).

A total of six one factor ANOVA with three levels were performed.

Results

Data were analyzed using one way ANOVA. When significant differences were found, tests for multiple comparisons

Table 2 Mean and standard error mean (SEM) of the different behaviors in male and female rats.

Interval	Behaviors	Male	Female	T-test value
3 s	Immobility	40.83 ± 10.28	53.14 ± 6.83	0.327
	Swim	43.33 ± 10.25	31.85 ± 6.75	0.355
	Climbing	15.83 ± 4.82	15 ± 2.89	0.878
5 s	Immobility	56.48 ± 9.36	62.65 ± 3.18	0.484
	Swim	28.70 ± 7.03	20.06 ± 4.52	0.307
	Climbing	14.81 ± 2.92	16.97 ± 2.59	0.602
10 s	Immobility	55.55 ± 2.61	67.90 ± 5.63	0.131
	Swim	28.70 ± 2.77	15.43 ± 4.62	0.064
	Climbing	15.74 ± 3.81	16.66 ± 3.13	0.856

(Tukey's tests) were used to identify differences. Normality and equal variances are assumed.

T-test for independent samples showed no significant differences between sexes in behavioral results of the FST at 3, 5 and 10 s intervals (Table 2).

We analyzed whether there were differences in each behavior, according to the different types of intervals used (Table 3). Immobility behavior shows no significant differences between intervals ($F(2, 27) = 2.816$; $p = .078$). Also, we obtain the same results for swim behavior ($F(2, 27) = 3.455$; $p = .046$) and climbing ($F(2, 27) = 0.053$; $p = .948$).

In general, we found significant differences between behaviors, where immobility is greater than swimming and climbing ($F(2, 99) = 9.204$; $p = .001$). Tukey test shown that these differences occur when behavioral observation was performed at 5 and 10 s interval ($p = .012$ and $p = .003$, respectively), but not for 3 s interval ($p = .462$).

Cohen's kappa coefficient was used to assess agreement between the two judges. This statistic showed substantial agreement ($k = 0.759$), $p < .05$, 95% CI [0.670–0.847], according to Landis and Koch (1977).

Discussion

Being aware of the lack of information relative to the use of different observational intervals in the FST, we examined if there are significant differences between the use of 3, 5 or 10 s intervals of recording in the behaviors observed in this test: swimming, climbing and immobility. Our results indicate that there are no differences between the recording intervals in any of the behaviors observed.

Despite the fact that most of the researches use only male animals when they assess behavior in the FST, we used both sexes, trying to assess behaviors in a more representative sample. We found no differences between sexes in behavioral results at any interval. This can be easily

explained by the fact that our animals are control animals and they are not modeling any disturbance or disease. However, in many animal models of disease both sexes should be analyzed, as sex differences in behaviors exist in some pathological conditions (Bai et al., 2014; Papaioannou, Geroziannis, Prokopiou, Bolaris, & Stylianopoulou, 2002).

We found no differences between the use of 3, 5 or 10 s intervals in any of the behaviors observed: swimming, climbing and immobility. Some authors conclude that the smaller the interval, the lower the absolute and relative errors (Wirth et al., 2014), however our results point out a different idea. One reason for this discrepancy could be that Wirth et al. (2014) explore longer intervals than us: they used intervals between 15 s and 450 s.

In other author's opinion, the length of the recording interval inversely affects the representativeness of the data (Repp et al., 1976). However, in our study we found that in the large recording intervals (10 s) and in the standard medium interval used in most of the studies (5 s), there are no differences for between behaviors. In both large and medium recording intervals, we found more immobility than swimming and climbing. In our case, the use of the short recording intervals (3 s), could be less representative because it might be lead to the loss of data. This loss of information could be due to the excessive atomization of the behaviors that usually occur for more than 3 s. Also, we should take into account that our behavioral data are high homogenous, as they are shown by a control sample, while, in a sample with more heterogeneous behaviors, the excessive long intervals, could lead to a loss of information. In an excessive long interval, the animal behavior might change many times. In this case, the recording interval length should be reconsidered.

Our objective in this work was to explore the consequences of using different intervals in the recording of the FST. Although there are many published works about the correct use of methodology in this test (Bogdanova, Kanekar,

Table 3 Means and standard error mean (SEM) of the behaviors in the different intervals.

Interval	Immobility	Swim	Climbing
3 s	48.222 ± 5.807	36.444 ± 5.721	15.333 ± 2.433
5 s	60.185 ± 4.012	23.519 ± 3.924	16.111 ± 1.873
10 s	62.963 ± 3.943	20.741 ± 3.587	16.296 ± 2.290

D'Anci, & Renshaw, 2013; Petit-Demouliere, Chenu, & Bourin, 2005), to the best of our knowledge, there is not any published the effect of the recording interval on the behaviors observed in the test.

The majority of the published works not even describe the interval used (see Table 1). Some authors indicate that they use 5 s interval when they analyze behaviors in the FST. The reason for using this interval instead of another is rarely described. However, according to our results, recording at 5 s interval seems to be a suitable method.

It is important to mention that we are using a subjective recording method and, for this reason, the observers had to be trained. At the same time, it is highly recommended that behaviors were analyzed by more than one examiner. Regarding examiners' training, it would be great interest to use more than one animal in order to be able to recognize behaviors that suffer mild changes between different subjects.

In our study, we used two trained observers and we get a substantial interjudge agreement. This agreement determines a great consistency among raters, and suggests a good implementation of the measurement system.

A limitation of our work is that it was performed with only control animals. The reason for choosing this non-pathological sample is that we wanted to assess the reliability of the interval used by the examiners of behavior without considering any other variable. Normally, this test is widely used for depression models (Kawai et al., 2012; Shaldubina, Einat, Bersudsky, & Belmaker, 2005; Ulloa et al., 2014), or related models, like stress-induced anhedonia (Briones et al., 2012). For this reason, future research should check whether these results are equal for pathological models, or if in these models we need to use a specific recording interval.

We conclude that there are no differences in the use of these three recording intervals (3, 5 and 10 s) in the registration method of the forced swimming test, for the main three behaviors measured: climbing, swimming and immobility. Perhaps, in some cases, the use of 3-s recording intervals could drive to a loss of relevant information.

Conflicts of interest

All authors declare that there are no actual or potential conflicts of interest including any financial, personal or other relationships with other people or organizations that could inappropriately influence this work.

Acknowledgements

This research was supported by Project Grant of the Spanish Ministry of Economy and Competitiveness PSI2013-45924-P and BES-2014-070562 to M.B.L., and Consejería de Economía y Competitividad del Gobierno del Principado de Asturias.

References

- Bai, M., Zhang, L., Zhu, X., Zhang, Y., Zhang, S., & Xue, L. (2014). Comparison of depressive behaviors induced by three stress paradigms in rats. *Physiology & Behavior*, 131, 81–86.
- Belzung, C., & Lemoine, M. (2011). Criteria of validity for animal models of psychiatric disorders: Focus on anxiety disorders and depression. *Biology of Mood & Anxiety Disorders*, 1, 9.
- Bogdanova, O. V., Kanekar, S., D'Anci, K. E., & Renshaw, P. F. (2013). Factors influencing behavior in the forced swim test. *Physiology & Behavior*, 118, 227–239.
- Borsini, F., & Meli, A. (1988). Is the forced swimming test a suitable model for revealing antidepressant activity? *Psychopharmacology*, 94, 147–160.
- Borsini, F., Volterra, G., & Meli, A. (1986). Does the behavioral despair test measure 'despair'? *Physiology & Behavior*, 38, 385–386.
- Briones, A., Gagno, S., Martisova, E., Dobarro, M., Aisa, B., Solas, M., et al. (2012). Stress-induced anhedonia is associated with an increase in Alzheimer's disease related markers. *British Journal of Pharmacology*, 165, 897–907.
- Citó, M. C. O., Silva, M. I. G., Santos, L. K. X., Fernandes, M. L., Melo, F. H. C., Aguiar, J. A. C., et al. (2015). Antidepressant-like effect of *Hoodia gordonii* in a forced swimming test in mice: Evidence for involvement of the monoaminergic system. *Brazilian Journal of Medical and Biological Research*, 48, 57–64.
- Deak, T., Bellamy, C., D'Agostino, L. G., Rosanoff, M., McElderry, N. K., & Bordner, K. A. (2005). Behavioral responses during the forced swim test are not affected by anti-inflammatory agents or acute illness induced by lipopolysaccharide. *Behavioural Brain Research*, 160, 125–134.
- El-Alfy, A. T., Ivey, K., Robinson, K., Ahmed, S., Radwan, M., Slade, D., et al. (2010). Antidepressant-like effect of Δ^9 -tetrahydrocannabinol and other cannabinoids isolated from *Cannabis sativa*. *Pharmacology Biochemistry and Behaviour*, 95, 434–442.
- Fiske, K., & Delmolino, L. (2012). Use of discontinuous methods of data collection in behavioral intervention: Guidelines for practitioners. *Behavior Analysis in Practice*, 5, 77–81.
- Gersner, R., Gordon-Kiwkowitz, M., & Zangen, A. (2009). Automated behavioral analysis of limbs' activity in the forced swim test. *Journal of Neuroscience Methods*, 180, 82–86.
- Grillo, C. A., Piroli, G. G., Kaigler, K. F., Wilson, S. P., Wilson, M. A., & Reagan, L. P. (2011). Downregulation of hypothalamic insulin receptor expression elicits depressive-like behaviors in rats. *Behavioural Brain Research*, 222, 230–235.
- Harrop, A., & Daniels, M. (1986). Methods of time sampling: A reappraisal of Momentary Time Sampling and Partial Interval Recording. *Journal of Applied Behavior Analysis*, 19, 73–77.
- Hawkins, J., Hichs, R. A., Phillips, N., & Moore, J. D. (1978). Swimming rats and human depression. *Nature*, 274–512.
- Hédou, G., Pryce, C., Di Iorio, L., Heidbreder, C. A., & Feldon, J. (2001). An automated analysis of rat behavior in the forced swim test. *Pharmacology Biochemistry and Behavior*, 70, 65–76.
- Holmes, A., Yang, R. J., Murphy, D. L., & Crawley, J. N. (2001). Evaluation of antidepressant-related behavioral responses in mice lacking the serotonin transporter. *Neuropsychopharmacology*, 27, 914–923.
- Hong, S., Flashner, B., Chiu, M., ver Hoeve, E., Luz, S., & Bhatnagar, S. (2012). Social isolation in adolescence alters behaviors in the forced swim and sucrose preference tests in female but not in male rats. *Physiology & Behavior*, 105, 269–275.
- Karson, A., Demirtas, T., Bayramgürler, D., Balci, F., & Utkan, T. (2013). Chronis administration of infliximab (TNF- α inhibitor) decreases depression and anxiety-like behavior in rat model of chronic mild stress. *Basic & Clinical Pharmacology & Toxicology*, 112, 335–340.
- Kawai, H., Ishibashi, T., Kudo, N., Kawashima, Y., & Mitsumoto, A. (2012). Behavioral and biochemical characterization of rats treated chronically with thioacetamide: Proposal of an animal model for hepatic encephalopathy associated with cirrhosis. *Journal of Toxicological Sciences*, 37, 1165–1175.

- Kołaczowski, M., Marcinkowska, M., Bucki, A., Śniecikowska, J., Pawłowski, M., Kazek, G., et al. (2014). Novel 5-HT₆ receptor antagonists/D₂ receptor partial agonists targeting behavioral and psychological symptoms of dementia. *European Journal of Medicinal Chemistry*, 92, 221–235.
- Krishnan, V., & Nestler, E. J. (2011). Animal models of depression: Molecular perspectives. *Current Topics in Behavioral Neurosciences*, 7, 121–147.
- Kuøemider, M., Solich, J., Pasach, P., & Dziedzicka-Wasylewska, M. (2007). Effect of citalopram in the modified forced swim test in rats. *Pharmacological Reports*, 59, 785–788.
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33, 159–174.
- Lucki, I., Dalvi, A., & Mayorga, A. J. (2001). Sensitivity to the effects of pharmacologically selective antidepressants in different strains of mice. *Psychopharmacology*, 155, 315–322.
- Martínez, A. R., Brunelli, S. A., & Zimmerberg, B. (2015). Communal nesting exerts epigenetic influences on affective and social behaviors in rats selectively bred for an infantile trait. *Physiology & Behavior*, 139, 97–103.
- McKinney, W. T., & Bunney, W. E. (1969). Animal model of depression: Review of evidence and implications for research. *Archives of General Psychiatry*, 21, 240–248.
- Papaioannou, A., Gerozissis, K., Prokopiou, A., Bolaris, S., & Stylianopoulou, F. (2002). Sex differences in the effects of neonatal handling on the animal's response to stress and the vulnerability for depressive behavior. *Behavioural Brain Research*, 129, 131–139.
- Petit-Demouliere, B., Chenu, F., & Bourin, M. (2005). Forced swimming test in mice: A review of antidepressant activity. *Psychopharmacology*, 177, 245–255.
- Porsolt, R. D., Bertin, A., & Jalfre, M. (1977). Behavioral despair in mice: A primary screening test for antidepressants. *Archives internationales de pharmacodynamie et de thérapie*, 229, 327–336.
- Porsolt, R. D., LePichon, M., & Jalfre, M. (1977). Depression: A new animal model sensitive to antidepressant treatments. *Nature*, 266, 730–732.
- Repp, A. C., Roberts, D. M., Slack, D. J., Repp, C. F., & Berkler, M. S. (1976). A comparison of frequency, interval and time-sampling methods of data collection. *Journal of Applied Behavior Analysis*, 9, 501–508.
- Shaldubina, A., Einat, H., Bersudsky, Y., & Belmaker, R. H. (2005). Epi-inositol is ineffective in Porsolt Forced Swim Test model of depression. *Neuropsychiatric Disease and Treatment*, 1, 180–190.
- Sirianni, R. W., Olausson, P., Chiu, A. S., Taylor, J. R., & Saltzman, W. M. (2010). The behavioral and biochemical effects of BDNF containing polymers implanted in the hippocampus of rats. *Brain Research*, 1321, 40–50.
- Su, J., Hato-Yamada, N., Araki, H., & Yoshimura, H. (2013). A test-retest paradigm of the forced swimming test in female mice is not valid for predicting antidepressant-like activity: Participation of acetylcholine and sigma-1 receptors. *Journal of Pharmacological Sciences*, 123, 246–255.
- Ulloa, R. E., Díaz-Valderrama, A., Herrera-Pérez, J., León-Olea, M., & Martínez-Mota, L. (2014). Age differences in the impact of forced swimming test on serotonin transporter levels in lateral septum and dorsal raphe. *Behavioral and Brain Functions*, 10, 1–8.
- Uz, T., Dimitrijevic, N., Imbesi, M., Manev, H., & Manev, R. (2008). Effects of MK-886, a 5-lipoxygenase activating protein (FLAP) inhibitor, and 5-lipoxygenase deficiency on the forced swimming behavior of mice. *Neuroscience Letters*, 436, 269–272.
- Wirth, O., Slaven, J., & Taylor, M. A. (2014). Interval sampling methods and measurement error: A computer simulation. *Journal of Applied Behavior Analysis*, 47, 83–100.
- Yang, C., Hu, Y. M., Zhou, Z. Q., Zhang, G. F., & Yang, J. J. (2013). Acute administration of ketamine in rats increases hippocampal BDNF and mTOR levels during forced swimming test. *Upsala Journal of Medical Sciences*, 118, 3–8.