



Erratum to: "Negative correlation between presence of reactive oxygen species and Sperm Motility Index in whole semen samples of infertile males"
[Rev Int Androl. 2017;15(3):84-89]

Fe de errores a "Correlación negativa entre la presencia de especies reactivas del oxígeno y el índice de movilidad espermática en muestras completas de semen de varones estériles"
[Rev Int Androl. 2017;15(3):84-89]

Shinnosuke Kuroda^{a,b}, Yasushi Yumura^{b,*}, Kohei Mori^b, Kengo Yasuda^b,
Teppei Takeshima^b, Takashi Kawahara^c, Yasuhide Miyoshi^c, Hiroji Uemura^c,
Akira Iwasaki^b, Kunitomo Takashima^d, Mario Ikeda^d, Yoshihito Kondo^d

^a Department of Urology, Yokohama Minato Red Cross Hospital, Kanagawa, Japan

^b Department of Urology, Reproductive Center, Yokohama City University Medical Center, Kanagawa, Japan

^c Department of Urology, Yokohama City University Medical Center, Kanagawa, Japan

^d Ebina Ladies' Clinic, Kanagawa, Japan

In the above published article, errors have been detected in the values of motility and concentration, and also in the caption of Figure 2.

The corrected values are:

Motility 0.538

Concentration 0.641

The corrected figure caption is:

Figure 2 The scatterplots of the relationship between SMI and each semen parameter using the Spearman correlation coefficient. There was significant correlation between SMI and motility ($\rho = 0.538$), concentration ($\rho = 0.641$), straight-line velocity (VSL) ($\rho = 0.352$), curvilinear velocity (VCL) ($\rho = 0.360$), mean amplitude of lateral head displacement (mALH) ($\rho = 0.60$), and progressive motility (PM) ($\rho = 0.578$), respectively ($p \leq 0.001$).

DOI of original article: <http://dx.doi.org/10.1016/j.androl.2016.08.002>

* Corresponding author.

E-mail address: yumura@yokohama-cu.ac.jp (Y. Yumura).