

4. Oelschläger BK, Lily Chang, Pellegrini CA. Improved outcome after extended gastric myotomy for achalasia. *Arch Surg.* 2003;138: 490-7.
5. Patti MG, Arcerito M, Tong J, et al. Pneumatic dilation versus transthoracic limited myotomy. *Am J Gastroenterol.* 1997;92:1109-12.
6. Topart P, Deschamps C, Taillefer R, Duranceau A. Long-term effect of total fundoplication on the myotomized esophagus. *Ann Thorac Surg.* 1992;54:1046-52.
7. Falkenback D, Johansson J, Oberg S, et al. Heller's esophagomyotomy with or without a 360 degrees floppy Nissen fundoplication for achalasia. Long-term results from a prospective randomized study. *Dis Esophagus.* 2003;16:284-90.
8. Lyass S, Thoman D, Steiner JP, et al. Current status of an antireflux procedure in laparoscopic Heller myotomy. *Surg Endosc.* 2003;17: 554-8.
9. Piccicocchi A, Cardillo G, D'Ugo D, Castrucci G, Mascellari L, Gragnone P. Surgical treatment of achalasia: a retrospective comparative study. *Surg Today.* 1993;23:855-9.
10. Richards WO, Torquati A, Holzman MD, Khaitan L, Byrne D, Lutfi R, et al. Heller myotomy versus Heller myotomy with dor fundoplication for achalasia. A prospective randomized double-blind clinical trial. *Ann Surg.* 2004;240:405-15.
11. Csendes A, Braghetto I, Burdiles P, Korn O, Csendes P, Henriquez A. Very late results of esophagomyotomy for patients with achalasia. *Ann Surg.* 2006;243:196-203.
12. Decker G, Borie F, Bouamrine D, Veyrac M, Guillon F, Fingerhut A, et al. Gastrointestinal quality of life before and after laparoscopic Heller myotomy with partial posterior fundoplication. *Ann Surg.* 2002;236:750-8.

Fe de errores

En el artículo "Esofagectomía transhiatal por vía abierta y vía laparoscópica para el cáncer de esófago: análisis de los márgenes de resección y ganglios linfáticos" (*Cirugía.* 2008;83[1]:24-7) se ha detectado un error. El nombre correcto del segundo autor es Sally Fares.