



REVISTA MÉDICA DEL
HOSPITAL GENERAL
DE MÉXICO

www.elsevier.es/hgmx



ORIGINAL ARTICLE

Exome sequencing analysis reveals homozygous GJB2 gene mutation in a Mexican family with profound hearing loss

M. Martínez-Saucedo^a, M.R. Rivera-Vega^a, L.M. Gonzalez-Huerta^a,
H. Urueta-Cuéllar^a, S.A. Cuevas-Covarrubias^{a,b,*}



CrossMark

^a Servicio de Genética, Hospital General de México, Mexico City, Mexico

^b Facultad de Medicina, Universidad Nacional Autónoma de México, Mexico City, Mexico

Received 18 July 2016; accepted 8 August 2016

Available online 22 August 2016

KEYWORDS

GJB2 gene;
Hypoacusia;
Gene mutations;
Connexins;
Homozygous

Abstract

Background: Sensorineural hearing loss (SNHL) is a clinically and genetically heterogeneous disease. In some populations, c.365delG mutation in the *GJB2* gene represents the most frequent cause of hereditary SNHL. The great diversity of mutations in the *GJB2* gene worldwide highlights the participation of ethnic background in SNHL.

Objective: To describe the presence of homozygous c.del35G mutation in the *GJB2* gene in a Mexican family with SNHL.

Materials and methods: A Mexican family with SNHL was included in the study. Analysis of the *GJB2* gene was performed through whole exome sequencing (WES) and DNA direct sequencing analysis in all members of the family and in 100 normal controls

Results: Affected sibs showed the homozygous c.del35G mutation in the *GJB2* gene. Parents of the families were heterozygous for the molecular defect and had normal audition.

Conclusion: We describe a homozygous c.del35G mutation in the *GJB2* gene through WES analysis, a homozygous mutation with a very low occurrence in Mexican population.

© 2016 Sociedad Médica del Hospital General de México. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Dr. Balmis 148 Col. Doctores, C.P. 06726 México, D.F., Mexico.

E-mail addresses: sergiocuevasunam@gmail.com, sercuezas@yahoo.com (S.A. Cuevas-Covarrubias).

PALABRAS CLAVE

Gene *GJB2*;
Hipoacusia;
Mutaciones génicas;
Conexinas;
Homocigoto

El análisis de secuenciación del exoma identifica una mutación homocigota en el gen *GJB2* en una familia mexicana con pérdida auditiva profunda

Resumen

Antecedentes: La pérdida auditiva neurosensorial (HNS) es una enfermedad clínica y genéticamente heterogénea. En algunas poblaciones, la mutación c.365delG en el gen *GJB2* representa la causa más frecuente de la HNS hereditaria. La gran diversidad de mutaciones presentes en el gen *GJB2* en todo el mundo pone de relieve la participación del genotipo étnico en HNS.

Objetivo: Describir la presencia de la mutación homocigota c.del35G en el gen *GJB2* en una familia mexicana con HNS.

Materiales y métodos: Una familia mexicana con HNS se incluyó en el estudio. El análisis del gen *GJB2* se realizó a través de la secuenciación del exoma (WES) y el análisis de la secuenciación directa del DNA en todos los miembros de la familia y en 100 controles normales. **Resultados:** Los hermanos afectados presentaron la mutación c.del35G homocigótica en el gen *GJB2*. Los padres de los pacientes fueron heterocigotos para el defecto molecular con una audición normal.

Conclusión: Se describe la mutación homocigota c.del35G en el gen *GJB2* a través del análisis de WES, una mutación homocigota observada en una muy baja incidencia en la población mexicana. © 2016 Sociedad Médica del Hospital General de México. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Non-syndromic sensorineural hearing loss (SNHL), a genetically heterogeneous affection, is a common hereditary disease that presents an occurrence of 1 in 1000 newborns. In relation to the onset, SNHL can be congenital, prelingual or postlingual. All types of inherited patterns have been observed in SNHL (autosomal dominant, autosomal recessive, mitochondrial and X-linked) being the autosomal recessive form the most frequently observed.¹ The syndromic form of sensorineural hearing loss is defined when other systemic manifestations are present; many syndromes with the presence of hearing loss are described in the OMIM (Online Mendelian Inheritance in Man). In some cases, differential diagnosis between syndromic and non-syndromic hearing loss is not easy to perform even with molecular studies because both types of hearing loss can be due to affections of the same gene.

It is considered that 200–250 genes are involved in SNHL.² Recently, more than 80 genes, around 1000 mutations, and 140 loci have been associated with SNHL (<http://hereditaryhearingloss.org/>).

Mutations in the *GJB2* gene are a frequent etiology of hereditary SNHL, they represent about 50% of such cases in several populations.^{3–5} Patients that carrier homozygous mutations in the *GJB2* gene harbor a wide spectrum of hearing loss that ranges from moderate to profound NSHL; this data supports the influence of epigenetic and environmental factors in the phenotypic manifestations.⁶ Previous studies show a rare presence of homozygous mutations in the *GJB2* gene in Mexican population.^{7,8}

The *GJB2* gene encodes the gap junction beta-2 protein connexin 26; six connexin molecules form connexons between adjacent cells, responsible for the exchange of ions and molecules among contiguous cells.⁹ Connexins are present in connective and epithelial tissues of the cochlea, which primordial function is to maintain the normal audition.¹⁰

Improvement of the technology plays an important role in the diagnosis of different pathologies, whole exome sequencing (WES) represents an important tool in the identification of the genetic etiology in individuals with SNHL. In the present study we describe a family with SNHL and homozygous mutation in the *GJB2* gene detected by WES.

Patient and methods

The proband was a 21-years-old man with profound congenital deafness in both ears. He was the first product of healthy and non-consanguineous parents. He has a 19-years-old sister with congenital deafness in both ears, as well (Figure 1). Family history was negative for intellectual disability or congenital malformations. No history of prenatal exposure to teratogens, maternal illness or use of aminoglycoside antibiotics was recorded. His mother had an




Figure 1 Pedigree of the family.




Figure 2 Electropherograms showing the mutation in the *GJB2* gene (c.del35G) in the proband (homozygous), parents (heterozygous) and normal control.

uneventful pregnancy with a vaginal delivery at 40 weeks of gestation. His psychomotor development was normal. There was no evidence of syndromic deafness. Computed tomography of the skull showed normal structures. Thyroid parameters were normal. Informed consent was obtained from the patients to participate in the study.

Whole exome sequencing (WES)

Two µg of genomic DNA was submitted to Axeq Technologies for human exome capture using SureSelect human All Exon kit (Agilent Technologies). Axeq Technologies performed sample validation, library preparation, exon enrichment, clustering and sequencing using illumina HiSeq 2000 Sequencer. Approximately, 63,000,000 reads of an average size of 107 bp per sample were returned to us as two fastq files (one file per orientation). Each pair of fastq files were aligned to the human genome (hg19) using Novoalign (<http://www.novocraft.com>). Common variants were removed using allele frequency information from the NHLBI ESP (<http://evs.gs.washington.edu/EVS>). The VCF files were annotated using snpEff (snpeff.sourceforge.net).

Mutation analysis

To confirm the WES findings, DNA sequencing analysis of the *GJB2* gene was performed through Sanger method. The coding exon and flanking intronic regions of the *GJB2* gene were amplified by PCR. All the PCR products were directly sequenced. DNA was analyzed on an ABI 3730XL automated sequencer (Applied Biosystems, Inc., Foster City, CA, USA). The *GJB2* gene was sequenced in all family members and in 100 normal controls to discard polymorphisms.

Results and discussion

From the initial raw variant file (vcf) contained more than 150,000 putative variants, we isolated the high quality, non-synonymous putative coding variants with our quality

filter in conjunction with the annotation tags generated by snpEff (Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, et al. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. Fly (Austin) 2012;6:80–92.). After exploring potential genes involved in NSHL, we found the homozygous mutation c.del35G in *GJB2*. This finding was confirmed through Sanger method (Fig. 2). Audiology studies revealed profound neurosensory hypoacusis in the affected sibs of the family.

Recently, several studies have been performed in the *GJB2* gene that encodes Cx26 connexin. *GJB2* mutations produce several degrees of hearing impairment even in the same family, demonstrating the clinical heterogeneity observed in SNHL. The genetic analysis of the *GJB2* gene (WES and DNA sequencing) in our family with SNHL identified the homozygous c.35delG mutation, this molecular affection was present in the sibs and absent in the normal controls. Both parents were heterozygous for this molecular defect.

Whereas previous studies reveal a very low frequency of this type of mutation in Mexican population, this molecular affection is the most prevalent *GJB2* mutation in Caucasian population.^{7,8,11} Conversely, analysis of the prevalence of *GJB2* mutations in eastern populations shows that c.235delC and not c.35delG is the most frequent cause of SNHL. This difference in the prevalence of the types of mutations in the *GJB2* gene indicates the influence of ethnic background.

The complexity of the genetic component in SNHL is denoted by the great quantity of genes involved in the pathogenesis of the disease. Currently, more than 80 genes have been associated to SNHL (<http://hereditaryhearingloss.org/>), which remarks the difficult to identify the molecular alteration in these patients. Even more, the great variability of the gene affections in each population makes no possible to establish a specific panel to identify the gene and the mutation involved in the molecular etiology of the SNHL.

In conclusion, we describe a Mexican family with SNHL and homozygous mutation c.35delG in the *GJB2* gene. The new techniques of the genomics era (such as whole genome

sequencing, whole exome sequencing or targeted next generation sequencing) have an important role in research or diagnostic areas; they represent excellent tools in the detection of molecular etiology as in SNHL due to its high genetic and clinical heterogeneity.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflict of interest

The authors declare that they have no conflict of interests.

References

1. Atik T, Bademci G, Diaz-Horta O, et al. Whole-exome sequencing and its impact in hereditary hearing loss. *Genet Res.* 2015;97:e4.
2. Rabbani B, Tekin M, Mahdieu N. The promise of whole-exome sequencing in medical genetics. *J Hum Genet.* 2014;59:5–15.
3. Hardisty R, Mburu P, Brown S. ENU mutagenesis and the search for deafness genes. *Br J Audiol.* 1999;33:279–84.
4. Scott D, Krat M, Carmi R, et al. Identification of mutations in the connexin 26 gene that cause autosomal recessive nonsyndromic hearing loss, *Hum. Mutat.* 1998;11:387–94.
5. Oguchi T, Ohtsuka AA, Hashimoto S, et al. Clinical features of patients with GJB2 (connexin 26) mutations: severity of hearing loss is correlated with genotypes and protein expression patterns. *J Hum Genet.* 2005;50:76–83.
6. Snoeckx RL, Huygen PL, Feldmann D, et al. GJB2 mutations and degree of hearing loss: a multicenter study. *Am J Hum Genet.* 2005;77:945–57.
7. Loeza-Becerra F, Rivera-Vega MR, Martínez-Saucedo M, et al. Particular distribution of the GJB2/GJB6 gene mutations in Mexican population with hearing impairment. *Int J Pediatr Otorhinolaryngol.* 2014;78:1057–60.
8. Hernández-Juárez AA, Lugo-Trampe Jde J, Campos-Acevedo LD, et al. GJB2 and GJB6 mutations are an infrequent cause of autosomal-recessive nonsyndromic hearing loss in residents of Mexico. *Int J Pediatr Otorhinolaryngol.* 2014;78:2107–12.
9. Segretain D, Falk MM. Regulation of connexin biosynthesis, assembly, gap junction formation, and removal. *Biochim Biophys Acta.* 2004;1662:3–21.
10. Sabag AD, Dagan O, Avraham KB. Connexins in hearing loss: a comprehensive overview. *J Basic Clin Physiol Pharmacol.* 2005;16:101–16.
11. Marlin S, Feldmann D, Blons H, et al. GJB2 and GJB6 mutations: genotypic and phenotypic correlations in a large cohort of hearingimpaired patients. *Arch Otolaryngol Head Neck Surg.* 2005;131:481–7.