



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Original article

Comparison of sporadic cases of *Salmonella* Typhimurium with other *Salmonella* serotypes in Castellon (Spain): case-case study



Alberto Arnedo-Pena^{a,b,h,*}, Iraya Vivas-Fornas^c, Noemí Meseguer-Ferrer^d, María Dolores Tirado-Balaguer^e, Alberto Yagüe-Muñoz^f, Silvia Herrera-León^g, Susana Sabater-Vidal^e, María Ángeles Romeu-García^d, Ana Vizcaino Batllés^d, Juan Bautista Bellido-Blasco^{d,h}, Rosario Moreno-Muñoz^e

^a Epidemiology Division, Public Health Center, Castellon, Spain

^b Department of Health Sciences, Public University of Navarra, Pamplona, Spain

^c Service of Surgery, Hospital General de Castellón, Castellón, Spain

^d Epidemiology Division, Public Health Center, Castellón, Spain

^e Microbiology Laboratory, Hospital General de Castellón, Castellón, Spain

^f Microbiology Laboratory, Hospital La Plana, Vila-real, Spain

^g Laboratorio de Referencia e Investigación en Enfermedades Bacterianas Transmitidas por agua y alimentos, Centro Nacional de Microbiología, Majadahonda, Madrid, Spain

^h CIBER-Salud Pública Epidemiología, Barcelona, Spain

ARTICLE INFO

Article history:

Received 16 May 2017

Accepted 2 August 2017

Available online 7 October 2017

Keywords:

Salmonella

Gastroenteritis

Salmonella Typhimurium

Children

Swine

Meat products

Dirt

Chickens

Eggs

ABSTRACT

Introduction: Salmonella infections (SI) are common in Spain. The aim of this study was to appraise risk factors and the clinical characteristics of sporadic *Salmonella* Typhimurium infections compared with other sporadic salmonella serotype infections (OSI).

Methods: From September 2014 to August 2015, a case-case study was carried out by the Epidemiology Division of the Public Health Centre of Castellon. Case 1 consisted of patients with sporadic *S. Typhimurium* infections, while case 2 comprised OSI patients, assessed according to the stool cultures analyzed by the Microbiology Laboratories of Hospital General de Castellón and Hospital de La Plana in Vila-real. Patients from detected outbreaks were not included. The salmonella serotype was identified by the National Centre of Microbiology (Madrid).

Results: The total number of SI patients reported was 327,242 of whom were studied (74.0%). 148 patients had sporadic *S. Typhimurium* infection and 64 had OSI, with median ages of 4 and 8.5 years, respectively. Sporadic *S. Typhimurium* infection patients presented more blood in feces and diarrhea episodes. Consumption of pork meat ($OR = 2.22$; 95% CI 1.12–4.43), cold pork meats ($OR = 2.49$; 95% CI 1.32–4.68) and playing in the dirt ($OR = 3.02$; 95% CI 1.55–5.88), were associated with sporadic *S. Typhimurium* infection. In the 0–4 year-old group, the associated factors were consumption of cold pork meats, omelets and female gender. In the 5-year-old and over group, only playing in soil was associated with sporadic *S. Typhimurium* infection.

Conclusions: The consumption of pork and omelets, as well as playing in the dirt, were the main factors associated with infection. Children were most affected by sporadic *S. Typhimurium* infection.

© 2017 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.

Comparación de casos esporádicos de *Salmonella typhimurium* con otros serotipos de *Salmonella* en Castellón (España): un estudio caso-caso

RESUMEN

Introducción: Las infecciones por *Salmonella* son frecuentes en España. El objetivo de este estudio fue estimar factores de riesgo y características de las infecciones esporádicas por *Salmonella typhimurium* (IET), comparadas con infecciones esporádicas por otros serotipos de *Salmonella*.

Palabras clave:

Salmonella

Gastroenteritis

Salmonella typhimurium

* Corresponding author.

E-mail address: albertoarnedopena@gmail.com (A. Arnedo-Pena).

Niños
Cerdo
Productos cárnicos
Tierra
Pollos
Huevos

Métodos: Desde septiembre de 2014 a agosto de 2015 se efectuó un estudio caso-caso por la Sección de Epidemiología del Centro de Salud Pública de Castellón. El caso 1 fueron los pacientes con IET y el caso 2 aquellos con infecciones esporádicas por otros serotipos de *Salmonella*, según los coprocultivos realizados por los laboratorios de Microbiología de los hospitales General de Castellón y La Plana de Vila-real, sin incluir los pacientes de brotes detectados. El serotipo de *Salmonella* fue identificado en el Centro Nacional de Microbiología (Madrid).

Resultados: Se notificaron 327 pacientes con infección por *Salmonella*, de los que 242 fueron estudiados (74,0%). Ciento cuarenta y ocho pacientes tenían IET y 64, infecciones esporádicas por otros serotipos de *Salmonella*, con una mediana de edad de 4 y 8,5 años, respectivamente. Los pacientes IET presentaron más episodios diarreicos y sangre en las heces. El consumo de carne de cerdo (OR 2,22; IC 95% 1,12-4,43), fiambres de cerdo (OR 2,49; IC 95% 1,32-4,68) y jugar en la tierra (OR 3,02; IC 95% 1,55-5,88) fueron asociados con IET. En el grupo de 0-4 años, los factores asociados fueron consumo de fiambres de cerdo, tortillas y ser mujer. En el grupo de 5 y más años, solo jugar en la tierra fue asociado con IET.

Conclusiones: El consumo de productos porcinos y tortillas, así como jugar en la tierra fueron los factores asociados, y los niños, los más afectados.

© 2017 Elsevier España, S.L.U. y Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica.
Todos los derechos reservados.

Introduction

Sporadic cases and outbreaks of *Salmonella* infections continue to be frequent in Spain.¹ During 2014, 241 food-borne outbreaks (FBO) of *Salmonella* with 1681 cases were notified in Spain, 10 FBO with 242 cases in the Valencia Community (VC), and 3 FBO with 25 cases in Castellon Health Department.² The incidence rates were 3,6, 4,8 and 5,2 per 100,000 inhabitants, respectively. However, in 2014 the numbers of notified cases with positive *Salmonella* culture were 7288 in Spain and 291 in Castellon with rates of 15,6 and 60,9 per 100,000 inhabitants, respectively.^{2,3} These data reflect only limited morbidity of salmonellosis considering the iceberg phenomenon of mandatory notified diseases. In addition, episodes of community FBO *Salmonella* continued to occur in Castellon.⁴ From studies of *Salmonella* epidemiology in Castellon in previous years, a change can now be detected with *S. Typhimurium* currently being the most common serotype isolated in culture compared to *Salmonella Enteritidis*.^{5,6}

The transmission of *Salmonella* follows a fecal-oral route and the consumption of foods, more rarely water, contaminated with the bacteria, is the most frequent vehicle of infection. Knowledge of the epidemiologic and the microbiological characteristics of the *Salmonella* species in the territory is the first step in preventing these infections. In this context, several epidemiologic designs could be used such as the case-case design,⁷ and the study of isolated *Salmonella* by serotype, phage-type, and molecular methods including pulsed-field gel electrophoresis and whole-genome sequencing may provide a deeper knowledge.

The aim of this study was to find associated factors and clinical characteristics of sporadic *S. Typhimurium* infections compare with other sporadic *Salmonella* serotypes infections using a case-case design in order to establish more appropriate measures of prevention.

Material and methods

In 2014, mandatory reporting of *Salmonella* was introduced in the VC, and results of positive *Salmonella* culture from the microbiology laboratories of public hospitals (MLPH) were sent to public health center epidemiology units. All positive *Salmonella* culture patients were then interviewed about the characteristics of the disease and potential risk factors as part of the epidemiologic surveillance of this infection. In addition, the MLPH sent *Salmonella* isolated in the culture to the *Salmonella* National Reference Laboratory (SNRL) of the Instituto Carlos III in Majadahonda (Madrid) to identify the serotyping and phage-typing.

From September 2014 to August 2015 a case-case study was carried out with all positive *Salmonella* cultures. These cases were divided in two groups: case1 as positive *S. Typhimurium* status and case2 as positive other *Salmonella* serotypes status. To obtain enhanced comparisons of patients only culture of feces was considered, and patients from detected outbreaks were not included. Staff at the Epidemiologic Division of the Public Health Center in Castellon contacted physicians to ask about the culture and request authorization to interview the patients. The patients or their parents were then contacted by telephone and their oral informed consent was obtained. The ethic committee of the Hospital General of Castellon approved this study.

The interview was the same for case 1 and case 2 patients and was carried out at which a mean of 61.1 ± 38.9 days from the *Salmonella* infections notification. The questionnaire asked about the clinical characteristics of the disease, its evolution, and risk factors including habitual food consumption, focusing on meat, chicken eggs and pork products, food preparation, hygiene practices in the kitchen, taking antibiotics, contacts with animals, among others.

Microbiological methods

The positive *Salmonella* culture arrived from the Microbiologic Laboratories of the Hospital General in Castellon and Hospital La Plana in Vila-real. In both Hospitals stool samples were inoculated on usual media. In the Microbiology Laboratory of the Hospital La Plana screening of suspected colonies was performed with an agglutination test (Difco *Salmonella* O Antiserum Poly A-I & Vi, BD). Strains with positive agglutination were identified with the automated system Walk Away (Beckman Coulter Inc.). In the Microbiology Laboratory of the Hospital General suspected colonies were directly identified with the automated system VITEK 2 (Biomérieux). They serotyped *Salmonella* isolates by agglutination, according to the Kauffmann-White scheme, using polyvalent and monovalent antisera to somatic and to flagellar antigens (Bio-Rad; Statens Serum Institute). It was possible to identify these different serotypes: *Salmonella Typhimurium*, *Salmonella Enteritidis*, *Salmonella Typhi*, *Salmonella Hadar* and *Salmonella Virchow*. The rest of *Salmonella* isolated were only characterized as group B, C and D or as *Salmonella* spp.

All *Salmonella* strains were also referred to the SNRL at Majadahonda (Madrid) for its phage typing and to complete its serotyping. *Salmonella* serotype was determined by the slide agglutination method using commercial antisera (Bio-Rad) and following the

Kauffman-White scheme for serotype attribution. Phage typing for serotypes Typhimurium, biphasic and monophasic variants, Enteritidis, Virchow, Hadar and Typhi was performed according to the international schemes with phages and interpreting criteria provided by the Health Protection Agency (HPA, formerly Public Health Laboratory Service, Colindale in London, England).

Statistical analysis

Two analyses were performed: a general analysis with all the cases of *Salmonella* infections, and other analysis according two age groups, considering one group from 0 to 4 years old and the other of 5 years old and over. These groups may be considered appropriate since more than 50% of *S. Typhimurium* cases were in children under the age of 5 years old.

The comparison between case 1 and case 2 was done by Chi2 and Fisher tests for the qualitative variables and by the Kruskal-Wallis test for the quantitative variables. Logistic regression was used to find factors associated with the *Salmonella* infections compare by odds ratio (OR) with a 95% confidence interval (CI). Multivariate logistic regression was used to analyze the associations with several risk factors. The factors were included in the initial models if they were associated with a significant *p* value <0.10. The final models presented a good of fit, *p* value >0.05. All calculations were performed with Stata® version 14.

Results

In the period of the study 327 positive *Salmonella* cultures were notified and 242 cases were included (74.0%). Of the 21 *Salmonella* isolated from outbreaks, 17 were *S. Typhimurium* and the remaining 4 other serotypes. From feces culture, 212 *Salmonella* were isolated, 148 (69.3%) *S. Typhimurium* and 64 (30.7%) other *Salmonella* serotypes. Nine *Salmonella* were isolated from other clinical samples (blood, urine, and skin abscess).

Table 1 presents the characteristics of these case 1 and case 2. The most affected age group was 1–4 years old for both case 1 and case 2. In case 1 with age median of 4.0 years, the second ranks more affected was 5–9 years old. In the case 2 with age median 8.5 years, the second group most affected was 35–65 years old. No significant differences were observed by gender, or country of origin. Other *Salmonella* serotypes isolated with more than one positive culture included the following: Derby (2), Rissen (7), Hadar (3), Newport (3), Mikawasima (5), Virchow (5), Goldcoast (4), Mbandaka (2), Bredeney (2) and Braenderup (2). The most frequent *S. Typhimurium* phage-types were DT104 (25), U311 (14), DT138 (12), RDNC (12), DT93 (11), and untypable (6).

The patients suffered acute gastroenteritis and their clinical symptoms are presented in **Table 2**. The group case 1 suffered more significant blood in the feces and higher numbers of diarrhea episodes by day than group case 2. However, nausea was more frequent in group case 2. The significance of these symptoms held when age was adjusted for. No other significant differences were observed. The hospital admissions in the case 1 and case 2 groups were 16.2% and 14.1%, respectively. Clinical evolution was good for all the patients.

Several factors were associated with *S. Typhimurium* infection in the logistic regression analysis (**Table 3**), namely, age, consumption of some specific foods once per week or more, including pork products, pork meat, pork sausages, cold pork meats, and dried pork sausages, minced meat, and chicken egg omelets, attending day-care/school, and playing in soil. Common risk factors of salmonellosis such as taking antibiotics 15 days before onset or contact with a patient with diarrhea were not associated.

Table 1
Characteristics of sporadic *S. Typhimurium* and other *Salmonella* serotypes patients.

Variables	<i>S. Typhimurium</i> <i>n</i> =148 <i>N</i> (%)	Other <i>Salmonella</i> serotypes <i>n</i> =64 <i>N</i> (%)	<i>p</i> -value
Age (years) mean ± SD	12.1 ± 20.6	22.8 ± 27.1	0.003
Age (years) median-range	4.0 (0.08–89)	8.5 (0.01–91)	
Age groups			
<1 year	5 (3.4)	8 (12.5)	
1–4 years	75 (50.7)	17 (26.6)	
5–9 years	32 (21.6)	8 (12.5)	
10–14 years	12 (8.1)	6 (9.4)	
15–34 years	5 (3.4)	6 (9.4)	
35–64 years	9 (6.1)	11 (17.2)	
65 years and more	10 (6.8)	8 (12.5)	0.001
Female	81 (54.7)	32 (50.0)	0.551
Country of origin			
Spain	135 (91.2)	59 (92.2)	
Romania	8 (5.4)	0 (0)	
Maghreb	2 (1.4)	3 (4.7)	
Other	3 (2.0)	2 (3.1)	0.075
Residence in Castellon department	85 (57.4)	34 (53.1)	0.651
Infection onset Summer season	55 (37.2)	22 (34.4)	0.757
Salmonella serotype			
<i>S. Typhimurium</i>	148		
<i>S. Enteritidis</i>		18 (28.1)	
Other <i>Salmonella</i> serotypes ^a		42 (65.6)	
<i>Salmonella</i> spp		4 (6.3)	

^a Derby, Rissen, Hadar, Newport, Mikawasima, Virchow, etc.

Table 2
Comparison of the clinical characteristic of sporadic *S. Typhimurium* and other *Salmonella* serotypes patients.

Variables	<i>S. Typhimurium</i> <i>n</i> =148 <i>N</i> (%)	Other <i>Salmonella</i> serotypes <i>n</i> =64 <i>N</i> (%)	<i>p</i> -value
Symptoms			
Abdominal cramps	103 (69.6)	38 (59.3)	0.243
Nauseas	33 (22.9)	24 (37.5)	0.015
Vomits	56 (37.8)	30 (46.9)	0.233
Diarrhea	145 (98.0)	61 (95.3)	0.014
No. diarrhea episodes by day ± SD	10.3±6.0	8.4±5.4	0.014
Blood in diarrhea feces	71 (48.0)	14 (21.9)	0.001
Fever ≥38 °C	103 (69.6)	36 (56.3)	0.060
Headache	34 (23.0)	16 (25.0)	0.750
Other symptoms	41 (27.7)	22 (34.4)	0.262
Hospital admission	24 (16.2)	9 (14.1)	0.691
Duration illness ± SD	10.2±7.5	13.3±14.3	0.302

SD= standard deviation.

Several logistic regression models were implemented from the associated factors with a significance level of *p*<0.10. In model 1 (**Table 4**), consumption of pork meat (OR=2.22 (95% CI 1.12–4.43)), cold pork meats (OR=2.49 (95% CI 1.32–4.68)), and playing in soil (OR=3.02 (95% CI 1.55–5.88)), were associated with *S. Typhimurium* infection. In model 2, pork products, considered as positive response answer to consumption of pork meat, pork sausages, dried pork sausages and cold pork meats (OR=5.03 (95% CI 2.05–12.35)), and playing in soil (2.23 (95% CI 1.09–4.57)) were associated with *S. Typhimurium*, adjusted by age (OR=0.98 (95% CI 0.97–0.99)).

Table 5 shows the 0–4 years old group and 5 years and over group considering the *S. Typhimurium* and other *Salmonella* serotypes in each group and the associated factors. In the 0–4

Table 3Comparison of *S. Typhimurium* and other *Salmonella* by consumption of different foods (frequency once per week or more) and factors.

Variables	<i>S. Typhimurium</i> N = 148 (%)	Other <i>Salmonella</i> serotypes N = 64 (%)	OR 95% CI	p value
Age (years) ± SD	12.1 ± 20.6	22.8 ± 27.1	0.98 (0.97–0.99)	0.003
Female	81 (54.7)	32 (50.0)	1.21 (0.67–2.17)	0.527
Country of origin				
Spain	135 (91.2)	59 (92.2)	0.88 (0.30–2.58)	0.816
Rumania	8 (5.4)	0 (0.0)	4.98 (0.75–∞)	0.106
Maghreb	2 (1.4)	3 (4.7)	0.29 (0.05–1.71)	0.167
Consumption frequency once by week or more				
Chicken	145 (98.0)	60 (93.8)	3.22 (0.70–14.23)	0.133
Turkey	53 (35.8)	15 (23.4)	1.87 (0.93–3.66)	0.079
Veal	89 (60.1.9)	38 (59.4)	1.03 (0.57–1.88)	0.917
Lamb	55 (37.2)	16 (25.0)	1.77 (0.92–3.42)	0.087
Pork meat	120 (81.1)	40 (62.5)	2.57 (1.34–4.73)	0.005
Pork sausages	94 (63.5)	30 (46.9)	1.97 (1.09–3.57)	0.025
Cold pork meats	92 (62.2)	26 (40.6)	2.64 (1.44–4.82)	0.002
Dried pork sausages	59 (39.9)	12 (18.9)	2.87 (1.49–5.84)	0.004
Pork products ^a	138 (93.2)	48 (75.0)	4.60 (1.96–10.82)	0.000
Minced meat	115 (77.7)	40 (62.5)	2.04 (1.11–3.95)	0.023
Fish	130 (87.8)	59 (92.2)	0.61 (0.22–1.73)	0.354
Chicken egg mayonnaise	30 (20.3)	12 (18.8)	1.10 (0.52–2.32)	0.799
Chicken egg omelets	133 (90.0)	49 (76.6)	2.71 (1.24–5.96)	0.013
Other foods with eggs	20 (13.5)	7 (10.9)	1.27 (0.51–3.18)	0.606
Drinking unsafe water	19 (12.8)	11 (17.2)	0.61 (0.28–1.37)	0.406
Raw meat in home kitchen	121 (81.8)	58 (90.6)	0.46 (0.18–1.18)	0.108
Handling raw meat	56 (37.8)	27 (42.2)	0.83 (0.46–1.51)	0.552
Restaurant visit 15 days before onset	59 (39.9)	17 (26.6)	1.83 (0.96–3.49)	0.066
Attend day-care/school	76 (51.4)	19 (29.7)	2.49 (1.34–4.67)	0.004
Play in soil	77 (52.0)	17 (26.6)	3.0 (1.58–5.70)	0.001
Play in places with pigeons	42 (28.4)	18 (28.1)	1.01 (0.53–1.94)	0.970
Animals at home	77 (52.0)	34 (53.1)	0.96 (0.53–1.72)	0.883
Dog, cat, other mammals	60 (40.5)	26 (40.6)	1.00 (0.55–1.81)	0.911
Birds	27 (18.2)	13 (20.3)	0.87 (0.41–1.83)	0.724
Reptiles	5 (3.4)	3 (4.7)	0.71 (0.16–3.07)	0.648
Take antibiotics 15 days before	30 (20.3)	15 (23.4)	0.83 (0.41–1.68)	0.605
Contact with diarrhea patients	19 (12.8)	7 (10.9)	1.20 (0.48–3.01)	0.699

OR = odds ratio; CI = confidence interval; SD = standard deviation.

^a Includes: pork meat, pork sausages, cold pork meats and dry pork sausages.**Table 4**Multivariate logistic regression models: *S. Typhimurium* associated factors.

Variables	Model 1		Model 2	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age (years) ± SD	–	–	0.98 (0.97–0.99)	0.035
Play in soil	3.02 (1.55–5.88)	0.001	2.23 (1.09–4.57)	0.028
Consumption frequency once per week or more				
Pork meat	2.22 (1.12–4.43)	0.023		
Cold pork meats	2.49 (1.32–4.68)	0.005		
Pork products ^a	–		5.03 (2.05–12.35)	0.000

OR = odds ratio; CI = confidence interval; SD = standard deviation.

^a Includes: pork meat, sausages, cold pork meats and dried sausages.

years old group, these factors were female gender, attending day-care/school, and consumptions once per week or more of pork meat, cold pork meats, dried pork sausages, pork products, minced meat, and chicken egg omelets. In the 5 years and over group, the associated factors were age and playing in soil. Table 6 shows the results of the logistic regression. In the 0–4 years old group, the associated factors were consumption of cold pork meats ($OR = 4.25$ (95% CI 1.39–12.96)) and chicken egg omelets ($OR = 5.00$ (95% CI 1.63–15.32)) and female gender ($OR = 4.13$ (95% CI 1.40–12.24)). In addition, the consumption of pork products ($OR = 12.84$ (95% CI 3.76–43.91)) and female gender ($OR = 4.09$ (95% CI 1.34–12.51)) were associated with *S. Typhimurium*. In the 5 years and over group, only playing in soil ($OR = 3.00$ (95% CI 1.10–8.17)) was associated with *S. Typhimurium*.

Discussion

The results of this study suggest that the consumption of pork products and playing in soil were associated with *S. Typhimurium* infections. However, when two age groups were considered, associated factors of *S. Typhimurium* infection in the 0–4 years old group were female gender, and consumption of pork products and chicken egg omelets. In the 5 years and over group, playing in soil was the only associated factor in the logistic model. In addition, some differences in clinical symptoms between *Salmonella* infections case 1 and case 2 groups could be highlighted.

In relation to intestinal symptoms, the patients with *S. Typhimurium* infections had more severe clinical manifestations than the patients with other *Salmonella* serotypes. *S. Typhimurium* produces inflammatory diarrhea with metabolic changes that affect the intestinal microbiota and the type III secretions systems induce the invasion, intracellular survival, and the immune response of the patient.^{8,9} Young children and old people had symptomatic infections, whereas other adults were asymptomatic or had few gastroenteritis symptoms. Salmonellosis may be considered a disease of children, since the 76% of *S. Typhimurium* cases were in patients under the age of 10; this percentage was 52% for other *Salmonella* serotypes. Higher incidence in children and old people has been observed in previous studies.⁵ Morbidity from positive *Salmonella* culture account for a small proportion of the total morbidity caused by *Salmonella* and only potential risk factors associated with these patients could be studied.

Table 5

S. Typhimurium associated factors for two age groups: 0–4 years old and 5 years and over.

Variables	0–4 years-old group n = 105		5 years and over group n = 107	
	S. Typhimurium n = 80 Other <i>Salmonella</i> serotypes n = 25		S. Typhimurium n = 68 Other <i>Salmonella</i> serotypes n = 39	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age (years) ± SD	1.54 (0.95–2.48)	0.078	0.98 (0.97–0.99)	0.024
Female gender	2.81 (1.11–7.14)	0.030	0.62 (0.28–1.37)	0.237
Attend daycare/school	3.35 (1.29–8.71)	0.013	1.68 (0.72–3.92)	0.234
Play in soil	2.50 (1.00–6.24)	0.050	3.00 (1.10–8.17)	0.032
<i>Consumption frequency once per week or more</i>				
Pork meat	5.17 (1.98–13.45)	0.001	1.74 (0.64–4.74)	0.279
Cold pork meats	4.51 (1.63–12.51)	0.004	2.20 (0.98–4.94)	0.056
Dried pork sausages	3.17 (1.15–8.76)	0.026	2.13 (0.77–5.91)	0.145
Pork products ^a	9.63 (3.19–29.01)	0.000	2.48 (0.52–11.69)	0.252
Minced meat	5.09 (1.95–13.31)	0.001	1.03 (0.42–2.56)	0.941
Chicken egg omelets	5.23 (1.93–14.16)	0.001	1.81 (0.35–9.41)	0.483

OR = odds ratio; CI = confidence interval; SD = standard deviation.

^a Includes: pork meat, pork sausages, cold pork meats and dried pork sausages.**Table 6**

Multivariate logistic regression models: S. Typhimurium associated factors by age groups; 0–4 years old and 5 years and over.

Variables	0–4 years old group		5 years and over group	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Female	4.13 (1.40–12.24)	0.01	—	—
Play in soil	—	—	3.00 (1.10–8.17)	0.032
<i>Consumption frequency once per week or more</i>				
Cold pork meats	4.25 (1.39–12.96)	0.011	—	—
Chicken egg omelets	5.00 (1.63–15.32)	0.005	—	—

OR = odds ratio; CI = confidence interval; SD = standard deviation.

Exposure to pork products, and playing in soil were higher in *S. Typhimurium* patients than other *Salmonella* patients, suggesting that they could be risk factors for the general population.¹⁰ Female gender and consumption of chicken egg omelets were associated with *S. Typhimurium* in the 0–4 years old group only, and it could be considered more hypothetical. In Castellon, the risk factors of sporadic *Salmonella* infection in 1–7 years old children were consumption of minced meat and contacts with animals in patients with diarrhea, and in 0–2 years-old children minced meat or poultry in the kitchen were associated with salmonellosis in a case-control study.^{11,12} In our study, minced meat was a risk factor in the 0–4 years old group but it lost significance in the logistic models. In Salamanca,¹³ a retrospective study of *Salmonella* infections found that *S. Typhimurium* was the most frequent serotype isolated, the patients had 7 years old as median of age, and consumption of pork products is suggested as a risk factor. In a case-control study in China,¹⁴ *S. Typhimurium* was more frequent than *S. Enteritidis* and pork and poultry products were risk factors. In a case-case in Germany, Ziehm and co-authors¹⁵ found that *S. Typhimurium* was associated with consumption of raw ground pork (OR = 2.8 95% CI 1.99–3.94) and rodents keeping (OR = 1.4 95% CI 1.05–1.89). This study included 867 *S. Typhimurium* versus 1034 other *Salmonella* serotypes patients.

Many animals are reservoirs of *S. Typhimurium*, including swine, cattle, chicken, birds, and reptiles and this serotype is considered ubiquitous as a result exposure is very varied. The pork products associated were pork meat and cold pork meats. The first product is eaten cooked and the second, which includes salami, chorizo, and cured dried pork sausage is eaten raw. In Spain, several studies of *Salmonella* infections have estimated high prevalence, 43.1% in swine farms,¹⁶ 31% in pigs¹⁷ and 10.8%

in processing plants.¹⁸ In addition, *Salmonella* was isolated in 23.7% of raw material for the production of fuet, a fermented pork sausage,¹⁹ 11.1% of cured dried sausage samples, a ready-to-eat food,²⁰ and 8.3% of pork meat in retail outlets.²¹ On the other hand, *S. Typhimurium* infections have been associated with chicken egg consumption in Spain²² and other countries such as Australia.²³ Another risk factor was playing in soil, which may be contaminated by animal feces. These risk factors have been indicated in studies of salmonellosis in children.²⁴ Female gender was associated with *S. Typhimurium* in the 0–4 years old group but not in the group 5 years and over more; its OR is elevated and it may indicate some gender differences in food preferences and unknown factors.

The study includes a high percentage of *Salmonella* positive culture cases and the patients were interviewed in acceptable period after the infection onset, which likely reduced interview and recall biases. The high sample size had a sufficient power. Foods consumed were identified as habitual preference.

This study has some limitations. First, there is a potential selection bias arising from the study of patients with *Salmonella* feces culture, since they may have more serious symptoms or other associated pathologies. Similarly, very young or very old patients could have some differences from the general population. Second, *S. Typhimurium* group had several phage-types that could differ in their reservoirs or transmission routes; in addition, other *Salmonella* group included serotypes associated with pork products²⁵ such as *S. Derby* and *S. Rissen*, these potential biases could have a deviation toward the null hypotheses. Third, in the case-case design, the common risk factors of *Salmonella* infections such as antibiotic use, immune-deficiencies, and other pathologies are difficult to detect. Fourth, the questionnaire focuses on foods consumed but other risk factors such as travels could also play some role. Fifth, the 0–4 years old group is homogenous (49.5% of total) and yielded obtain more significant associations than the 5 years and over group (51.5% of total), which had a higher age dispersion. Sixth, the absence of molecular methods to study *S. Typhimurium* such as pulsed-field gel electrophoresis hampers a more precise epidemiology for finding outbreaks and it may be case that some of what were considered to be sporadic cases were actually from unknown outbreaks.²⁶

Case-case methodology has been used to study salmonellosis outbreaks and presents some advantages over the case-control design. Information bias may be lower but selection bias may increase. The use of case-case design in parallel with epidemiologic surveillance may be indicated; since it reduces the time needed to

find appropriate controls for the case, and may allow more adequate measures to stop *Salmonella* infections.

Increased health education about the consumption of pork products may be required considering that the potential risk of salmonellosis associated with pork products is little known in the Spanish population and consumption of this food is very common; Spain was the second producer of pig meat in Europe in 2014 and the third pork consumer in 2015 with 54.4 kg per capita/year.^{27,28} The risk of salmonellosis by chicken eggs consumption is present and it need more specific actions. In addition, other measures include improvements of food hygiene in the swine industry, a national control program for *Salmonella* in pigs and pork products²⁹ and the appropriate maintenance of children playgrounds.³⁰ Children are the most affected by STI; pork products and chicken egg omelets consumption and playing in soil are the main associated factors.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgment

The authors thank all the patients for their cooperation, which has made this study possible.

References

- Sala-Farré MR, Osorio-Sánchez D, Arias-Varela C, Simó-Sanahuja M, Recasens-Recasens A, Pérez-Jové J. Aspectos epidemiológicos y carga asistencial de gastroenteritis aguda por *Campylobacter* y *Salmonella*. Med Clin (Barc). 2015;145:294–7.
- Centro Nacional de Epidemiología. Resultados de la vigilancia epidemiológica de las enfermedades transmisibles e Informes del sistema de información microbiológica. Año 2014. Instituto de Salud Carlos III. Ministerio de Economía, Industria y Competitividad. Madrid. <http://www.isciii.es/ISCIII/es/contenidos/fd-el-instituto/publicaciones.shtml> [accessed 10.03.17].
- Sección de Epidemiología. Centro de Salud Pública. Memoria anual 2014. Castellón: Generalitat Valenciana; 2015.
- Arnedo-Peña A, Sabater-Vidal S, Herrera-León S, Bellido-Blasco JB, Silvestre-Silvestre E, Meseguer-Ferrer N, et al. An outbreak of monophasic and biphasic *Salmonella* Typhimurium and *Salmonella* Derby associated with the consumption of dried pork sausage in Castellón (Spain). Enferm Infect Microbiol Clin. 2016;34:544–50.
- Bellido-Blasco JB, Celades-Porcar ME, Tirado-Balaguer MD, González-Cano JM, Gil-Ortuño M, Arnedo-Peña A. Estudio de la diarrea infecciosa en Castellón (EDICS): incidencia poblacional de casos esporádicos en 2004 y su comparación con el año 2000. Med Clin (Barc). 2006;127:448–50.
- Tirado MD, Moreno RM, Celades ME, Bellido-Blasco J, Pardo FJ. Evolución de los serotipos, fagotipos y resistencia a antimicrobianos de *Salmonella* spp en el departamento 02 de la provincia de Castellón, España (2000–2006). Rev Chilena Infectol. 2009;26:520–7.
- McCarthy N, Giesecke J. Case-case comparison to study causation of common infectious diseases. Int J Epidemiol. 1999;28:764–8.
- Patel S, McCormick BA. Mucosal inflammatory response to *Salmonella* Typhimurium. Front Immunol. 2014;5:311.
- LaRock DL, Chaudhary A, Miller S. *Salmonellae* interactions with host processes. Nat Rev Microbiol. 2015;13:191–205.
- Pogreba-Brown K, Ernst K, Harris RB. Case-case methods for studying enteric diseases: a review and approach for standardization. OA Epidemiol. 2014;2:7.
- Bellido-Blasco JB, González-Cano JM, Galiano-Arlandis JV, Bernat S, Arnedo A, González-Morán F. Factores asociados con casos esporádicos de salmonelosis en niños de 1 a 7 años. Estudio de casos y controles. Gac Sanit. 1998;12:118–45.
- Bellido-Blasco JB, González-Cano JM, Galiano-Arlandis JV, Herrero-Carot C, Tirado-Balaguer MD, Arnedo-Peña A, et al. Factores de riesgo de los casos esporádicos de diarrea por *Campylobacter*, *Salmonella* y rotavirus en niños preescolares. An Pediatr (Barc). 2007;66:357–74.
- Cores-Calvo O, Valero-Juan LF, García-Sánchez E, García-Sánchez JE, García-García MI. Cambios en la epidemiología de las gastroenteritis causadas por *Salmonella* durante el periodo 2005–2014 en Salamanca, España. Rev Esp Quimioter. 2016;29:91–8.
- Yang X, Jin K, Yang F, Yuan G, Liu W, Xiang L, et al. Nontyphoidal *Salmonella* gastroenteritis in Baoshan, Shanghai, China, 2010 to 2014: an etiological surveillance and case-control study. J Food Prot. 2017;16:482–7.
- Ziehm D, Dreesman J, Rabsch W, Fruth A, Pulz M, Kreienbrock L, et al. Subtype specific risk factor analyses for sporadic human salmonellosis: a case-case comparison in Lower Saxony, Germany. Int J Hyg Environ Health. 2013;216:428–34.
- García-Feliz C. Salmonelosis porcina en España: prevalencia, factores de riesgo y resistencias antimicrobianas. Departamento de Sanidad Animal, Universidad de León; 2011. Tesis doctoral.
- Vico JP, Roll I, Garrido V, San Román B, Grilló MJ, Mainar-Jaime RC. Salmonellosis in finishing pigs in Spain: prevalence, antimicrobial agent susceptibilities, and risk factor analysis. J Food Prot. 2011;74:1070–8.
- Hernández M, Gómez-Laguna J, Luque I, Herrera-León S, Maldonado A, Reguillo L, et al. *Salmonella* prevalence and characterization in a free-range pig processing plant: tracking in trucks, lairage, slaughter line and quartering. Int J Food Microbiol. 2013;162:48–54.
- Martín B, Garriga M, Aymerich T. Prevalence of *Salmonella* spp. and *Listeria* monocytogenes at small-scale Spanish factories producing traditional fermented sausages. J Food Prot. 2011;74:812–5.
- Cabedo L, Picart i Barrot L, Teixido i Canelles A. Prevalence of *Listeria* monocytogenes and *Salmonella* in ready-to-eat food in Catalonia, Spain. J Food Prot. 2008;71:855–9.
- Valero A, Hernández M, De Cesare A, Manfreda G, García-Gimeno RM, González-García P, et al. Probabilistic approach for determining *Salmonella* spp and *L. monocytogenes* concentration in pork from presence/absence microbiological data. Int J Food Microbiol. 2014;184:60–3.
- Arnedo A, Bellido JB, Pac MR, Criado J, Usera MA, Mesanza I, et al. Brotes epidémicos de salmonelosis causados por consumo de huevos. Enferm Infect Microbiol Clin. 1998;16:408–12.
- Moffatt CR, Musto J, Pingault N, Miller M, Stafford R, Gregory J, et al. *Salmonella* Typhimurium and outbreaks of egg-associated disease in Australia, 2001–2011. Foodborne Pathog Dis. 2016;13:379–85.
- Doorduyn Y, Van der Brandhof WE, Van Duynhoven YT, Wannet WJB. Risk factors for *Salmonella* Enteritidis and Typhimurium (DT104 and non.DT104) infections in the Netherlands: predominant roles for raw eggs in Enteritidis and sandboxes in Typhimurium infections. Epidemiol Infect. 2006;134:617–26.
- Bonardi S. *Salmonella* in the pork production chain and its impact on human health in the European Union. Epidemiol Infect. 2017;28:1–14.
- Arnedo-Peña A, Bellido-Blasco JB, Romeu-García MA, Meseguer-Ferrer N. Detection of foodborne *Salmonella* Typhimurium outbreaks. Enferm Infect Microbiol Clin. 2017 Feb 22, <http://dx.doi.org/10.1016/j.eimc.2017.01.001>.
- European Commission. http://ec.europa.eu/eurostat/statistics-explained/index.php/Meat_production_statistics [accessed 10.05.17].
- Agriculture and Horticulture Development Board. <http://pork.ahdb.org.uk/prices-stats/consumption/eu-per-capita-consumption/> [accessed 10.05.17].
- Arguello H. Tesis doctoral Salmonelosis porcina en España: factores de riesgo en reproductores, estrategias de control de cerdos de cebo y la importancia del sacrificio. Departamento de Sanidad Animal, Universidad de León; 2013.
- Olmedo-Lucerón C, Pérez-Meixeira A, Abad-Sanz I, Cid-Delyeto V, Herrera-León S, Gutierrez-Ruiz L. An outbreak of *Salmonella* Typhimurium associated with playground sand in a preschool setting-Madrid, Spain, September-October 2016. MMWR. 2017;66:256–7.