



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Original article

Epidemiology of dermatophytic infections between 2008 and 2017 in Barcelona, Spain



Adrián Antuori^{a,*}, Gema Fernández^a, Anabel Fernández^a, Mireya Alcaide^a, Aram Boada^b, María Isabel Bielsa^b, Nona Romaní^a, Lurdes Matas^{a,c}

^a Microbiology Department, Clinical Laboratory North Metropolitan Area, University Hospital “Germans Trias i Pujol”, Departament of Genetics and Microbiology, Autonomous University of Barcelona, Badalona, Spain

^b Dermatology and Venereology Service, Germans Trias i Pujol University Hospital, Badalona, Spain

^c CIBER Epidemiología y Salud Pública, Spain

ARTICLE INFO

Article history:

Received 20 November 2018

Accepted 20 February 2019

Available online 7 May 2019

Keywords:

Dermatophyte infections

Epidemiology

Spain

Mycosis

Trichophyton

Tinea

ABSTRACT

Objective: The aim of this study was to evaluate the epidemiological profile of dermatophytoses from 2008 to 2017 in the area of “Barcelonès Nord”, located in north-eastern Spain.

Methods: From 2008 to 2017, 13,419 specimens obtained from patients with suspected superficial mycosis were subjected to direct microscopy and culture. Clinical and sociodemographic data were recorded. Proportions were compared using Chi-square and Fisher’s exact tests. A logistic regression model was used for multivariate analysis.

Results: *Trichophyton rubrum* was the most frequently isolated fungus (76.1%), followed by *Trichophyton mentagrophytes/Trichophyton interdigitale* (11.9%) and *Microsporum canis* (2.9%). Among the population over 15 years of age, *tinea unguium pedum* (40.4%) and *tinea corporis* (29.1%) were the predominant dermatophyte infections. *Tinea capitis* was mostly prevalent (53.6%) among patients up to 15 years of age, followed by *tinea corporis* (21.4%). We observed an increase in non-endemic anthropophilic dermatophytes (*Trichophyton soudanense*, *Microsporum audouinii* and *Trichophyton violaceum*) in the last few years. These species were associated with the population up to 15 years of age ($p < 0.001$), having *tinea capitis* ($p = 0.0017$) and being of African/Hindustani origin ($p < 0.001$).

Conclusions: We confirmed the spread of *T. rubrum* as the predominant dermatophyte in our area and reported an increase in non-endemic anthropophilic dermatophytes in the last few years, especially in the African and Hindustani population up to 15 years of age.

© 2019 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.

Epidemiología de las dermatofitosis entre 2008 y 2017 en Barcelona, España

RESUMEN

Objetivo: La finalidad del presente estudio fue evaluar el perfil epidemiológico de las dermatofitosis entre los años 2008 y 2017 en el área de «Barcelonès Nord» ubicada en el noreste de España.

Métodos: Entre los años 2008 y 2017, 13.419 muestras obtenidas de pacientes con sospecha de micosis superficial fueron remitidas para microscopía directa y cultivo. Se registraron datos clínicos y sociodemográficos. Las proporciones se compararon mediante las pruebas de Chi-cuadrado y Fisher. Para el análisis de variables múltiples, se utilizó un modelo de regresión logística.

Resultados: *Trichophyton rubrum* fue el hongo más frecuentemente aislado (76,1%), seguido de *Trichophyton mentagrophytes/Trichophyton interdigitale* (11,9%) y *Microsporum canis* (2,9%). Entre la población mayor de 15 años, la *tinea unguium pedum* (40,4%) y la *tinea corporis* (29,1%) fueron las dermatofitosis predominantes. La *tinea capitis* prevaleció principalmente (53,6%) entre los pacientes menores de 15 años, seguida de la *tinea corporis* (21,4%). Se percibió un aumento de los dermatofitos antropófilos no

Palabras clave:

Dermatofitosis

Epidemiología

España

Micosis

Trichophyton

Tinea

* Corresponding author.

E-mail address: adrian.antuori@gmail.com (A. Antuori).

endémicos (*Trichophyton soudanense*, *Microsporum audouinii* y *Trichophyton violaceum*) en los últimos años. Estas especies se asociaron con la población menor de 15 años ($p < 0,001$), la presencia de *tinea capitis* ($p = 0,0017$) y el origen africano/indostánico ($p < 0,001$).

Conclusiones: Confirmamos el predominio de *Trichophyton rubrum* como el dermatofito más prevalente en nuestra área, y describimos un aumento en los dermatofitos antropófilos no endémicos en los últimos años, especialmente en población africana e indostana menor de 15 años.

© 2019 Elsevier España, S.L.U. y Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica.

Todos los derechos reservados.

Introduction

Dermatophytes are a group of filamentous fungi able to invade keratinized tissues such as hair, skin and nails, resulting in superficial infections.¹ According to the adaptation of each species to different animals or other ecological reservoirs, they have also been classified into geophilic, zoophilic and anthropophilic species.¹ The identification of current main genera *Epidermophyton*, *Microsporum* and *Trichophyton* are mainly based on the microscopic appearance of the reproductive organs in culture, called conidia (L, s: conidium), as well as in some other secondary structures of the vegetative mycelium (coiled, pectinate or antler-like hyphae, chlamydoconidia or nodular organs).² These fungi represent a frequent reason of consultation, and it is estimated that around 20–25% of the global population are, at some stage, affected by these mycoses.³

The epidemiology of superficial dermatophytoses has changed radically throughout the 20th century in relation to factors such as life conditions, socio-economic status or migration.³ In the 1930s, anthropophilic species such as *Microsporum audouinii* and *Trichophyton schoenleinii* were the principal agents of *tinea capitis* in Europe and America, while *Trichophyton mentagrophytes* was the dermatophyte most frequently isolated in *tinea pedis* and *tinea corporis*.⁴ After World War II, *tinea pedis* increased together with global immigration.⁴ The epidemiology of these infections changed with the spread of *Microsporum canis*, *T. mentagrophytes* and *Trichophyton verrucosum* in Europe and *Trichophyton tonsurans* in North America as agents of *tinea capitis*. In the last decades, however, *Trichophyton rubrum* has become the predominant species in most of the studies performed, particularly in skin and nails.^{3,4}

In Spain, the lack of recent data on superficial mycoses makes it difficult to develop a reliable map of dermatophytoses. However, a change from zoophilic to anthropophilic species has been observed over the last decades.^{5,6} Species such as *T. tonsurans*, *Trichophyton violaceum* and *M. audouinii* are being reintroduced in this and other European countries, often related to people from endemic countries,^{5,7,8} and their capacity to produce outbreaks and to persist in healthy carriers represents a public health problem.^{9–11}

In this context, the aim of the present study was to evaluate the epidemiology of the dermatophytoses occurring from 2008 to 2017 in the health area of Barcelonès Nord, located in Northeastern Spain, and to compare it with the results of previous studies conducted in Spain and other European countries.

Material/Methods

From 2008 to 2017, 13,419 specimens obtained from patients with a superficial mycosis suspicion were submitted to the “Microbiology” Department, Laboratori Clínic Metropolitana Nord, Hospital Universitari Germans Trias i Pujol, “Department of Genetics and Microbiology”, Universitat Autònoma de Barcelona, Spain, the reference hospital of “Barcelonès Nord”, an area of 1,400,000 inhabitants. The samples came both from patients at the hospital and from primary health facilities. Only one sample per

patient and location was included to avoid bias in the epidemiological analysis, consequently, the positive samples included in the study belonged to different patients. Cultures with isolated non-dermatophyte fungi were excluded.

All specimens (hair, skin and nail) were examined by direct wet mount microscopy using 20% KOH. Samples were cultured on Sabouraud dextrose agar plates with chloramphenicol (BioMérieux, Marcy-L'Étoile, France), Sabouraud-dextrose agar plates with chloramphenicol and gentamicin (BioMérieux, Marcy-L'Étoile, France) and Sabouraud dextrose agar tubes with and without cycloheximide (BioMérieux, Marcy-L'Étoile, France). All cultures were incubated at 25 °C for at least 4 weeks and examined weekly. When no sporulation was observed, a subculture was incubated on Borelli's Lactrimel media (wheat flour 2% (w/v), cow's milk 20% (v/v), honey 1% (v/v), agar 2% (w/v), chloramphenicol 0,010% (w/v), water). The identification of dermatophytes was based on microscopic and macroscopic features of the fungi.

Patients were categorized depending on their age (until 15 and >15 years old) and their origin. Patients from South America and China were clustered with Spaniards under the heading of Caucasian. Most of these patients were long-term residents without documented travels and, accordingly, it was assumed that the dermatophytic infection took place in Spain. Patients from Africa, India and Pakistan were grouped together because, in many cases, the country of origin was difficult to determine. However, various sociodemographic data supported an African/Hindustani origin.

Clinical and sociodemographic data in positive cases, such as age, sex, origin, type of sample, species isolated and location of the *tinea* were recorded if possible.

Proportions were compared with Chi-square and Fisher's exact tests. The odds ratio (OR) was calculated when possible. A logistic regression model was used for multivariate analysis; the level of statistical significance was $p < 0.05$. Statistical data were generated using the software package Stata 14.


Results

Out of the 13,419 samples submitted, 607 (4.5%) were included in the study [313 from males (51.6%) and 294 from females (48.4%)]. Among these, 84 (13.8%) specimens were obtained from patients up to 15 years of age. Regarding the participants' origins, 458 were Spaniards, 10 were from South America, 10 from China, 17 from Morocco, 14 from Pakistan, 6 from India and 11 from sub-Saharan countries. The country of origin of the remaining 81 patients could not be determined, although various sociodemographic data supported an African/Hindustani origin.

The dermatophyte species isolated per year are shown in [Table 1](#). *T. rubrum* was the predominant fungus (76.1%) in all dermatophytoses unless *tinea barbae/tinea faciei*, showing a higher prevalence among the population older than 15 ([Fig. 1](#)). *T. mentagrophytes/Trichophyton interdigitale* was isolated in 11.9% of cases, mainly in *tinea corporis* and *tinea unguium pedum* ([Table 2](#)). The increase of non-endemic dermatophytes over time was also

Table 1
Number of dermatophytes isolated per year.

	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	Total
<i>Epidermophyton floccosum</i>	–	–	–	–	–	1	–	1	1	–	3
<i>Microsporium canis</i>	1	1	2	2	2	1	1	1	–	7	18
<i>Trichophyton mentagrophyte/interdigitale</i>	2	8	4	5	3	6	1	8	10	25	72
<i>Trichophyton rubrum</i>	4	14	23	20	14	18	8	27	126	208	462
<i>Microsporium gypseum</i>	–	–	1	1	2	1	–	1	1	3	10
<i>Trichophyton soudanense</i>	–	–	–	–	2	1	–	1	3	–	7
<i>Trichophyton tonsurans</i>	1	1	–	2	1	–	–	2	4	5	16
<i>Trichophyton schoenleinii</i>	–	–	–	–	–	–	1	–	–	–	1
<i>Microsporium audouinii</i>	–	–	–	–	–	1	–	–	4	3	8
<i>Trichophyton violaceum</i>	–	–	–	–	–	1	–	–	5	2	8
<i>Trichophyton terrestre</i>	–	–	–	1	–	–	–	–	–	–	1
<i>Trichophyton erinacei</i>	–	–	–	–	–	–	–	–	–	1	1

**Fig. 1.** Frequency of dermatophytes by location and age group. Statistical significance is indicated by * ($p < 0.01$); ** ($p < 0.001$).**Table 2**
Number of dermatophytes isolated by location.

	<i>Tinea corporis</i>	<i>Tinea cruris</i>	<i>Tinea capitis</i>	<i>Tinea pedis</i>	<i>Tinea barbae/tinea faciei</i>	<i>Tinea unguium pedum</i>	<i>Tinea unguium manuum</i>	Not located
<i>Epidermophyton floccosum</i>	1 (1) ^a	–	–	1	–	–	–	–
<i>Microsporium audouinii</i>	3 (3)	–	5 (5)	–	–	–	–	–
<i>Microsporium gypseum</i>	7 (1)	–	–	1	1	1	–	–
<i>Microsporium canis</i>	10 (4)	–	5 (2)	–	2 (2)	1	–	–
<i>Trichophyton mentagrophyte/interdigitale</i>	23 (2)	4	9 (5)	3	4	20	4	5 (1)
<i>Trichophyton rubrum</i>	122 (5)	7	23 (13)	46	–	196 (10)	65 (3)	3 (3)
<i>Trichophyton soudanense</i>	1 (1)	–	6 (6)	–	–	–	–	–
<i>Trichophyton tonsurans</i>	–	–	12 (8)	1 (1)	–	2	1	–
<i>Trichophyton schoenleinii</i>	–	–	1 (1)	–	–	–	–	–
<i>Trichophyton violaceum</i>	3 (1)	–	5 (5)	–	–	–	–	–
<i>Trichophyton terrestre</i>	–	–	–	–	–	1	–	–
<i>Trichophyton erinacei</i>	1 (1)	–	–	–	–	–	–	–
Total	171 (19)	11	66 (45)	52 (1)	7 (2)	221 (10)	71 (3)	8 (4)

^a Numbers in brackets correspond to dermatophytes isolated from patients up to 15 years of age.

noteworthy (11 *T. tonsurans*, 7 *M. audouinii*, 4 *Trichophyton soudanense* and 7 *T. violaceum* isolated in the last three years) (Table 1). The location of the tinea could not be determined in eight patients (five *T. rubrum* and three *T. mentagrophytes/T. interdigitale*).

Among the population older than 15, *tinea unguium pedum* (40.3%) was the main dermatophyte infection, followed by *tinea corporis* (29.1%) (Fig. 1). *Tinea capitis* was the most prevalent (53.6%) among patients until 15 years of age (Fig. 1).

Accordingly, *tinea pedis* ($p < 0.001$), *tinea unguium manuum* ($p = 0.010$; OR = 3.25) and *tinea unguium pedum* ($p < 0.001$; OR = 4.9)

were significantly linked to those over 15 (Fig. 1). *Tinea capitis* was statistically associated with patients up to 15 years of age ($p < 0.001$; OR = 20.74) and caused by a wide variety of fungi, highlighting anthropophilic dermatophytes such as *M. audouinii* ($n = 5$), *T. soudanense* ($n = 6$), *T. tonsurans* ($n = 12$), *T. schoenleinii* ($n = 1$) and *T. violaceum* ($n = 5$) (Table 2).

Epidemiological data were recorded from non-endemic dermatophytes. Among the eight *T. violaceum* specimens isolated, six belonged to patients until 15 (none of them exceeded 7 years) (Table 2). Five patients were of African origin, one was from




Fig. 2. Number of non-endemic dermatophytes by origin. Statistical significance is indicated by * ($p < 0.01$); ** ($p < 0.001$).

Pakistan and two were Spaniards (Fig. 2). Further epidemiological data could be obtained from five patients, of which four had moved to Spain a few months ago.

T. soudanense was the causal agent of seven dermatophytoses, of which five corresponded to African children (three from Gambia and one from Senegal) and two from Spanish children (Fig. 2). Among the latter, one shared school with another African child and the other had a mother of African origin. No data about travels or time of residence could be recorded from African patients.

T. schoenleinii was the dermatophyte agent of *tinea capitis favosa* ($n = 1$) (Table 2) in a Moroccan woman of 41 years, diagnosed in early childhood in his country of origin (Fig. 2).

M. audouinii was the causative agent of eight dermatophytoses (Table 2). Out of eight patients, seven were African children (two from Guinea and one from Mali) and one from Spain (Fig. 2). Further epidemiological information could only be recorded from four patients, of which two became infected in their country of origin.

T. tonsurans was the etiological agent of 16 dermatophyte infections (Table 2). Nine patients were Spaniards, four from South America and three from Africa (Fig. 2). Further epidemiological data could be recorded in eight cases. One local patient contracted the disease in South America, while another reported an outbreak within the family.

T. soudanense ($p < 0.0001$), *M. audouinii* ($p < 0.0001$), *T. tonsurans* ($p = 0.0002$; OR = 7.18) and *T. violaceum* ($p = 0.0002$; OR = 18.68) were significantly associated with patients up to 15 years of age.

Moreover, *T. soudanense* ($p = 0.0047$; OR = 10.20), *M. audouinii* ($p = 0.0004$; OR = 24.74) and *T. violaceum* ($p = 0.0002$; OR = 18.68) were associated with patients from Africa/Hindustan. An analysis of *T. schoenleinii* could not be performed due to the scarcity of isolates from this species.

An analysis of non-endemic dermatophytes grouping *T. soudanense*, *M. audouinii*, *T. violaceum* and *T. schoenleinii* as non-endemic dermatophytes was conducted (*T. tonsurans* was excluded from this analysis based on the univariate analysis). In the univariate analysis, non-endemic dermatophytes were associated with African/Hindustani origin ($p < 0.0001$; OR = 16.47) and patients until 15 years of age ($p < 0.0001$; OR = 50.20). When stratified by age, non-endemic dermatophytes were associated with African/Hindustani origin only in the population up to 15 years ($p < 0.0001$). In the multivariate analysis, non-endemic dermatophytes (all except *T. tonsurans* were considered non-endemic) were linked to ages until 15 years ($p < 0.001$), *tinea capitis* ($p = 0.0017$) and African/Hindustani origin ($p < 0.001$).

Discussion

Dermatophytoses are a major public health problem, affecting 20–25% of the global population.^{3,4} Their epidemiology has changed significantly throughout the 20th century,⁴ and their spread has been linked to factors such as economic status,^{3,8,12,13} population growth,⁴ immigration,^{5,7,8,11,14} living conditions⁴ or contact with animals.^{8,15}

In the present study *tinea unguium pedum* was the predominant dermatophytic infection (40.4%) among the population over 15 years of age, unlike previous studies performed in Spain,^{16,17} but consistent with a national study performed by Monzon et al.⁵ in 2001 and with recent surveys in Europe.^{13,18} The frequency of *tinea pedis* and *tinea unguium pedum* increased after World War II,⁴ and several sociocultural factors, such as occlusive footwear or the use of swimming pools, have been related to this increase.^{13,18} As expected, *tinea unguium manuum* ($p = 0.010$; OR = 3.25) and *tinea unguium pedum* ($p < 0.001$; OR = 4.9) were associated with patients over 15 years of age (Fig. 1). The increase of *tinea unguium* with age is well-known. Factors related to this dermatophytose in the elderly include poor peripheral circulation, repeated nail trauma and slower nail growth. By contrast, children and young people have a faster nail growth, a smaller nail surface and are more likely to start treatment at an earlier stage of the disease.¹³

In parallel, *T. rubrum* was the main dermatophyte in the present study. This anthropophilic species is distributed worldwide⁴ and its growth is favored in humid environments.⁶ It has become the most common etiological agent in the majority of the European surveys performed.^{5,6,8,19,20} The rise of this dermatophyte and its close relationship with *tinea unguium* has been widely discussed. Sociodemographic factors, such as the cohabitation of more than one generation, sharing towels and socks or the presence in family members have been related to the spread of the fungus.¹³

Tinea pedis was poorly represented in our study (9.8% among the population over 15 years and 1.2% up to 15 years), which is remarkable regarding other epidemiological surveys.^{5,6,16,18,21,22} The lack of clinical signs in a great percentage of cases could explain this finding. A study conducted by Perea et al.²³ in Madrid found that 55% of *tinea pedis* infections in healthy individuals were asymptomatic; the authors also observed a high prevalence of concomitant *tinea unguium* and *tinea pedis*, explaining the underrepresentation of *tinea pedis* in our work. Besides, in many cases, this *tinea* does not need for microbiological confirmation, and specimens are not usually sent to the laboratory. Similar to *tinea*

unguium, *tinea pedis* was associated with the population over 15 years ($p < 0.0001$).

Tinea capitis was the most prevalent (53.6%) dermatophytosis among patients up to 15 years, according to other Spanish and international surveys.^{5,7,12} *T. rubrum* was the most frequent dermatophyte in *tinea capitis*, which contrasts to previous Spanish studies, in which *M. canis* predominated.^{5,16,17,24} Generally, cats and dogs are the reservoir of *M. canis*,²⁴ and the low prevalence of this fungus in the present study could be explained by the absence of stray animals in this area. We observed other anthropophilic fungi as etiological agents of *tinea capitis*, such as *T. tonsurans*, *T. soudanense*, *M. audouinii* and *T. violaceum*. Kieliger and colleagues remarked the spread of these fungi in Zurich, mostly isolated in patients of African origin. They suggested contagion within the family unit as a necessary condition for their transmission.²⁵ In another French study, the authors also noted the high prevalence of such *Trichophyton* species as etiological agents of *tinea capitis*, mainly in patients from West Africa and coinciding with the return of children from their holidays in Africa.²⁶

T. rubrum was also the main dermatophyte in *tinea corporis*, which has been observed in other surveys from Spain, Sweden and Poland.^{5,10,27} However, in other studies from Italy²⁸ and Greece,²² *M. canis* was still the leading cause of *tinea corporis*.

T. mentagrophytes/T. interdigitale was the second most common dermatophyte and was mainly found in *tinea corporis* and *tinea unguium pedum*. Zoophilic strains of *T. mentagrophytes* are difficult to distinguish from anthropophilic *T. interdigitale* by optic microscopy, requiring molecular assays for this purpose,²⁹ which is a limitation of this study. In India, however, Chowdhary and colleagues have suggested that the epidemic observed in their country was more likely due to an anthropophilic species.²⁹ In the work carried out by Mazon et al. from 1989 to 1994, *T. mentagrophytes* was the main dermatophyte in *tinea corporis*,¹⁶ while in another survey performed by Monzon et al. in 2001, *T. mentagrophytes* was the second etiological agent in *tinea unguium* and *tinea pedis* and the most isolated fungus in *tinea barbae/tinea faciei* and *tinea manuum*.⁵ Boz and collaborators described *T. mentagrophytes* (var. *Mentagrophytes*) as the second most common agent of *tinea capitis* among the pediatric population in Southern Spain. The spread of zoophilic species such as *T. mentagrophytes* has been widely described in the past century in Spain, mainly associated with rural environments and close contact with animals.^{3,24}

The presence of anthropophilic species such as *T. soudanense*, *T. schoenleinii*, *M. audouinii*, *T. tonsurans* or *T. violaceum* as etiological agents of *tinea capitis* and *tinea corporis* (mostly in patients up to 15 years of age) is remarkable. Although extensive epidemiological data was difficult to obtain, most of patients were of African origin, except for *T. tonsurans* (the majority were Spanish). The association of *T. tonsurans* with Spanish patients is not surprising. In other European countries, such as England and France, *T. tonsurans* has become the most common cause of *tinea capitis*.^{3,26} However, further studies are needed to understand the prevalence and origin of this fungus among the study population.

Univariate and multivariate analyses revealed an association between these species with patients up to 15 years, *tinea capitis* and with patients from Africa/Hindustan (*T. tonsurans* was the only one not associated with African/Hindustani origin). Moreover, in some cases (four *T. violaceum*, one *T. schoenleinii* and two *M. audouinii*) there was evidence that these patients contracted the infection in their country of origin. These dermatophytes were highly prevalent in Spain and the rest of Europe at the beginning of the 20th century, mainly associated with *tinea capitis*.^{4,6} Nowadays, they are endemic throughout the African continent.¹² With advanced globalization, the reappearance of anthropophilic dermatophytes has been observed in Europe, provoking outbreaks in kindergartens and schools.^{11,14}

Previous surveys have already reported the emergence of these anthropophilic species across Europe.^{5,7–9,21,30} In Spain, a study performed by Monzón et al. documented the significant prevalence of *T. violaceum* and *T. tonsurans* among patients of African origin, mainly in the scalp.⁵ In another study, Teodolinda et al. observed a predominance of *T. violaceum* as etiological agent of *tinea capitis* among African patients, although they seemingly contracted the infection in Italy.⁷

The presence of asymptomatic carriers in some studies is also noteworthy.^{9,11,14} In a prospective study conducted by Cuetara et al. from 1994 to 1996 with 10,000 healthy children, the authors detected 19 asymptomatic carriers (13 were *T. tonsurans*).⁹ In another study in Paris, Gits-Muselli reported a high number of asymptomatic carriers of *M. audouinii*.²⁶

This study has some limitations. Firstly, it only reflects positive cultures from specimens delivered to the laboratory. Therefore, it may not be fully representative of all clinically diagnosed dermatophytoses. Secondly, the use of partial epidemiological data hampers to determine where the disease was contracted, which would allow us to a better understanding of the epidemics dynamic of the above mentioned non-endemic dermatophytes. Moreover, the African/Hindustani group reflects a heterogeneous population with very diverse origins. This is mainly due to the lack of sociodemographic information for many patients, of whom it was difficult to determine the country of origin. However, the objective of the present study was not to describe the epidemiology of superficial mycoses in these countries, but to study the evolution and possible causes of dermatophytoses in our environment.

In conclusion, we confirmed the spread of *T. rubrum* as the predominant dermatophyte in our area, replacing zoophilic species such as *M. canis* or *T. mentagrophytes*. Moreover, we reported an increase in non-endemic anthropophilic dermatophytes in the last few years, especially in the African and Hindustani population up to 15 years of age. These agents are highly contagious and require continuous monitoring. However, prospective population-based studies are needed to understand the determinants of dermatophytosis in the study area.

Ethical approval

The project related to this manuscript was approved by the Ethical Committee of the Hospital Germans Trias i Pujol.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of interest

None declared.

References

- de Diego A. Clinical, diagnostic and therapeutic aspects of dermatophytosis. *Enferm Infecc Microbiol Clin.* 2011;29 Suppl. 3:33–9.
- Crespo Erhiga V, Gómez Moyano E, Crespo Palomo M. *Dermatomycosis. The Laboratory diagnosis within the reach of the dermatologists.* 1st ed. Madrid: Creación, Ergón; 2015.
- Havlickova B, Czaika VA, Friedrich M. Epidemiological trends in skin mycoses worldwide. *Mycoses.* 2008;51 Suppl. 4:2–15.
- Zhan P, Liu W. The changing face of dermatophytic infections worldwide. *Mycopathologia.* 2017;182:77–86.
- de la Torre A, Cuenca-Estrella M, Rodríguez-Tudela JL. Epidemiological survey of dermatophytosis in Spain (April–June 2001). *Enferm Infecc Microbiol Clin.* 2003;21:477–83.
- García-Martos P, García-Agudo L, Agudo-Pérez E, de Sola F, Linares M. Dermatomycoses due to anthropophilic fungi in Cadiz, Spain, between 1997 and 2008. *Actas Dermosifiliogr.* 2010;101:242–7.

7. Mapelli ET, Cerri A, Bombonato C, Menni S. *Tinea capitis* in the paediatric population in Milan, Italy: the emergence of *Trichophyton violaceum*. *Mycopathologia*. 2013;176:243–6.
8. Farina C, Fazio P, Imberti G, Lombardi G, Passera M, Andreoni S, et al. *Trichophyton violaceum* and *T. soudanense*: re-emerging pathogens in Italy, 2005–2013. *New Microbiol.* 2015;38:409–15.
9. Cuetara MS, del Palacio A, Pereiro M, Amor E, Alvarez C, Noriega AR. Prevalence of undetected *tinea capitis* in a school survey in Spain. *Mycoses*. 1997;40:131–7.
10. Budak A, Bogusz B, Tokarczyk M, Trojanowska D. Dermatophytes isolated from superficial fungal infections in Krakow, Poland, between 1995 and 2010. *Mycoses*. 2013;56:422–8.
11. Durán-Valle MT, Regodón-Domínguez M, Velasco-Rodríguez MJ, Aragón A, Gómez-Garcés JL. Outbreak of *Trichophyton tonsurans* ringworm in a health area of the Community of Madrid (Spain). *Rev Iberoam Micol.* 2016;33:126–8.
12. Coulibaly O, L'Ollivier C, Piarroux R, Ranque S. Epidemiology of human dermatophytes in Africa. *Med Mycol.* 2018;56:145–61.
13. Dubljanin E, Džamić A, Vujčić I, Grujičić S, Arsenijević VA, Mitrović S, et al. Epidemiology of onychomycosis in Serbia: a laboratory-based survey and risk factor identification. *Mycoses*. 2017;60:25–32.
14. Zink A, Papanagioutou V, Todorova A, Seidl HP, Niedermeier A, Ring J, et al. Outbreak of *Microsporium audouinii* in Munich – the return of infectious fungi in Germany. *Mycoses*. 2014;57:765–70.
15. Farag AGA, Hammam MA, Ibrahim RA, Mahfouz RZ, Elnaidany NF, Qutubuddin M, et al. Epidemiology of dermatophyte infections among school children in Menoufia Governorate, Egypt. *Mycoses*. 2018;61:321–5.
16. Mazón A, Salvo S, Vives R, Valcayo A, Sabalza MA. Etiologic and epidemiologic study of dermatomycoses in Navarra (Spain). *Rev Iberoam Micol.* 1997;14:65–8.
17. Padilla A, Sampedro A, Sampedro P, Delgado V. Clinical and epidemiological survey of dermatophytoses in Jaen (Spain). *Rev Iberoam Micol.* 2002;19:36–9.
18. Vena GA, Chieco P, Posa F, Garofalo A, Bosco A, Cassano N. Epidemiology of dermatophytoses: retrospective analysis from 2005 to 2010 and comparison with previous data from 1975. *New Microbiol.* 2012;35:207–13.
19. Rodríguez-Pazos L, Pereiro-Ferreirós MM, Pereiro M, Toribio J. Onychomycosis observed in children over a 20-year period. *Mycoses*. 2011;54:450–3.
20. Nasr A, Vyzantiadis TA, Patsatsi A, Louka A, Ioakimidou A, Zachrou E, et al. Epidemiology of superficial mycoses in Northern Greece: a 4-year study. *J Eur Acad Dermatol Venereol.* 2016;30:837–9.
21. Faure-Cognet O, Fricker-Hidalgo H, Pelloux H, Leccia MT. Superficial fungal infections in a French teaching hospital in Grenoble area: retrospective study on 5470 samples from 2001 to 2011. *Mycopathologia*. 2016;181:59–66.
22. Maraki S, Tselentis Y. Survey on the epidemiology of *Microsporium canis* infections in Crete, Greece over a 5-year period. *Int J Dermatol.* 2000;39:21–4.
23. Perea S, Ramos MJ, Garau M, Gonzalez A, Noriega AR, del Palacio A. Prevalence and risk factors of *tinea unguium* and *tinea pedis* in the general population in Spain. *J Clin Microbiol.* 2000;38:3226–30.
24. Del Boz J, Crespo V, Rivas-Ruiz F, De Troya M. A 30-year survey of paediatric *tinea capitis* in southern Spain. *J Eur Acad Dermatol Venereol.* 2011;25:170–4.
25. Kieliger S, Glatz M, Cozzio A, Bosshard PP. *Tinea capitis* and *tinea faciei* in the Zurich area – an 8-year survey of trends in the epidemiology and treatment patterns. *J Eur Acad Dermatol Venereol.* 2015;29:1524–9.
26. Gits-Muselli M, Benderdouche M, Hamane S, Mingui A, De Chauvin MF, Guigee N, et al. Continuous increase of *Trichophyton tonsurans* as a cause of *tinea capitis* in the urban area of Paris, France: a 5-year-long study. *Med Mycol.* 2017;55:476–84.
27. Drakensjö IT, Chryssanthou E. Epidemiology of dermatophyte infections in Stockholm, Sweden: a retrospective study from 2005–2009. *Med Mycol.* 2011;49:484–8.
28. Panasiti V, Devirgiliis V, Borroni RG, Mancini M, Curzio M, Rossi M, et al. Epidemiology of dermatophytic infections in Rome, Italy: a retrospective study from 2002 to 2004. *Med Mycol Mycol.* 2007;45:57–60.
29. Chowdhary A, Singh A, Singh PK, Khurana A, Meis JF. Perspectives on misidentification of *Trichophyton interdigitale/Trichophyton mentagrophytes* using internal transcribed spacer region sequencing: urgent need to update the sequence database. *Mycoses*. 2019;62:11–5.
30. Donghi D, Hauser V, Bosshard PP. *Microsporium audouinii tinea capitis* in a Swiss school: assessment and management of patients and asymptomatic carriers. *Med Mycol.* 2011;49:324–8.