



# Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Original article

## Methicillin-resistant *Staphylococcus aureus* in swine housed indoors in Galicia, Spain



Antonio Moreno-Flores<sup>a,b</sup>, Carmen Potel-Alvarellos<sup>a,b</sup>, Mónica Francisco-Tomé<sup>b</sup>, Lucia Constenla-Caramés<sup>b</sup>, Eduardo Pérez-Roth<sup>c</sup>, Carolina López-Cotón<sup>d</sup>, Estefanía Comesaña-Da Vila<sup>d</sup>, Lourdes Eiroa-de la Puente<sup>d</sup>, Maximiliano Álvarez-Fernández<sup>a,b,\*</sup>

<sup>a</sup> Servicio de Microbiología, Hospital Álvaro Cunqueiro (CHUVI), Vigo, Pontevedra, Spain

<sup>b</sup> Instituto de Investigación Sanitaria Galicia Sur, Grupo Microbiología e Infectología, Spain

<sup>c</sup> Departamento de Bioquímica, Microbiología, Biología Celular y Genética, Facultad de Ciencias, Universidad de La Laguna, La Laguna, Spain

<sup>d</sup> Ciclo Superior de Laboratorio Clínico, I.E.S. Manuel Antonio Vigo, Vigo, Spain

### ARTICLE INFO

#### Article history:

Received 11 January 2019

Accepted 8 March 2019

Available online 10 May 2019

#### Keywords:

Methicillin-resistant *Staphylococcus aureus*

(MRSA)

Swine

Antimicrobial resistance

Farming cleaning procedures

### ABSTRACT

**Introduction:** Livestock are known reservoirs of methicillin-resistant *Staphylococcus aureus* (MRSA) and this constitutes an important public health issue. The prevalence of nasal MRSA carriers in swine housed indoors in Galicia, Spain, was studied.

**Methods:** 197 samples from swine aged three, eight, 12, 16 and 24 weeks, and from adult pigs, were obtained from four farms. The cleaning procedures implemented to clean the barns and antimicrobial consumption were analyzed. Antimicrobial susceptibility and antimicrobial resistance genes were studied. PFGE, *spa* typing and MLST were used to classify the isolates. *SCCmec*, *agr* and *pvl* were analyzed.

**Results:** MRSA prevalence was 12.7%. Swine younger than 16 weeks had a higher colonization rate; 22.9% vs 3.5% (OR, 8.16; 95% CI, 2.47–29.79;  $p < 0.01$ ). The only farm found to be MRSA-free used disinfectants as part of its cleaning procedure. All MRSA were tetracycline-resistant (identifying the *tetK* and *tetM* genes), 80% were resistant to erythromycin and clindamycin and 16% were only clindamycin-resistant. The *ermC* and *vgaA* genes were identified in these two phenotypes. A single genotype (PFGE type A) and ST398 – *spa* t011 (84%) and t1451 (16%) were identified. *SCCmec* type V and *agrI* were identified in all isolates, and all were *pvl*-negative.

**Conclusion:** A correlation between swine age and MRSA colonization was observed. Appropriate cleaning procedures could have an impact on MRSA colonization in farming. Resistance to antibiotics used in human health was identified. Clinicians should be aware if their patients have come into contact with farm animals.

© 2019 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.

## **Staphylococcus aureus resistente a la meticilina en cerdos estabulados en Galicia, España**

### R E S U M E N

**Introducción:** Los animales de granja son reservorios de *Staphylococcus aureus* resistente a la meticilina (SARM), y constituyen un problema de salud pública. Se estudia la prevalencia de portadores nasales de SARM en cerdos estabulados en Galicia, España.

**Métodos:** En 4 explotaciones se obtuvieron 197 muestras de cerdos con edades en semanas de 3, 8, 12, 16, 24 y adultos. Se analizaron los métodos empleados para limpiar los establos y el consumo de antimicrobianos. Se estudió la resistencia a antimicrobianos, y los genes involucrados en esta. Los aislamientos fueron clasificados mediante PFGE, *spa* y MLST. Se analizaron *SCCmec*, *agr* y *pvl*.

**Resultados:** La prevalencia de SARM fue del 12,7%. Los cerdos de < 16 semanas presentaron las frecuencias de colonización más elevadas 22,9 vs. 3,5% (OR: 8,16; IC 95%: 2,47-29,79;  $p < 0,01$ ). En la única explotación libre de SARM se empleaban desinfectantes en la limpieza. Todos los SARM fueron resistentes a

#### Palabras clave:

*Staphylococcus aureus* resistente a la

meticilina

Cerdos

Resistencia antimicrobianos

Procedimientos de limpieza en

explotaciones ganaderas

\* Corresponding author.

E-mail addresses: Maximiliano.alvarez.fernandez@sergas.es, mxalvfer@gmail.com (M. Álvarez-Fernández).

tetraciclina identificándose los genes *tetK* y *tetM*, el 80% fueron resistentes a eritromicina y clindamicina, y el 16% fueron únicamente resistentes a clindamicina. Se identificaron los genes *ermC* y *vgaA* en estos 2 fenotipos. Se identificó un único genotipo (PFGE-A) y ST398, siendo *spa* t011 (84%) y t1451 (16%). En todos los aislamientos se identificó *SCCmec V* y *agrI*, siendo estos *pvl* negativos.

**Conclusiones:** Se observó la asociación entre edad y colonización SARM. La limpieza adecuada podría modificar la colonización por SARM. Se detectaron resistencias a antibióticos empleados en humanos. Los médicos deberían conocer si los pacientes tienen contacto con animales de granja.

© 2019 Elsevier España, S.L.U. y Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Todos los derechos reservados.

## Introduction

In 2005 a methicillin-resistant *Staphylococcus aureus* (MRSA) strain associated with pig farming was identified in France and The Netherlands.<sup>1,2</sup> The MRSA strain belonged to sequence type (ST) 398 clone which has also been further identified in animals and humans worldwide.<sup>3–6</sup> Indeed, 20% of all MRSA identified in humans in The Netherlands belonged to the ST398, documenting the importance of considering livestock and other animals when studying the epidemiology of MRSA.<sup>7</sup>

Spain is the fourth porcine producer in the world and the second in Europe behind Germany.<sup>8</sup> The current information about MRSA colonization in live swine in Spain is scarce. Porrero et al.<sup>9</sup> studied the MRSA epidemiology in free living Iberian pigs and Reynaga et al.<sup>3</sup> studied MRSA colonization in farm pigs. Other Spanish studies were done in slaughter pigs<sup>10</sup> and food samples.<sup>11</sup> Nevertheless, pigs could have become MRSA colonized during transportation from farm to slaughterhouse<sup>12</sup> not representing the real farm prevalence.

In 2006, ST398 MRSA was first isolated from patients with different infections in our hospital.<sup>13</sup> We concluded that the ST398 clone was part of the MRSA population in Galician region (Spain), hypothesizing that pigs could be a potential reservoir of MRSA from a public health perspective.

The aim of this study was to determine the prevalence of nasal MRSA carriage in live swine hosted in indoor production systems in Galician region, Spain. The isolates were genetically classified and the antimicrobial resistance was studied. The cleaning procedures used in each of the studied farms were questioned.

## Materials and methods

### Farms description

The four sampled farms were located in an area of 80 km<sup>2</sup> in Galicia (Spain). From March to April 2010 we collected a total of 197 nasal swabs from two farrow-to-rearing farms (farms A and B, pigs were up to 8 weeks), a rearing farm (farm C; 8–24 weeks, at the moment of the study the pigs were 16 weeks), and a farrow-to-finish farm (farm D; 90 sows, 140 piglets 3 weeks old, 170 grower pigs 12–24 weeks old). The sample comprised 28 piglets 3 weeks old, 45 weaned piglets 8 weeks old, 10 grower pigs 12 weeks old, 75 grower pigs 16 weeks old, 25 finisher pigs 24 weeks old, and 14 adult breeding sows (Table 1). The farms were questioned about the cleaning procedures and the antimicrobial consumption. The origin of the swine was also inquired. The permissions for this study were obtained from the farm owners allowing us to sample a previously defined number of animals. For the development of this study the European Directive 2010/63/EU was fulfilled, which is transposed to Spanish national legislation by means of Real Decreto (RD) 53/2013 for the protection of animals for experimentation and other scientific purposes.

**Table 1**

Swine age and methicillin-resistant *Staphylococcus aureus* (MRSA) isolated by farm. Farm denomination and MRSA/total *S. aureus* (%).

Age (weeks)	A <sup>a</sup>	B <sup>a</sup>	C <sup>a</sup>	D <sup>a</sup>	Total
3	2/4 (50)	1/4(25)		0/20	3/28 (10.7)
8	9/21(43)	6/24(25)			15/45 (33.3)
12				3/10(30)	3/10 (30)
16			0/75		0/75
24				2/25(8)	2/25 (8)
Adult	1/2 (50)	1/2(50)		0/10	2/14 (14.3)
Total	12/27(44)	8/30(26.6)	0/75	5/65(7.7)	25/197 (12.7)

<sup>a</sup> Swine present in each farm: A, 750; B, 1000; C, 750; D, 400.

### Sample collection, bacterial isolation, and identification


A total of 197 nasal swine samples were collected using sterile swabs and inserted in Stuart's medium at 4 °C for transportation. The two nares were sampled with the same swab. The number of swine sampled represented 6.8% of the total pigs hosted in the farms at the moment of the study. Samples were not pooled, so they were inoculated one by one into 2 mL enrichment broth containing 10 g tryptone/L, 75 g mannitol/L and 2.5 g yeast extract/L.<sup>4</sup> After 24 h of incubation at 35 °C, a loopful of broth was inoculated onto selective MRSA agar plate (ChromID MRSA, bioMérieux) and colistin-nalidixic acid agar with 5% lamb blood (CNA) (bioMérieux). The plates were incubated for 24–48 h at 35 °C and examined for MRSA. Only one MRSA isolate was selected for further analysis. Isolates were confirmed to be *S. aureus* by the catalase test, the tube coagulase test and biochemically identified using the VITEK 2 GP card (bioMérieux). Methicillin resistance was confirmed by testing for the presence of penicillin binding protein 2 (PBP2') (MRSA latex agglutination test, Oxoid).

### Pulsed field gel electrophoresis (PFGE) analysis

Whole DNA from each MRSA isolate was analyzed by *SmaI*, *Cfr9I* and *EagI*, as previously described using the CHEF DR-III System (Bio-Rad).<sup>4,14,15</sup> *S. aureus* NCTC 8325 was included as reference strain. Genetic similarity between profiles was determined by computer analysis of the banding patterns using the InfoQuest FP software (Bio-Rad). Percent similarities were identified on a dendrogram derived from the unweighted pair group method using arithmetic averages and based on Dice coefficients. Band position tolerance and optimization were set at 1.0 and 0.5%, respectively. A similarity coefficient of 80% was selected to define the pulsed-field type clusters. Every band difference within a PFGE type resulted in adding a numerical order to the pulsed field cluster.<sup>16</sup>

### *Spa* typing, multilocus sequence typing (MLST), staphylococcal chromosome cassette (SCCmec) typing, and *agr* typing

DNA was extracted using the RTP-bacteria minikit (STRATEK Molecular). All isolates were analyzed by *spa* typing and MLST.<sup>17,18</sup>



**Fig. 1.** Dendrogram showing the PFGE relatedness using *EagI* and *Cfr9I*. The Dice coefficient of similarity (D) was 0.84. One cluster, A, was detected and four PFGE subtypes were identified (A1, A2, A3, and A4). The *spa* types, susceptibility to erythromycin (EM) and clindamycin (CM), number of isolates per each PFGE and *spa* types, and the farms where each PFGE and *spa* were isolated are shown. \* Farm A, 12 strains; farm B, 8 strains; and farm D, 5 strains. R and S mean resistant and susceptible respectively. -, no susceptibility studied, \*\*NCTC 8325, its chromosomal DNA was digested with *SmaI*.

MRSA isolates were *SCCmec* typed.<sup>19</sup> The *agr* type was also identified.<sup>20</sup> The presence of the virulence *pvl* gene was determined by PCR.<sup>21</sup>

#### Antimicrobial susceptibility testing and detection of antibiotic resistance genes

All MRSA were tested for antimicrobial susceptibility by the VITEK 2 AST-P588 card (bioMérieux, France). Susceptibility to tetracycline and daptomycin was studied using Etest on Mueller-Hinton agar (bioMérieux, France). The results were interpreted following the EUCAST recommendations.<sup>22</sup> *S. aureus* ATCC 29213 was used for quality assurance. Antimicrobial resistance genes were detected by PCR amplification as previously described.<sup>23–25</sup>

#### Data analysis

Chi-square test was used to analyze the statistical significance. A significance level of 0.05 was used in the analysis.

## Results

#### Detection of MRSA in the studied samples, cleaning procedures used, pig's origin, and antimicrobial consumption

Nasal swabs were taken from 197 swine hosted in four farms representing 6 different age groups (Table 1). The overall prevalence was 12.7% ( $n=25$ ). Swine younger than 16 weeks old had higher odds of MRSA colonization 22.9 vs 3.5% (OR, 8.16; 95% CI, 2.47–29.79;  $p < 0.01$ , Chi square test) when compared to swine older than 16 weeks. No MRSA was isolated in the farm C where all the pigs were 16 weeks old.

The cleaning procedure used in farm C included farm emptying, wash using high-pressure water, disinfection using 250-PPM bleach solution pulverization, washed again with high-pressure

water, and finally painted with quicklime. This procedure was repeated three times per year. The cleaning procedures in the other three farms comprised just cleaning with high-pressure water.

The farms informed that swine came solely from other local farms in Galicia. As well, they denied any antimicrobial use or consumption in the farms.

#### PFGE, *spa*, MLST, and *SCCmec* typing of MRSA isolates

Twenty-five MRSA isolates were subjected to PFGE. The isolates were non typeable when the DNA was digested with *SmaI* but they were typeable with *EagI* and *Cfr9I*. All isolates were closely related, Dice coefficient of similarity  $\geq 0.84$ , being identified one PFGE cluster and four PFGE subtypes (Fig. 1).

The *spa*-type t011 was identified in 21 isolates (84%). The *spa*-type t1451 was identified in 4 isolates (16%) (Fig. 1). All isolates belonged to the ST398 clone.

All isolates were *SCCmec* type V, *agrI* type and *pvl*-negative.

#### Antimicrobial susceptibility and resistance genes

All isolates were methicillin and tetracycline resistant. They were susceptible to trimethoprim-sulfamethoxazole, gentamicin, tobramycin, levofloxacin, linezolid, daptomycin, vancomycin, teicoplanin, rifampicin, mupirocin, fusidic acid, fosfomicin, nitrofurantoin, and tigecycline. Twenty strains (80%) were resistant to erythromycin (EM) and clindamycin (CM) and this resistance was constitutive. Four strains (16%) were resistant to CM but susceptible to EM, and one strain was EM and CM susceptible (4%) (Fig. 1).

The *tetK* and *tetM* genes were detected in all isolates, the *tetL* gene was not identified. The *ermC* gene was detected in all EM<sup>R</sup>-CM<sup>R</sup> isolates. The *ermA* and *B* genes were not detected in any isolate. The *vgaA* gene was identified in four EM<sup>S</sup>-CM<sup>R</sup> isolates, all belonging to *spa*-type t1451.

## Discussion

Galicia is one of the western regions in the European Union. The overall prevalence of MRSA carriage in pigs (12.7%) was lower to the one reported by other series.<sup>3–5,26</sup> Remarkably, no MRSA was identified in farm C where all pigs were 16 weeks old. In this production, pigs were hosted for four months before being delivered for consumption. Once the facility was totally emptied, an intensive cleaning procedure was always performed with cheap products like water, bleach and quicklime painting. As has been suggested by others, the presence of MRSA among pigs is related to the introduction of MRSA positive animals in a MRSA negative scenario, also the herd size could influence in the MRSA prevalence.<sup>5,27</sup> The origin of the pigs was questioned, and it was verified that local farms were the only pig suppliers. That makes us to think that MRSA is a usual colonizer among swine in Galicia. The different cleaning procedures and the pig's age could be an important factor in the absence of swine's MRSA colonization in farm C. Moreover, MRSA has been identified in dust samples obtained from farms and from the trucks used for pig transportation. Therefore, these studies underline the role of the environment as a source of MRSA in the pig production chain.<sup>6,12,27</sup>

In this work the rate of colonization by SARM was higher in younger pigs, as previously showed by Smith TC et al.<sup>4</sup>

According to other national studies,<sup>3,9</sup> the *spa*-type t011 was the most prevalent (84%). In this study, t1451 was isolated from 4 pigs only in farm D and four PFGE subtypes in the t011 *spa*-type were identified (Fig. 1). Therefore, we agree with other studies proposing that the transmission and outbreaks of MRSA ST398 could be better studied using PFGE which showed a higher discriminatory power.<sup>15,16</sup> All isolates contained the SCCmec type V, which is the most common SCCmec type identified in MRSA ST398.<sup>7</sup>

All isolates were resistant to tetracycline and carried the *tetM* and *tetK* genes. These genes, alone or in combination, are the most frequently identified in ST398 isolates.<sup>28</sup> The resistance to tetracycline was highly expected and this resistance has been proposed as a marker to suspect the presence of MRSA ST398 in hospitals.

Regarding the macrolide-lincosamides resistance genes, *ermC* was the only *erm* gene detected. The unusual macrolide-lincosamide resistance phenotype EM<sup>S</sup>-CM<sup>R</sup> was identified in four MRSA in farm D being all classified as *spa*-type 1451, reflecting a local endemism of this resistance phenotype. The *vga(A)* gene was identified as the responsible for this resistance.<sup>29</sup>

Our work has several limitations. Neither swine were sampled when arriving at the four farms nor environmental samples were taken along the time when pigs were hosted in the farms. Therefore, we do not know the real influence of the cleaning procedures on the MRSA colonization of the swine. Besides, the resistance genes studied were limited we cannot exclude the presence of others. Moreover, only four farms were studied, our results may not be generalized to other swine farms.

Moreover, the adaptation of ST398 to the human being makes MRSA ST398 a matter of concern among humans and the public health representing a risk to themselves and the community.<sup>30</sup> So, given the prevalence of swine MRSA colonization clinicians should be aware of the risk of MRSA colonization in people related with animal farming.

In summary, this study shows the MRSA colonization of pigs in indoor swine farms, being the MRSA prevalence lower in older swine. All MRSA were resistant to antimicrobials currently used in the clinical setting. The identification of a MRSA negative farm that used disinfecting and cleaning procedures suggests that appropriate hygienic practices could also be important in farming, as has been shown in the clinical setting.

## Sources of financing

This work was financed by grants from Fondo de Investigaciones Sanitarias and Fondo Europeo para el Desarrollo Regional (FEDER) (07-0812) and Consejería de Sanidad de La Junta de Galicia (PS08/34).

## Conflict of interest

The authors declare that there are not any conflicts of interest of any nature.

This work was partially presented at the 21st European Congress of Clinical Microbiology and Infectious Diseases (ECCMID) Milan, Italy.

## Acknowledgments

We are grateful to farm workers and veterinarians for their help.

## References

- Armand-Lefevre L, Ruimy R, Andrement A. Clonal comparison of *Staphylococcus aureus* isolates from healthy pig farmers, human controls, and pigs. *Emerg Infect Dis.* 2005;11:711–4.
- Voss A, Loeffen F, Bakker J, Klaassen C, Wulf M. Methicillin-resistant *Staphylococcus aureus* in pig farming. *Emerg Infect Dis.* 2005;11:1965–6.
- Reynaga E, Navarro M, Vilamala A, Roure P, Quintana M, García-Núñez M, et al. Prevalence of colonization by methicillin-resistant *Staphylococcus aureus* ST398 in pigs and pig farm workers in an area of Catalonia, Spain. *BMC Infect Dis.* 2016;16:716.
- Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers. *PLoS ONE.* 2009;4:e4258.
- Broens EM, Graat EAM, Van der Wolf PJ, Van de Giessen AW, De Jong MCM. Prevalence and risk factor analysis of livestock associated MRSA-positive pig herds in The Netherlands. *Prev Vet Med.* 2011;102:41–9.
- European Food Safety Authority. Analysis of the baseline survey on the prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in holdings with breeding pigs, in the EU, 2008 [1]-Part A: MRSA prevalence estimates. *EFSA J.* 2009;7:1376–458.
- van Loo I, Huijsdens X, Tiemersma E, de Neeling A, van de Sande-Bruinsma N, Beaujean D, et al. Emergence of methicillin-resistant *Staphylococcus aureus* of animal origin in humans. *Emerg Infect Dis.* 2007;13:1834–9.
- Gerosa S. The state of food and agriculture. Livestock in the balance. Part III, Statistical Annexes. Food and Agriculture Organization of The United Nations. Rome: Electronic Publishing Policy and Support Branch Communication Division FAO; 2009.
- Porrero MC, Wassenaar TM, Gómez-Barrero S, García M, Bárcena C, Alvarez J, et al. Detection of methicillin-resistant *Staphylococcus aureus* in Iberian pigs. *Lett Appl Microbiol.* 2012;54:280–5.
- Gómez-Sanz E, Torres C, Lozano C, Fernández-Pérez R, Aspiroz C, Ruiz-Larrea F, et al. Detection, molecular characterization, and clonal diversity of methicillin-resistant *Staphylococcus aureus* CC398 and CC97 in Spanish slaughter pigs of different age groups. *Foodborne Pathog Dis.* 2010;7:1269–77.
- Lozano C, López M, Gómez-Sanz E, Ruiz-Larrea F, Torres C, Zarazaga M. Detection of methicillin-resistant *Staphylococcus aureus* ST398 in food samples of animal origin in Spain. *J Antimicrob Chemother.* 2009;64:1325–6.
- Broens EM, Graat EAM, Van der Wolf PJ, Van de Giessen AW, De Jong MCM. Transmission of methicillin resistant *Staphylococcus aureus* among pigs during transportation from farm to abattoir. *Vet J.* 2011;189:302–5.
- Potel C, Alvarez-Fernández M, Constenla L, Alvarez P, Perez S. First human isolates of methicillin-resistant *Staphylococcus aureus* sequence type 398 in Spain. *Eur J Clin Microbiol Infect Dis.* 2010;29:351–2.
- Huijsdens XW, Bosch T, van Santen-Verheul MG, Spalburg E, Pluister GN, van Luit M, et al. Molecular characterisation of PFGE non-typable methicillin-resistant *Staphylococcus aureus* in The Netherlands, 2007. *Euro Surveill.* 2009;14, pii=19335.
- Argudín MA, Fetsch A, Tenhagen B-A, Hammerl JA, Hertwig S, Kowall J, et al. High heterogeneity within methicillin-resistant *Staphylococcus aureus* ST398 isolates, defined by Cfr91 macrorestriction-pulsed-field gel electrophoresis profiles and *spa* and SCCmec types. *Appl Environ Microbiol.* 2010;76:652–8.
- Bosch T, de Neeling AJ, Schouls LM, van der Zwaluw KW, Kluytmans JAJW, Grundmann H, et al. PFGE diversity within the methicillin-resistant *Staphylococcus aureus* clonal lineage ST398. *BMC Microbiol.* 2010;10:40.
- Shopsin B, Gomez M, Montgomery SO, Smith DH, Waddington M, Dodge DE, et al. Evaluation of protein A gene polymorphic region DNA sequencing for typing of *Staphylococcus aureus* strains. *J Clin Microbiol.* 1999;37:3556–63.

18. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol.* 2000;**38**:1008–15.
19. Kondo Y, Ito T, Ma XX, Watanabe S, Kreiswirth BN, Etienne J, et al. Combination of multiplex PCRs for staphylococcal cassette chromosome *mec* type assignment: rapid identification system for *mec*, *ccr*, and major differences in junkyard regions. *Antimicrob Agents Chemother.* 2007;**51**:264–74.
20. Tenover FC, McDougal LK, Goering RV, Killgore G, Projan SJ, Patel JB, et al. Characterization of a strain of community-associated methicillin-resistant *Staphylococcus aureus* widely disseminated in the United States. *J Clin Microbiol.* 2006;**44**:108–18.
21. Lina G, Piémont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, et al. Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis.* 1999;**29**:1128–32.
22. EUCAST. Breakpoint tables for interpretation of MICs and zone diameters. European Committee on Antimicrobial Susceptibility Testing (EUCAST); 2011. p. Version 1.3 January 5. Available from: <http://www.eucast.org/Version1.3>
23. Lozano C, Aspiroz C, Ezpeleta AI, Gómez-Sanz E, Zarazaga M, Torres C. Empyema caused by MRSA ST398 with atypical resistance profile, Spain. *Emerg Infect Dis.* 2011;**17**:138–40.
24. Ng LK, Martin I, Alfa M, Mulvey M. Multiplex PCR for the detection of tetracycline resistant genes. *Mol Cell Probes.* 2001;**15**:209–15.
25. Martineau F, Picard FJ, Lansac N, Ménard C, Roy PH, Ouellette M, et al. Correlation between the resistance genotype determined by multiplex PCR assays and the antibiotic susceptibility patterns of *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Antimicrob Agents Chemother.* 2000;**44**:231–8.
26. Morcillo A, Castro B, Rodríguez-Álvarez C, González JC, Sierra A, Montesinos MI, et al. Prevalence and characteristics of methicillin-resistant *Staphylococcus aureus* in pigs and pig workers in Tenerife, Spain. *Foodborne Pathog Dis.* 2012;**9**:207–10.
27. Fromm S, Beißwanger E, Käsbohrer A, Tenhagen B-A. Risk factors for MRSA in fattening pig herds – a meta-analysis using pooled data. *Prev Vet Med.* 2014;**117**:180–8.
28. Argudín MA, Tenhagen B-A, Fetsch A, Sachsenröder J, Käsbohrer A, Schroeter A, et al. Virulence and resistance determinants of German *Staphylococcus aureus* ST398 isolates from nonhuman sources. *Appl Environ Microbiol.* 2011;**77**:3052–60.
29. Kadlec K, Pomba CF, Couto N, Schwarz S. Small plasmids carrying *vga(A)* or *vga(C)* genes mediate resistance to lincosamides, pleuromutilins and streptogramin A antibiotics in methicillin-resistant *Staphylococcus aureus* ST398 from swine. *J Antimicrob Chemother.* 2010;**65**:2692–3.
30. Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. *Staphylococcus aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. *MBio.* 2013;**4**:e00520–612.