



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Original article

Serologic study of *Bartonella* sp. infection among human population of Southern Spain



Sonia Santibáñez^b, Antonio Caruz^c, Julia Márquez-Constán^d, Aránzazu Portillo^b, Jose Antonio Oteo^b, Francisco J. Márquez^{a,*}

^a Departamento de Biología Animal, Biología Vegetal y Ecología, Universidad de Jaén, Jaén, Spain

^b Departamento de Enfermedades Infecciosas, Hospital Universitario San Pedro-Centro de Investigación Biomédica de La Rioja (CIBIR), Logroño, Spain

^c Departamento de Biología Experimental, Universidad de Jaén, Jaén, Spain

^d Hospital "Arnau de Vilanova", Lleida, Spain

ARTICLE INFO

Article history:

Received 28 August 2020

Accepted 9 October 2020

Available online 15 December 2020

Keywords:

Bartonella henselae

Bartonella quintana

Immunofluorescent test

Iberian Peninsula

ABSTRACT

Introduction: The purpose of this study was to determine the prevalence of IgG antibodies against *Bartonella* sp. in a randomly selected sample from the population of the patients of North Sanitary District of Jaén.

Methods: We used a commercially available immunofluorescent test (Focus-Technology IFA *Bartonella quintana* and *B. henselae* test).

Results: Six hundred five healthy individuals were divided by sex into three age groups. We detected that 13.55% and 11.07% subjects were IgG seropositive to *B. henselae* and *B. quintana*, respectively.

Conclusions: Our data show that the prevalence of both *Bartonella* species in Andalusia (Southern Spain) is relatively high. No statistical difference in the seropositivity was observed among these groups. In both cases, the IgG antibody titers ranged from 1/128 to 1/512.

© 2020 Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Published by Elsevier España, S.L.U. All rights reserved.

Estudio serológico de la infección por *Bartonella* sp. en la población humana del sur de España

RESUMEN

Introducción: El propósito de este estudio fue determinar la prevalencia de anticuerpos IgG frente a *Bartonella* sp. en una muestra escogida al azar de la población de pacientes del Distrito Sanitario Norte de Jaén.

Métodos: Se ha utilizado una prueba de inmunofluorescencia disponible comercialmente (Focus-Technology IFA *Bartonella quintana* y prueba de *B. henselae*). Seiscientos cinco individuos sanos se dividieron por sexo en 3 grupos de edad.

Resultados: Detectamos que el 13,55% y el 11,07% de los sujetos eran IgG seropositivos a *B. henselae* y *B. quintana*, respectivamente. En ambos casos, los títulos de anticuerpos IgG variaron de 1/128 a 1/512.

Conclusión: Nuestros datos muestran que la prevalencia de ambas especies de *Bartonella* en Andalucía (sur de España) es relativamente alta. No se observaron diferencias estadísticas en la seropositividad entre grupos de edad.

© 2020 Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Bartonella henselae

Bartonella quintana

Test de inmunofluorescencia

Península ibérica

Introduction

Bartonella infections have gained considerably importance in international public health during the last decades, mainly due to their worldwide distribution and their morbidity rate among

* Corresponding author.

E-mail address: jmarquez@ujaen.es (F.J. Márquez).

Table 1

Pattern of IgG antibody titers to *B. henselae* and *B. quintana* in the study population (N = 605).

<i>B. henselae</i> titers	<i>B. quintana</i> titers				ΣBh
	≤1/64	1/128	1/256	1/512	
≤1/64	517	4	0	2	523
1/128	18	45	0	0	63
1/256	1	1	10	0	12
1/512	2	1	2	2	7
ΣBq	538	51	12	4	605

human populations. In addition, these bacteria cause a growing clinical spectrum, which often results in a challenge for clinicians due to the limitation of the microbiological tests used for diagnosis.¹

The aim of this paper is to evaluate the prevalence of past infection due to *Bartonella* spp. in Jaén province (Andalusia, Spain).

Methods: Serum samples from 605 healthy individuals (222 men, 383 women) from North Sanitary District of Jaén province (Andalusia) were tested for the presence of IgG antibodies against *Bartonella* by indirect immunofluorescence assay (IFA) using commercially available antigen for *B. henselae* and *B. quintana* (Focus Technologies, Cypress, CA). The average age in the sample was 52.4 years (55.2 for men, rank 3–106 years; 50.8 for women, rank 2–97 years), with a standard deviation of 21 years. The kit for detecting IgG antibodies uses Vero cells infected with either *B. henselae* or *B. quintana*. The serum samples were initially diluted 1/64 for the detection of IgG antibodies. Any serum sample found to be positive at the initial dilution was further titrated. Positive and negative controls were included in each test. Endpoint titers were obtained by serial dilution on positive specimens, with titers ≥ 1/128 considered indicative of past infection to *Bartonella* spp.

For statistical analysis we considered gender and three age groups (less than 19, between 19 and 65, and 66 and more years old). Statistical analysis was performed using statistical tools of Microsoft Office Excel 2007. The Student's *t*-test was used to compare mean ages of subjects seropositive and seronegative for *B. henselae* and *B. quintana*. The chi-square test and Fisher's exact method for small samples were used for comparison of prevalence rates in the subgroups.² Also we compared the results of the serological assays with the results of a previous serological study carried out with the same sera for investigating the seroprevalence against *Rickettsia typhi*.³

Results

We found a global prevalence of infection of 13.55% (82 positives) and 11.07% (67 positives) to *B. henselae* and *B. quintana* respectively, corresponding with 63 (51), 12 (12) and 7 (4) positive samples with titer 1/128, 1/256 and 1/512 respectively for *B. henselae* (and *B. quintana*) (Table 1). We do not find significant differences when we studied the distribution of positives in relation with age and gender. The number of positive serologies considering both *B. henselae* and *B. quintana* was 68 (11.18% over total population). Fig. 1 shows the distribution of the titles obtained in the municipalities of origin of the individuals studied, both for *B. henselae* and *B. quintana*.

When we compared titles against *Bartonella* spp. and *Rickettsia typhi* over the same human serum samples³ over all positive serum (titer ≥ 1/128) for *R. typhi*, *B. henselae* and *B. quintana*, we observed that the 51.7% of positive sera react only with *R. typhi*, 23.86% of serums react against *B. quintana* and *B. henselae*, and 10.79% shows positive reaction against these three pathogens (Table 2).

Discussion

Our study confirms the widespread distribution of *Bartonella* sp. in Jaén province, as reflected by moderate prevalence of past infection due to *Bartonella* spp. agents in a representative sample of general population. Since in our study the titers of antibodies to *B. henselae* and *B. quintana* are very similar we cannot know the real prevalence of both microorganisms due to cross reactions among the different species of *Bartonella*. Only in 4 sera we have found difference more than 2 dilutions among the samples.

Several studies developed in Europe and America found comparable prevalence. In Greece, the observed seroprevalence for IgG antibodies were of 19.8% and 15% to *B. henselae* and to *B. quintana*, respectively.² In a rural area of Brazil, da Costa et al.⁴ found a seroprevalence of 13.7% and 12.8% for *B. henselae* and *B. quintana*. Moreover, in a study conducted on blood donors shows that seropositivity rate of IgG antibodies against *B. henselae* range from of 3.3% (*n* = 333) in Turkey,⁵ 3.61% (*n*: 140), in New Zealand,⁶ 4.4% (*n*: 498) in Sweden⁷ and 11.4% (*n*: 122) in Italy.⁸ In the first studies carry out in Spain (La Rioja) the prevalence of antibodies to *B. henselae* in cat owners, healthy population and HIV patients was respectively 28.9%, 5.9% and 17.3%.^{9,10} In a study carry out in Catalonia, *B. henselae* seroprevalence was signaled as 8.7% in a study carried out on a healthy population.¹¹ Studies developed on veterinarian personnel shows high seroprevalence values than in blood donors or health people (33% for *B. henselae* and 10% for *B. quintana*; *n* = 87).¹² Seroprevalence to *Bartonella quintana* and *Bartonella henselae* among urban homeless and marginalized people in Europe and United States, range from 0–37.5% to 0–10.3% respectively.¹³

Multiple reactivity or cross-reactivity with other rickettsial antigens was frequently observed especially between closely related antigens such as *B. henselae* and *B. quintana*.^{14,15} For this test, Maurin et al.¹⁶ established a specificity of 87%, which can be improved using a cut-off titer of 128. In the case of *B. henselae*, a comparative study that evaluated different serological assays for diagnosis of cat-scratch disease comparing several IgM and IgG IFA test and ELISA test determined in the case of the test used in this study obtained a 98% of sensitivity and a 69% of specificity.¹⁷ Also is very common and well documented cross reactivity with *Coxiella burnetii* infections.¹⁴ Our results suggest an elevated frequency of asymptomatic carriers of antibodies against *Bartonella* spp. among healthy population of studied area. We can also confirm that most part of the infections by those *Bartonella* are subclinical or misdiagnosed. There are few published data regarding the seroprevalence in blood donors in our country, so determining if *B. henselae* is present in the blood of donors at the time of donation is very important, since this microorganism can survive up to 35 days in the red blood cells stored in a blood bank at 4 °C.¹⁸

Our understanding of the transmission of *B. quintana* has changed in recent years. Although body louse is a well-known vector of trench fever, the growth kinetics of *B. quintana* in body lice has not been fully understood. It had been reported that multiplication rate. *B. quintana* started proliferation in body lice 4 days after ingestion and was constantly excreted in the feces for at least 3 weeks.¹⁹ Studies carried on body lice from homeless in San Francisco found a prevalence of *B. quintana* ranging from 12.3% to 33.3%.²⁰ However, they also noted the appearance of this organism in head lice in the presence,²¹ but also in absence of body lice in the sampled individuals.^{20,22} All these questions have been solved when studies aimed at determining the genetic structure of populations based on analysis of mitochondrial genome sequences have been able to determine the existence of up to six distinct genetic clades.^{23,24} Despite having raised different hypotheses,^{23,25} the genetic basis and evolutionary relationships among body and head lice remain obscure.²⁶ The change in the phylogenetic paradigm of lice also determines an adaptation of the epidemiological paradigm,




Fig. 1. Titles obtained for *B. henselae* and *B. quintana* in the municipalities of North Sanitary District of Jaén considered in this study.

Table 2

Pattern of IgG antibody titers $\geq 1/128$ against *R. typhi*, *B. quintana* and *B. henselae* (showed as Rt/Bq/Bh) in the study population ($N=605$).

Key	Men	Women	Total	%/total	%/M	%/W
Rt/-/-	11	14	25	4.13	4.95	3.66
Rt/Bq/-	0	0	0	0.00	0.00	0.00
Rt/-/Bh	1	1	2	0.33	0.45	0.26
Rt/Bq/Bh	5	1	6	0.99	2.25	0.26
-/Bq/-	3	2	5	0.83	1.35	0.52
-/-/Bh	6	7	13	2.15	2.70	1.83
-/Bq/Bh	14	28	42	6.94	6.31	7.31
-/-/-	182	330	512	84.63	81.98	86.16

so that it is currently considered that of both vector ecotypes (head and body lice) may be involved in the transmission of *B. quintana*.^{27,28} This is consistent with the results of several molecular studies detected *B. quintana* DNA in head lice worldwide, usually in people infested with both head and body lice,^{20,21,29} as well as those infected exclusively by head lice.^{30–32} Also it is possible that other vectors such as bed bugs or fleas could be involved in the transmission of *B. quintana*.^{33,34}

Bartonella henselae has been signaled as the etiological agent of Cat scratch disease (CSD)^{35,36} in conjunction of other *Bartonella* species.^{37,38} The main transmission process enroll the scratch or, less likely, by the bite or lick of cats as well as arthropod (cat flea was generally signaled).^{39,40} CSD is characterized by local inflammation and significant enlargement of regional lymph nodes are usually seen within 10–14 days of scratching or biting by a cat. Systemic symptoms such as general discomfort and fever also develop, which can last several weeks. While most cases have a benign course and could resolve spontaneously, in some more severe cases various complications occur (endocarditis, encephalitis, meningitis, osteomyelitis and oculoglandular syndrome).^{41–45}

Several *Bartonella* sp. different to *B. henselae* has been reported in continental Spain on potential vectors and mammal host, concerning rodents,^{46,47} lagomorphs,^{37,48} domestic and wild carnivore,^{49–52} chiropteran,⁵³ ruminants⁵⁴ and to veterinarian personnel working with domestic animals¹² or sanitary workers.⁵⁵ In this scenario we can consider that serologic test alone can

determined the presence of antibodies against *Bartonella* spp., and no to the involved specie or species that should be interpreted in the adequate clinical and epidemiological context.

Anyway although new tools are available for diagnosing *Bartonella* infections such as molecular and culture ones^{12,55} it is necessary to know the prevalence of antibodies to *Bartonella* spp. in a determinate area since serology keep going the more used technique for diagnosing patients with *Bartonella* infections.

Conflicts of interest

The authors declare that they have no conflict of interest.

Acknowledgments

This project was partially supported by a gram from the Universidad de Jaen Research Program (Acción 16).

References

- Breitschwerdt EB. Bartonellosis, one health and all creatures great and small. *Vet Dermatol*. 2017;28, 96–e21.
- Tea A, Alexiou-Daniel S, Arvanitidou M, Diza E, Antoniadis A. Occurrence of *Bartonella henselae* and *Bartonella quintana* in a healthy Greek population. *Am J Trop Med Hyg*. 2003;68:6–554. <http://dx.doi.org/10.4269/ajtmh.2003.68.554>.
- Santibáñez S, Astasio A, Villa-Real R, Cámara JA, Oteo JA, Márquez FJ. Serologic study of *Rickettsia typhi* infection among the human population of southern Spain. *Clin Microbiol Infect*. 2009;15:8–247. <http://dx.doi.org/10.1111/j.1469-0691.2008.02171.x>.
- da Costa PS, Brigatte ME, Greco DB. Antibodies to *Rickettsia rickettsii*, *Rickettsia typhi*, *Coxiella burnetii*, *Bartonella henselae*, *Bartonella quintana* and *Ehrlichia chaffeensis* among healthy population in Minas Gerais, Brazil. *Mem Inst Oswaldo Cruz*. 2005;100:9–853. <http://dx.doi.org/10.1590/s0074-02762005000800006>.
- Aydin N, Bülbül R, Tellı M, Gültekin B. Seroprevalence of *Bartonella henselae* and *Bartonella quintana* in blood donors in Aydin province, Turkey. *Mikrobiyol Bul*. 2014;48:3–477.
- Zarkovic A, McMurray C, Deva N, Ghosh S, Whitley D, Guest S. Seropositivity rates for *Bartonella henselae*, *Toxocara canis* and *Toxoplasma gondii* in New Zealand blood donors. *Clin Exp Ophthalmol*. 2007;35:4–131. <http://dx.doi.org/10.1111/j.1442-9071.2006.01406.x>.
- McGill S, Wesslen L, Hjelm E, Holmberg M, Auvinen MK, Berggren K, et al. *Bartonella* spp. seroprevalence in healthy Swedish blood donors. *Scand J Infect Dis*. 2005;37:30–723. <http://dx.doi.org/10.1080/00365540510012152>.

8. Mansueto P, Ilenia Pepe, Cillari E, Arcoleo F, Micalizzi A, Bonura F, et al. Prevalence of antibodies anti-*Bartonella henselae* in Western Sicily: children, blood donors, and cats. *J Immunol* 2012;188:23:18–25, <http://dx.doi.org/10.1080/15321819.2011.591476>.
9. Blanco-Ramos JR, Oteo-Revuelta JA, Martínez de Artola V, Ramalle-Gómara E, García-Pineda A, Ibarra-Cucalón V. Seroepidemiology of *Bartonella henselae* infection in a risk Group. *Rev Clin Esp*. 1998;198:9–805.
10. Blanco JR, Oteo JA, Martínez V, Ramalle E, García A, Ibarra V, et al. Seroepidemiology of *Bartonella henselae* infection in HIV-infected patients. *Enferm Infecc Microbiol Clin*. 1999;17:8–434.
11. Pons I, Sanfeliu I, Cardenosa N, Noguera MM, Font B, Segura F. Serological evidence of *Bartonella henselae* infection in healthy people in Catalonia, Spain. *Epidemiol Infect*. 2008;25:1–5, <http://dx.doi.org/10.1017/S0950268808000368>.
12. Oteo JA, Maggi R, Portillo A, Bradley J, García-Álvarez L, San-Martín M, et al. Prevalence of *Bartonella* spp. by culture, PCR and serology, in veterinary personnel from Spain. *Parasit Vect*. 2017;10:553, <http://dx.doi.org/10.1186/s13071-017-2483-z>.
13. Leibler JH, Zakhour CM, Gadhoke P, Gaeta JM. Zoonotic and vector-borne infections among urban homeless and marginalized people in the United States and Europe, 1990–2014. *Vect-Borne Zoonotic Dis*. 2016;16:4–435.
14. La Scola B, Raoult D. Serological cross-reactions between *Bartonella quintana*, *Bartonella henselae*, and *Coxiella burnetii*. *J Clin Microbiol*. 1996;34:4–2270.
15. Maurin M, Eb F, Etienne J, Raoult D. Serological crossreactions between *Bartonella* and *Chlamydia* species: implications for diagnosis. *J Clin Microbiol*. 1997;35:7–2283.
16. Maurin M, Rolain JM, Raoult D. Comparison of in-house and commercial slides for detection by immunofluorescence of immunoglobulins G and M against *Bartonella henselae* and *Bartonella quintana*. *Clin Diagn Lab Immunol*. 2002;9:9–1004.
17. Vermeulen MJ, Verbakel H, Notermans DW, Reimerink JH, Peeters MF. Evaluation of sensitivity, specificity and cross-reactivity in *Bartonella henselae* serology. *J Med Microbiol*. 2010;59:5–743, <http://dx.doi.org/10.1099/jmm.0.015248-0>.
18. Magalhães RF, Pitassi LU, Salgado M, De Moraes AM, Barjas-Castro ML, Velho PENF. *Bartonella henselae* survives after the storage period of red blood cell units: is it transmissible by transfusion? *Tranfus Med*. 2008;18:91–287, <http://dx.doi.org/10.1111/j.1365-3148.2008.00871.x>.
19. Seki N, Kasai S, Saito N, Komagata O, Mihara M, Sasaki T, et al. Quantitative analysis of proliferation and excretion of *Bartonella quintana* in body lice, *Pediculus humanus* L. *Am J Trop Med Hyg*. 2007;77:6–562.
20. Bonilla DL, Kabeya H, Henn J, Kramer VL, Kosoy MY. *Bartonella quintana* in body lice and head lice from homeless persons, San Francisco, California, USA. *Emerg Infect Dis*. 2009;15:5–912, <http://dx.doi.org/10.3201/eid1506.0900054>.
21. Sasaki-Fukutsu K, Koga R, Nikoh N, Yoshizawa K, Kasai S, Mihara M, et al. Symbiotic bacteria associated with stomach discs of human lice. *Appl Environ Microb*. 2006;72:52–7349, <http://dx.doi.org/10.1128/AEM.01429-06>.
22. Sangaré AK, Boutellis A, Drali R, Socolovschi C, Barker SC, Diatta G, et al. Detection of *Bartonella quintana* in African body and head lice. *Am J Trop Med Hyg*. 2014;91:294–301, <http://dx.doi.org/10.4269/ajtmh.13-0707>.
23. Drali R, Boutellis A, Raoult D, Rolain JM, Brouqui P. Distinguishing body lice from head lice by multiplex real-time PCR analysis of the Phum.PHUM540560 gene. *PLoS One*. 2013;8:e58088, <http://dx.doi.org/10.1371/journal.pone.0058088>.
24. Boutellis A, Abi-Rached L, Raoult D. The origin and distribution of human lice in the world. *Infect Genet Evol*. 2014;23:17–209, <http://dx.doi.org/10.1016/j.meegid.2014.01.017>.
25. Li W, Ortiz G, Fournier PE, Gimenez G, Reed DL, Pittendrigh B, et al. Genotyping of human lice suggests multiple emergencies of body lice from local head louse populations. *PLoS Negl Trop Dis*. 2010;4:e641, <http://dx.doi.org/10.1371/journal.pntd.0000641>.
26. Amanzougaghene N, Fenollar F, Raoult D, Mediannikov O. Where are we with human lice? A review of the current state of knowledge. *Front Cell Infect Microbiol*. 2019;9:474, <http://dx.doi.org/10.3389/fcimb.2019.00474>.
27. Amanzougaghene N, Fenollar F, Sangaré AK, Sissoko MS, Doumbo OK, Raoult D, et al. Detection of bacterial pathogens including potential new species in human head lice from Mali. *PLoS One*. 2017;12:e0184621, <http://dx.doi.org/10.1371/journal.pone.0184621>.
28. Louni M, Mana N, Bitam I, Dahmani M, Parola P, Fenollar F, et al. Body lice of homeless people reveal the presence of several emerging bacterial pathogens in northern Algeria. *PLoS Negl Trop Dis*. 2018;12:e0006397, <http://dx.doi.org/10.1371/journal.pntd.0006397>.
29. Koehler JE, Sanchez MA, Garrido CS, Whitfield MJ, Chen FM, Berger TG, et al. Molecular epidemiology of *Bartonella* infections in patients with bacillary angiomatosis-peliosis. *N Engl J Med*. 1997;337:83–1876, <http://dx.doi.org/10.1056/NEJM199712253372603>.
30. Angelakis E, Rolain JM, Raoult D, Brouqui P. *Bartonella quintana* in head louse nits. *FEMS Immunol Med Microbiol*. 2011;62:6–244, <http://dx.doi.org/10.1111/j.1574-695X.2011.00804.x>.
31. Boutellis A, Veracx A, Angelakis E, Diatta G, Mediannikov O, Trape J-F. *Bartonella quintana* in head lice from Senegal. *Vector-Borne Zoonotic Dis*. 2012;12:7–564, <http://dx.doi.org/10.1089/vbz.2011.0845>.
32. Diatta G, Mediannikov O, Sokhna C, Bassene H, Socolovschi C, Ratmanov P, et al. Prevalence of *Bartonella quintana* in patients with fever and head lice from rural areas of Sine-Saloum, Senegal. *Am J Trop Med Hyg*. 2014;91:3–291, <http://dx.doi.org/10.4269/ajtmh.13-0685>.
33. Kernif T, Leulmi H, Socolovschi C, Berenger JM, Lepidi H, Bitam I, et al. Acquisition and excretion of *Bartonella quintana* by the cat flea, *Ctenocephalides felis felis*. *Mol Ecol*. 2014;23:12–204, <http://dx.doi.org/10.1111/mec.12663>.
34. Leulmi H, Bitam I, Berenger JM, Lepidi H, Rolain JM, Almeras L, et al. Competence of *Cimex lectularius* bed bugs for the transmission of *Bartonella quintana*, the agent of trench fever. *PLoS Negl Trop Dis*. 2015;9:e0003789, <http://dx.doi.org/10.1371/PMC>.
35. Welch DF, Pickett DA, Slater LN, Steigerwalt AG, Brenner DJ. *Rochalimaea henselae* sp. nov., a cause of septicemia, bacillary angiomatosis, and parenchymal bacillary peliosis. *J Clin Microbiol*. 1992;30:275–80.
36. Regnery RL, Anderson BE, Clarridge JE III, Rodriguez-Barradas M, Jones DC, Carr JH. Characterization of a novel *Rochalimaea* species *R. henselae* sp. nov., isolated from blood of a febrile, human immunodeficiency virus-positive patient. *J Clin Microbiol*. 1992;30:265–74.
37. Koehler JE, Glaser CA, Tappero JW. *Rochalimaea henselae* infection. A new zoonosis with the domestic cat as reservoir. *JAMA*. 1994;271:531–5.
38. Rolain JM, Franc M, Davoust B, Raoult D. Molecular detection of *Bartonella quintana*, *B. koehlerae*, *B. henselae*, *B. clarridgeiae*, *Rickettsia felis*, and *Wolbachia pipiensis* in cat fleas, France. *Emerg Infect Dis*. 2003;9:338–42, <http://dx.doi.org/10.3201/eid0903.020278>.
39. Chomel BB, Kasten RW, Floyd-Hawkins K, Chi B, Yamamoto K, Roberts-Wilson J, et al. Experimental transmission of *Bartonella henselae* by the cat flea. *J Clin Microbiol*. 1996;34:1952–6.
40. Finkelstein JL, Brown TP, O'Reilly KL, Wedincamp J Jr, Foil LD. Studies on the growth of *Bartonella henselae* in the cat flea (*Siphonaptera: Pulicidae*). *J Med Entomol*. 2002;39:915–9, <http://dx.doi.org/10.1603/0022-2585-39.6.915>.
41. Florin TA, Zaoutis TE, Zaoutis LB. Beyond cat scratch disease: widening spectrum of *Bartonella henselae* infection. *Pediatrics*. 2008;121:e1413–25, <http://dx.doi.org/10.1542/peds.2007-1897>.
42. Nelson CA, Saha S, Mead PS. Cat-scratch disease in the United States, 2005–2013. *Emerg Infect Dis*. 2016;22:1741–6, <http://dx.doi.org/10.3201/eid2210.160115>.
43. Aziz HA, Plessek TP, Sabella C, Udayasankar UK, Singh AD. Cat scratch disease: expanded spectrum. *Ocul Oncol Pathol*. 2016;2:246–50, <http://dx.doi.org/10.1159/000447063>.
44. Okaro U, Addisu A, Casanas B, Anderson B. *Bartonella* species, an emerging cause of blood-culture-negative endocarditis. *Clin Microbiol Rev*. 2017;30:709–46, <http://dx.doi.org/10.1128/CMR.00013-17>.
45. Canneti B, Cabo-López I, Puy-Núñez A, García JG, Cores FJ, Trigo M, et al. Neurological presentations of *Bartonella henselae* infection. *Neurol Sci*. 2019;40:261–8, <http://dx.doi.org/10.1007/s10072-018-3618-5>.
46. Márquez FJ, Rodríguez-Liébana JJ, Pachón-Ibáñez ME, Docobo-Pérez F, Hidalgo-Fontiveros A, Bernabeu-Wittel M, et al. Molecular screening of *Bartonella* species in rodents from south western Spain. *Vector Borne Zoonotic Dis*. 2008;8:695–700, <http://dx.doi.org/10.1089/vbz.2007.0257>.
47. Gil H, García-Esteban C, Barandika JF, Peig J, Toledo A, Escudero R, et al. Variability of *Bartonella* genotypes among small mammals in Spain. *Appl Environ Microbiol*. 2010;76:8062–70, <http://dx.doi.org/10.1128/AEM.01963-10>.
48. Márquez FJ. Molecular detection of *Bartonella alsatica* in European wild rabbits (*Oryctolagus cuniculus*) in Andalusia (Spain). *Vector Borne Zoonotic Dis*. 2010;10:4–731, <http://dx.doi.org/10.1089/vbz.2009.0135>.
49. Roura X, Breitschwerdt E, Lloret A, Ferrer L, Hegarty B. Serological evidence of exposure to *Rickettsia*, *Bartonella*, and *Ehrlichia* species in healthy or Leishmania infantum-infected dogs from Barcelona, Spain. *J Appl Res Vet Med*. 2005;3:37–129.
50. Blanco JR, Pérez-Martínez L, Vallejo M, Santibáñez S, Portillo A, Oteo JA. Prevalence of *Rickettsia felis*-like and *Bartonella* spp. in *Ctenocephalides felis* and *Ctenocephalides canis* from La Rioja (Northern Spain). *Ann N Y Acad Sci*. 2006;1078:4–270, <http://dx.doi.org/10.1196/annals.1374.050>.
51. Márquez FJ, Millán J, Rodríguez-Liebana JJ, García-Egea I, Muniaín MA. Detection and identification of *Bartonella* sp. in fleas from carnivorous mammals in Andalusia, Spain. *Med Vet Entomol*. 2009;23:8–393, <http://dx.doi.org/10.1111/j.1365-2915.2009.00830.x>.
52. Gerrikagoitia X, Gil H, García-Esteban C, Anda P, Juste RA, Barral M. Presence of *Bartonella* species in wild carnivores of northern Spain. *Appl Environ Microbiol*. 2012;78:8–885, <http://dx.doi.org/10.1128/AEM.05938-11>.
53. Stuckey MJ, Boulouis HJ, Cliquet F, Picard-Meyer E, Servat A, Aréchiga-Ceballos N, et al. Potentially zoonotic *Bartonella* in bats from France and Spain. *Emerg Infect Dis*. 2017;23:539, <http://dx.doi.org/10.3201/eid2303.160934>.
54. Antequera-Gómez ML, Lozano-Almendral L, Barandika JF, González-Martín-Niño RM, Rodríguez-Moreno I, García-Pérez A, et al. *Bartonella chomelii* is the most frequent species infecting cattle grazing in communal mountain pastures in Spain. *Appl Environ Microbiol*. 2015;81:9–623, <http://dx.doi.org/10.1128/AEM.03159-14>.
55. Portillo A, Maggi R, Oteo JA, Bradley J, García-Álvarez L, San-Martín M, et al. *Bartonella* spp. prevalence (serology, culture, and PCR) in sanitary workers in La Rioja Spain. *Pathogens*. 2020;9:189, <http://dx.doi.org/10.3390/pathogens9030189>.