



# NEUROLOGÍA

[www.elsevier.es/neurologia](http://www.elsevier.es/neurologia)



## ORIGINAL ARTICLE

# Evaluation of the anticonvulsant and neuroprotective effect of intracerebral administration of growth hormone in rats

I. Zamora-Bello<sup>a</sup>, A. Martínez<sup>b</sup>, L. Beltrán-Parrazal<sup>c</sup>, I. Santiago-Roque<sup>d</sup>,  
E. Juárez-Aguilar<sup>e</sup>, M.L. López-Meraz<sup>f,\*</sup>

<sup>a</sup> Centro de Investigaciones Cerebrales, Universidad Veracruzana, Médicos y Odontólogos s/n, Col. Unidad del Bosque Xalapa, Veracruz C.P. 91010, Mexico

<sup>b</sup> Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, Calzada México Xochimilco No. 101, Col. San Lorenzo Huipulco, Tlalpan, Ciudad de México C.P. 14370, Mexico

<sup>c</sup> Centro de Investigaciones Cerebrales, Universidad Veracruzana, Médicos y Odontólogos s/n, Col. Unidad del Bosque Xalapa, Veracruz C.P. 91010, Mexico

<sup>d</sup> Laboratorio de Neurotoxicología, Universidad Veracruzana, Médicos y Odontólogos s/n, Col. Unidad del Bosque Xalapa, Xalapa, Veracruz C.P. 91010, Mexico

<sup>e</sup> Instituto de Ciencias de la Salud, Universidad Veracruzana, Av. Dr. Luis Castelazo Ayala s/n, Col. Industrial Animas, Xalapa, Veracruz C.P. 91190, Mexico

<sup>f</sup> Centro de Investigaciones Cerebrales, Universidad Veracruzana, Médicos y Odontólogos s/n, Col. Unidad del Bosque Xalapa, Veracruz C.P. 91010, Mexico

Received 13 November 2020; accepted 1 March 2021

## KEYWORDS

Growth hormone;  
*Status epilepticus*;  
Neurodegeneration;  
Pilocarpine

## Abstract

**Introduction:** The growth hormone (GH) has been reported as a crucial neuronal survival factor in the hippocampus against insults of diverse nature. *Status epilepticus* (SE) is a prolonged seizure that produces extensive neuronal cell death. The goal of this study was to evaluate the effect of intracerebroventricular administration of GH on seizure severity and SE-induced hippocampal neurodegeneration.

**Methodology:** Adult male rats were implanted with a guide cannula in the left ventricle and different amounts of GH (70, 120 or 220 ng/3  $\mu$ l) were microinjected for 5 days; artificial cerebrospinal fluid was used as the vehicle. Seizures were induced by the lithium–pilocarpine model (3 mEq/kg LiCl and 30 mg/kg pilocarpine hydrochloride) one day after the last GH administration. Neuronal injury was assessed by Fluoro-Jade B (F-JB) staining.

**Results:** Rats injected with 120 ng of GH did not have SE after 30 mg/kg pilocarpine, they required a higher number of pilocarpine injections to develop SE than the rats pretreated with the vehicle, 70 ng or 220 ng GH. Prefrontal and parietal cortex EEG recordings confirmed that

\* Corresponding author.

E-mail address: [leonorlopez@uv.mx](mailto:leonorlopez@uv.mx) (M.L. López-Meraz).

<https://doi.org/10.1016/j.nrl.2021.03.010>

0213-4853/© 2021 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: I. Zamora-Bello, A. Martínez, L. Beltrán-Parrazal et al., Evaluation of the anticonvulsant and neuroprotective effect of intracerebral administration of growth hormone in rats, *Neurología*, <https://doi.org/10.1016/j.nrl.2021.03.010>

I. Zamora-Bello, A. Martínez, L. Beltrán-Parrazal et al.

latency to generalized seizures and SE was also significantly higher in the 120 ng group when compared with all the experimental groups. FJ-B positive cells were detected in the hippocampus after SE in all rats, and no significant differences in the number of F-JB cells in the CA1 area and the hilus was observed between experimental groups.

**Conclusion:** Our results indicate that, although GH has an anticonvulsive effect in the lithium–pilocarpine model of SE, it does not exert hippocampal neuroprotection after SE.

© 2021 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## PALABRAS CLAVE

Hormona de crecimiento;  
*Status epilepticus*;  
Neurodegeneración;  
Pilocarpina

## Evaluación del efecto anticonvulsivante y neuroprotector de la administración intracerebral de la hormona de crecimiento en ratas

### Resumen

**Introducción:** La hormona de crecimiento (HC) es un factor que favorece la supervivencia neuronal en el hipocampo ante agresiones de diversa naturaleza. El *status epilepticus* (SE) es un tipo de crisis epiléptica de larga duración que produce muerte neuronal. El objetivo de este estudio fue evaluar el efecto de la administración intracerebroventricular de HC en la severidad de las convulsiones y la neurodegeneración hipocampal debida al SE.

**Metodología:** A ratas macho adultas se les implantó una cánula guía en el ventrículo lateral izquierdo y se les microinyectaron diferentes cantidades de HC (70, 120 o 220 ng/3 µl) durante 5 días; como vehículo se inyectó líquido cefalorraquídeo artificial. Las convulsiones se generaron con el modelo de litio-pilocarpina (3 mEq/kg LiCl y 30 mg/kg clorhidrato pilocarpina) un día después de la última inyección de HC. La neurodegeneración se identificó con la tinción de Fluoro-Jade B (F-JB).

**Resultados:** Las ratas a las que se les inyectaron 120 ng de HC requirieron 2 o 3 inyecciones de pilocarpina para desarrollar SE, en comparación con el resto de los grupos experimentales que requirieron solo una aplicación del convulsivante. Los registros EEG de la corteza prefrontal y parietal confirmaron que la latencia a las crisis generalizadas y al SE fue mayor en dicho grupo experimental. Todas las ratas con SE presentaron células positivas al FJ-B en el área CA1 e hilus del hipocampo, y no se identificaron diferencias entre los tratamientos.

**Conclusión:** Nuestros resultados muestran que, aunque la HC tiene un efecto anticonvulsivante, una vez que se ha desarrollado el SE no promueve neuroprotección en el hipocampo.

© 2021 Sociedad Española de Neurología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The growth hormone (GH) is produced by the adenohypophysis and its most reported functions include body growth and lipid and protein metabolism (Moller and Jorgensen, 2009). In the Central Nervous System (CNS), GH promotes cell proliferation and survival of neurons and neural progenitors,<sup>1,2</sup> brain growth, myelination, neuronal arborization, glial differentiation<sup>3,4</sup> and spinogenesis.<sup>5</sup> Interestingly, GH has been reported as an endogenous relevant factor in the hippocampus where it is locally secreted and regulated by age, sex and stress as well as by insults of different nature.<sup>6–9</sup> Thus, exogenous GH administration exerts neuroprotective effect in this brain structure against aging, hypoxia-ischemia and sleep deprivation insults.<sup>3,10–12</sup>

On the other hand, *status epilepticus* (SE) is a non-self-limited type of epileptic seizure characterized by an enduring epileptic state.<sup>13</sup> In humans and experimental rats, SE produces neuronal cell death in different brain

regions including the hippocampus, the piriform cortex, the mediodorsal thalamic nucleus, the amygdala, and the cerebellum.<sup>14–16</sup> Although some alterations in the serum levels of GH have been reported during seizures,<sup>17,18</sup> information regarding the effect of GH on epilepsy or its possible neuroprotective effect on SE-induced neuronal damage is limited. In a study by Kato, intrahippocampal administration of GH in mice enhanced the progression of amygdala kindling, whereas pegvisomant, an antagonist of GH receptors, delayed it.<sup>19</sup> However, it has not been explored whether GH could protect against neuronal damage induced by SE. Therefore, the main goal of this study was to evaluate the effect of intracerebroventricular (i.c.v.) administration of GH on hippocampal neuronal injury in the lithium–pilocarpine model of SE. We performed a detailed evaluation of the GH effect on seizure severity considering that this may affect the subsequent neuronal damage. This study expands our knowledge on the effects of GH in the hippocampus, and more specifically on epilepsy.

## Materials and methods

### Animals

This study was carried out in strict accordance with the Mexican guidelines on the care and use of laboratory animals (NOM-062-ZOO-1999), the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and under the ARRIVE guidelines.<sup>20</sup> Internal Committee for the Care and Use of Laboratory Animals of the Instituto de Ciencias de la Salud at Universidad Veracruzana approved this protocol (CICUAL-ICS 2017-0002). All efforts were made to minimize the number of animals used and their suffering. Male rats Wistar were obtained from the facilities of the Centro de Investigaciones Cerebrales (Universidad Veracruzana) or the Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz. Rats were housed in groups of 4 and kept in environmental conditions of temperature and humidity (20–28 °C and 40–70% RH, respectively), with light–dark cycles of 12/12 h (07:00–19:00) and free access to water and food.

### Surgical procedures

All surgical procedures were performed entirely in aseptic conditions and under anesthesia with isoflurane (1.5–2%, Vedco, Inc., USA). Rats were stereotaxically implanted with a stainless-steel guide cannula (gauge 21, C313G/SPC Plastics One, Roanoke, VA, USA) in the left lateral ventricle (AP = 1.4 mm, ML = 2 mm, DV = –4.3 mm, relative to bregma).<sup>21</sup> For rats that underwent Electroencephalogram (EEG) recording, two nail-shaped stainless steel electrodes were also implanted on the prefrontal (PFC) and parietal cortices (PC). The body temperature of the rats was always controlled with a temperature regulation system (FHC brand, model 41-90-D8) and after the surgery, rats were rehydrated with glucose saline (equivalent to 5% body weight, s.c.). An analgesic (Meglumine, 2.5 mg/kg s.c., for 2 days) and an antimicrobial (Enrofloxacin, 5 mg/kg s.c., for 5 days) were applied in the postoperative period. Rats underwent experimental protocols 5 days after surgery. Only rats correctly implanted in the left lateral ventricle were included in the study.

### Intracerebroventricular microinjection of growth hormone

For the dose–response curve experiments, the GH (Lilly laboratories) was administered through the guide cannula using a microinjector connected to an insulin syringe (BD ultra-Fine, 1 ml), which was placed in a microperfusion pump (Pump 11 Elite, Harvard Apparatus) with a flow rate of 0.3  $\mu$ l/min. Three different concentrations of GH were tested: 70 ng/3  $\mu$ l ( $n$  = 5), 120 ng/3  $\mu$ l ( $n$  = 10) and 220 ng/3  $\mu$ l ( $n$  = 6). The control group received 3  $\mu$ l of ACSF as the vehicle ( $n$  = 8). Each rat was treated with a single dose of GH or the vehicle, which was injected daily during five consecutive days. SE was induced the following day after the last GH administration and the behavioral manifestation of seizures was evaluated. For the EEG recordings, two

additional groups of rats were injected with 120 ng/3  $\mu$ l ( $n$  = 5) or vehicle ( $n$  = 8) as described before.

### Induction of *status epilepticus* with lithium–pilocarpine

All rats were treated with lithium chloride (3 mEq/kg i.p.) and 24 h later, they were injected with pilocarpine hydrochloride (30 mg/kg s.c.) to induce SE or saline solution for control rats (volume of administration 1 ml/kg). If SE was not induced after sixty min of the first pilocarpine injection, the animals were given one or two additional doses of pilocarpine to generate it. The seizures were stopped 1 h after the SE onset with diazepam (10 mg/kg i.p.) and rats were rehydrated with glucose saline. The behavioral manifestation of seizures in the presence and absence of GH were video-recorded and evaluated according to the Racine scale.<sup>22</sup> The parameters evaluated included the number of injections of pilocarpine required for induction of SE and seizure latency, duration and frequency. At the end of these experiments, rats were processed as indicated below for brain histology.


EEG recordings were performed using an analog EEG amplifier Grass Model 7 (Astro Med., Warwick, USA, 2002). Signals were amplified and band-pass filtered (1–100 Hz), acquired and digitally stored for later detailed analysis with GAMMA, Grass data acquisition software, version 3.1 (Astro Med., Warwick, USA, 2003). The EEG was continuously obtained in basal conditions (a baseline of 20 min before pilocarpine injection) and 4 h after the first injection of pilocarpine. Latency and duration to the first generalized seizure and spike frequency during generalized seizures were analyzed in both PC and PFC. To evaluate the frequency of epileptic spikes, only those with an amplitude at least three times above the baseline were considered. At the end of the EEG experiments, rats were killed by an overdose of sodium pentobarbital and the implant site was verified histologically.

### Tissue processing for histology

Twenty-four hours after SE, rats were anesthetized with sodium pentobarbital (100 mg/kg i.p.) and transcardially perfused with 0.9% NaCl, followed by 4% paraformaldehyde (in 0.1 M phosphate solution, pH 7.4). Brains were post-fixed in 4% paraformaldehyde for 2 h and processed for paraffin embedding. Sequential coronal sections (10  $\mu$ m thickness) were obtained at the level of the dorsal hippocampus using a microtome (Leica) and mounted on gelatin-coated slides.

### Fluoro-jade B staining

Degenerative cells were identified by Fluoro-Jade B (F-JB) staining as described previously.<sup>23</sup> Sections were analyzed with a fluorescence microscope (Olympus AX70) at an excitation wavelength of 450–490 nm. A semi-quantitative analysis to count F-JB positive (F-JB+) cells per mm<sup>2</sup> was performed in 4 consecutive coronal sections for each animal in both hemispheres. Cell counting in the dorsal hippocampus was performed in the CA1 pyramidal layer and the



**Figure 1** Effect of growth hormone (GH) on *status epilepticus* (SE) induced by lithium–pilocarpine. Data are shown as the percentage of rats that received 1 or more (2 or 3) injections of pilocarpine (30 mg/kg per administration) to induce SE, and were analyzed by a proportion test. \* $p < 0.05$ .

hilus. Regions of interest were photographed, and cell counting was performed using the open-source platform for biological-image analysis Fiji.

### Statistical analysis

Seizure and histology data were analyzed using a one-way ANOVA to identify differences between the experimental groups and subsequently, when necessary, a Tukey post hoc test was performed. Data from EEG recordings were analyzed with a *Student's t-test* to identify differences between groups. To determine the difference between two proportions, it was used a proportion test. Graph Pad Prism 6.0 or Statistica 7.0 software were used for statistical analysis. A  $p$ -value  $< 0.05$  was considered statistically significant.

## Results


### Seizure severity

Our results unexpectedly showed that any of the rats pre-treated with 120 ng GH had SE after 30 mg/kg of pilocarpine. Thus, all rats in the group of 120 ng GH required 2 or 3 injections of pilocarpine to develop SE, a proportion significantly greater than the rest of the experimental groups (0–16%) ( $p < 0.05$ ) (Fig. 1). Consequently, the latency to the first generalized seizure was larger in the 120 ng GH group ( $79.8 \pm 3.3$  min) compared to the

rest of the experimental groups (ACSF =  $44.4 \pm 6.1$  min; 70 ng GH =  $50.5 \pm 2.4$  min; 220 ng GH =  $46.2 \pm 6.3$  min) ( $F(3, 25) = 14, p = 0.0001$ ). Similarly, the latency to SE was greater in the group of 120 ng GH ( $89.2 \pm 2.9$  min) compared to the other groups (ACSF =  $50.8 \pm 5.4$  min; 70 ng GH =  $55.6 \pm 2.7$ ; 220 ng GH =  $52.5 \pm 6.3$  min) ( $F(3, 25) = 19.7, p = 0.0001$ ). No differences were observed among experimental groups in the duration of the first generalized seizure (ACSF =  $29.3 \pm 3.1$  s; 70 ng GH =  $38.4 \pm 6.8$  s, 120 ng GH =  $31.9 \pm 4.1$  s; 220 ng GH =  $44.0 \pm 9.7$  s) ( $F(3, 25) = 1.3, p = 0.3$ ) or of the SE (ACSF =  $5.6 \pm 0.3$  h; 70 ng GH =  $5.6 \pm 0.2$  h; 120 ng GH =  $4.6 \pm 0.3$  h; 220 ng GH =  $5.5 \pm 0.4$  h) ( $F(3, 24) = 2.6, p = 0.1$ ); all experimental groups had a similar number of generalized seizures until the application of diazepam (Stage IV seizures: ACSF =  $2.6 \pm 0.5$ ; 70 ng GH =  $2.4 \pm 0.4$ ; 120 ng GH =  $1.6 \pm 0.3$ ; 220 ng GH =  $2.8 \pm 0.5$ ) ( $F(3, 25) = 1.9, p = 0.2$ ); Stage V seizures: ACSF =  $2.8 \pm 0.3$ ; 70 ng GH =  $2.6 \pm 0.4$ ; 120 ng GH =  $2.8 \pm 0.2$ ; 220 ng GH =  $2.3 \pm 0.2$ ) ( $F(3, 25) = 0.7, p = 0.6$ ) and no differences in mortality were observed.

EEG recordings confirmed that rats pre-treated with 120 ng GH required 2 or 3 injections of pilocarpine to develop SE, and that latency to the first electrographic seizure and SE was significantly higher in the GH group ( $84.7 \pm 6.7$  and  $102.2 \pm 4.5$  min, respectively) than in the ACSF group ( $31.8 \pm 3.9$  and  $43.4 \pm 3.9$  min, respectively) ( $t = 6.4, 9, p < 0.05$ , and  $t = 9.5, 9, p < 0.05$ , respectively). No difference in the duration of the first EEG generalized seizure was observed between GH group ( $34.8 \pm 7.5$  s) and vehicle group ( $31.3 \pm 6.8$  s) ( $t = 0.34, 7, p > 0.05$ ). Finally,





**Figure 2** Effect of growth hormone (GH, 120 ng) on electrographic activity recorded from Prefrontal (PFC) and Parietal (PC) cortex during the *status epilepticus* (SE) induced by lithium–pilocarpine. (A) Basal line, (B) Electrographic activity 45 min after the first application of pilocarpine (the rat injected with artificial cerebrospinal fluid [ACSF] is already in SE), (C) Electrographic activity 85 min after the first application of pilocarpine (at this time, both rats have electrographic and behavioral SE). Graphs on the right panel show the spike frequency during the first generalized electrographic seizure recorded in the PFC and PC after the injection of ACSF or GH. Data are expressed as the mean  $\pm$  S.E.M. and were analyzed by a *Student's t-test*. No differences were identified between experimental groups. Scale bars: 5 s (horizontal) and 300  $\mu$ V (vertical).

120 ng GH treatment did not affect the spike frequency during generalized seizures either in the PC ( $3.6 \pm 0.4$  Hz) or in the PFC ( $3.6 \pm 0.9$  Hz) when compared with the vehicle ( $3.5 \pm 0.5$  Hz and  $3.1 \pm 0.3$  Hz, respectively) (PC,  $t=0.1$ , 8,  $p=0.8$ ; PFC,  $t=0.5$ , 8,  $p=0.1$ ) (Fig. 2).

### Hippocampus neurodegeneration after SE


FJ-B+ cells were detected in the hippocampus in all rats in the CA1 area and the hilus, but not in CA2, CA3 or dentate gyrus. No significant differences in the number of F-JB+ cells/mm<sup>2</sup> in CA1 were observed between the groups, either in the ipsilateral (ASCF =  $428.5 \pm 101.5$ ; 70 ng GH =  $478 \pm 150.1$ ; 120 ng GH =  $779.5 \pm 95.1$ ; 220 ng GH =  $862.1 \pm 187.1$ ) ( $F(3, 25)=2.9$ ,  $p=0.05$ ) or in the contralateral hemisphere (ASCF =  $493.9 \pm 129.9$ ; 70 ng GH =  $457.8 \pm 104.7$ ; 120 ng GH =  $715.8 \pm 78.3$ ; 220 ng GH =  $955.6 \pm 194.0$ ) ( $F(3, 25)=2.7$ ,  $p=0.07$ ) (Fig. 3). The number of F-JB+ cells/mm<sup>2</sup> in the hilus was similar after all the treatments and for both hemispheres, ipsilateral (ASCF =  $13.8 \pm 4$ ; 70 ng GH =  $22.3 \pm 5.6$ ; 120 ng GH =  $27.5 \pm 5$ , 220 ng GH =  $20.2 \pm 8.2$ ) ( $F(3, 25)=1.2$ ,  $p=0.3$ ) and contralateral to the site of injection (ASCF =  $24 \pm 3.4$ ; 70 ng GH =  $32.4 \pm 7.6$ , 120 ng GH =  $42.2 \pm 7.5$ , 220 ng GH =  $22 \pm 8.62$ ) ( $F(3, 25)=2$ ,  $p=0.1$ ) (Fig. 3).

### Discussion

Our study shows that sub-chronic i.c.v. administration of 120 ng GH has an anticonvulsant effect against SE induced by pilocarpine; however, once SE is triggered, the pattern of behavioral and electrographic seizures and the subsequent hippocampal neurodegeneration are not affected.

A growing body of evidence suggest a central role of the GH in the hippocampal function. Interestingly, the expression of this hormone and its receptor are locally up regulated in this brain region by insults of different nature, while GH exogenous administration counteracts their negative effects.<sup>3,11,12</sup> The present study aimed to analyze the effect of the GH on SE and the subsequent neurodegeneration in the hippocampus. As experimental strategy, we decided to deliver the GH through i.c.v. microinjection to avoid pharmacokinetic issues associated to the systemic administration, and as a potential therapeutic route.<sup>24</sup> Moreover, since it has been reported that daily injection of GH increases overall transcript abundance in the hippocampus more than continuous infusion,<sup>25</sup> GH was administrated five days before the SE insult to evaluate the potential anticonvulsant and hippocampal neuroprotective effect of GH, as previously described.<sup>11</sup>

The role of GH on epilepsy and seizures has not been deeply investigated. The work by Kato and colleagues<sup>19</sup> showed a pro-convulsive effect of GH in the electrical



**Figure 3** Effect of growth hormone (GH) on hippocampal neurodegeneration after *status epilepticus* (SE) induced by lithium–pilocarpine. Microphotographs show Fluoro-Jade B positive (F-JB+) cells (in white) in the dorsal hippocampus of rats treated with artificial cerebrospinal fluid (ACSF) (upper left side panels) or GH (70, 120, or 220 ng) (upper right side panels). Arrowheads in high and low magnification microphotographs point to F-JB+ cells in CA1 and the hilus. Original pictures were converted to 8-bit greyscale images with Fiji. The graphs represent the number of FJ-B+ cells/mm<sup>2</sup> in CA1 and the hilus for each experimental group. Data are expressed as the mean  $\pm$  S.E.M. and were analyzed by a one-way ANOVA. No difference between experimental groups was identified.

kindling model of epilepsy. In that study, chronic administration of GH (12 mg/ml) in the mouse hippocampus during the induction of kindling, increased the number of spikes during the afterdischarge, which suggests that GH decreases the threshold for seizure generation.<sup>19</sup> In contrast, our study showed an anticonvulsant effect of GH pretreatment. We tested three different amounts of GH, but unexpectedly, only 120 ng had an anticonvulsant effect. Previous studies have reported that daily i.c.v. administration of 120 ng of GH for seven days significantly increased the expansion rate in long-term neurosphere cultures derived from wild-type mice<sup>26</sup> and induces changes in the length of the dendritic trees and the density of dendritic spines in the CA1 region of the dorsal hippocampus and in the PFC (detected 21 days after GH treatment).<sup>5</sup> However, considering that GH can exert its multiple biological activities at different concentrations and that those effects are characterized by a bell-shaped dose-response curve, where the optimal concentration could vary,<sup>27–30</sup> we decided to probe two additional amounts of GH. Our results confirmed that 120 ng per day is the optimal amount for exert an anticonvulsant effect in the SE model and that inferior or superior amounts are ineffective. Differences in the effect produced by GH in both epilepsy models may also be explained by the unlike mechanisms of seizure generation for each one;

whereas the electrical kindling is a model of epileptogenesis (chronic), pilocarpine-induced SE is an acute event. Additionally, the effect of GH pretreatment versus its co-administration with the pro-convulsant stimulus may affect differently the mechanism of seizure generation.

Our results suggest that the subchronic administration of GH decreases the susceptibility of the brain to the neuronal excitability initiated by pilocarpine, but once seizures generate, there is no difference in SE severity (seizure latency, frequency, and duration). Several studies have reported the participation of GH on structural<sup>5</sup> and synaptic plasticity in the hippocampus.<sup>31–34</sup> Specifically, GH enhance excitatory synaptic transmission in the CA1 area of hippocampus, increasing both AMPA and NMDA receptor-mediated excitatory post-synaptic potentials without alteration of GABA-receptor mediated inhibitory synaptic transmission.<sup>31–34</sup> GH also regulates NMDA receptor subunit mRNA expression levels in the rat hippocampus.<sup>35,36</sup> We do not know if five days of GH treatment is enough to promote changes in structural plasticity that may explain its anticonvulsant effect. It is possible that under our experimental conditions GH may have modified excitatory synaptic plasticity through changes in glutamate receptor expression. GH can activate the PI 3 kinase-Akt pathway,<sup>37</sup> which is involved in the expression of glial and neuronal excitatory amino acid transporter.<sup>38,39</sup>

Thus, GH could also increase the expression of glutamate transporters and regulate the primary mechanism for the removal of glutamate from the extracellular space to promote its anticonvulsant effect. Further experiments are needed to understand the anticonvulsant mechanism of GH.

As mentioned above, once the SE was induced, there was no difference between GH or control groups in seizure latency, frequency or duration, suggesting an equivalent extension of neural damage in the hippocampus. Contrary to what was expected, the number of degenerative cells as long as their histological pattern in the hippocampus were unaffected after GH treatments, but similar to formerly reported by others in the lithium–pilocarpine model.<sup>40,41</sup> Hippocampal neuronal cell death induced by lithium–pilocarpine SE can be necrotic and apoptotic, as a result of excessive activation of glutamatergic receptors.<sup>42–44</sup> Previous studies have demonstrated the neuroprotective effect of GH against different insults to the brain, associated with its capability to regulate apoptosis (i.e., an anti-apoptotic effect).<sup>3,11,45–49</sup> In those experiments, GH was administered either before or after the damage inducer and the neuroprotective effect was evaluated at different times, suggesting that this effect is an intrinsic property of this hormone. We initially expected that the i.c.v. administration of GH would stimulate an anti-apoptotic response in the hippocampus that will protect the neurons against the SE insult as we have reported previously.<sup>11</sup> Our results showed the absence of neuroprotection under our experimental conditions, but it is worth noting that the current report is the first study designed to assess the anticonvulsant and neuroprotective effect of GH against neuronal damage caused by SE. Further studies are needed in this field to characterize the exact role of GH in epilepsy and its consequences.

In conclusion, the i.c.v. administration of GH has an anticonvulsant effect since it increases the amount of pilocarpine necessary to induce SE. However, once SE is triggered, the pattern of hippocampal neurodegeneration is not modified, i.e., GH has not a neuroprotective effect. Our findings are relevant in understanding the role of GH in epilepsy.

## Authors' contributions

MLLM, EJA, and IZB conceived and designed the study. IZB and AM performed the research. IZB, MLLM, and LBP analyzed the experimental data. MLLM, IZB, EJA and ISR wrote the paper.

## Sources of support

This work was supported by CONACYT Mexico (fellowship awarded to IZB, 309011).

## Conflict of interests

The authors declare no conflict of interests.

## References

1. Aberg ND, Brywe KG, Isgaard J. Aspects of growth hormone and insulin-like growth factor-I related to neuroprotection, regeneration, and functional plasticity in the adult brain. *Sci World J.* 2006;6:53–80, <http://dx.doi.org/10.1100/tsw.2006.22>.
2. Regalado-Santiago C, López-Meraz ML, Santiago-García J, Fernández-Pomares C, Juárez-Aguilar E. Growth hormone (GH) is a survival rather than a proliferative factor for embryonic striatal neural precursor cells. *Growth Horm IGF Res.* 2013;23:179–86, <http://dx.doi.org/10.1016/j.ghir.2013.07.002>.
3. Scheepens A, Sirimanne ES, Breier BH, Clark RG, Gluckman PD, Williams CE. Growth hormone as a neuronal rescue factor during recovery from CNS injury. *Neuroscience.* 2001;104:677–87, [http://dx.doi.org/10.1016/s0306-4522\(01\)00109-9](http://dx.doi.org/10.1016/s0306-4522(01)00109-9).
4. Arce VM, Devesa P, Devesa J. Role of growth hormone (GH) in the treatment on neural diseases: from neuroprotection to neural repair. *Neurosci Res.* 2013;76:179–86, <http://dx.doi.org/10.1016/j.neures.2013.03.014>.
5. Olivares-Hernández JD, García-García F, Camacho-Abrego I, Flores G, Juárez-Aguilar E. Intracerebroventricular administration of growth hormone induces morphological changes in pyramidal neurons of the hippocampus and prefrontal cortex in adult rats. *Synapse.* 2018;72:1–8, <http://dx.doi.org/10.1002/syn.22030>.
6. Donahue CP, Kosik KS, Shors TJ. Growth hormone is produced within the hippocampus where it responds to age, sex, and stress. *Proc Natl Acad Sci USA.* 2006;103:6031–6, <http://dx.doi.org/10.1073/pnas.0507776103>.
7. Vander Weele CM, Saenz C, Yao J, Correia SS, Goosens KA. Restoration of hippocampal growth hormone reverses stress-induced hippocampal impairment. *Front Behav Neurosci.* 2013;7:66, <http://dx.doi.org/10.3389/fnbeh.2013.00066>.
8. Arellanes-Licea EC, Ávila-Mendoza J, Ramírez-Martínez EC, Ramos E, Uribe-González N, Arámburo C, et al. Upregulation of GH, but not IGF1, in the hippocampus of the lactating dam after kainic acid injury. *Endocr Connect.* 2018;7:258–67, <http://dx.doi.org/10.1530/EC-17-0380>.
9. Ueda Y, Kitamoto A, Willmore LJ, Kojima T. Hippocampal gene expression profiling in a rat model of post-traumatic epilepsy reveals temporal upregulation of lipid metabolism-related genes. *Neurochem Res.* 2013;38:1399–406, <http://dx.doi.org/10.1007/s11064-013-1037-9>.
10. Shin DH, Lee E, Kim JW, Kwon BS, Jung MK, Jee YH, et al. Protective effect of growth hormone on neuronal apoptosis after hypoxia–ischemia in the neonatal rat brain. *Neurosci Lett.* 2004;354:64–8, <http://dx.doi.org/10.1016/j.neulet.2003.09.070>.
11. García-García F, Alberto K, Juárez-Aguilar E, Regalado-Santiago C, Millán-Aldaco D, Blanco-Centurión C, et al. Growth hormone improves hippocampal adult cell survival and counteracts the inhibitory effect of prolonged sleep deprivation on cell proliferation. *Brain Res Bull.* 2011;84:252–7, <http://dx.doi.org/10.1016/j.brainresbull.2011.01.003>.
12. Srimontri P, Hirota H, Kanno H, Okada T, Hirabayashi Y, Kato K. Infusion of growth hormone into the hippocampus induces molecular and behavioral responses in mice. *Exp Brain Res.* 2014;232:2957–66, <http://dx.doi.org/10.1007/s00221-014-3977-y>.
13. Chen JW, Naylor DE, Wasterlain CG. Advances in the pathophysiology of status epilepticus. *Acta Neurol Scand Suppl.* 2007;186:7–15, <http://dx.doi.org/10.1111/j.1600-0404.2007.00803.x>.
14. Fujikawa DG, Itabashi HH, Wu A, Shinmei SS. Status epilepticus–induced neuronal loss in humans without systemic complications or epilepsy. *Epilepsia.* 2000;41:981–91, <http://dx.doi.org/10.1111/j.1528-1157.2000.tb00283.x>.



15. Kubová H, Druga R, Lukasiuk K, Suchomelová L, Haugvicová R, Jirmanová I, et al. Status epilepticus causes necrotic damage in the mediodorsal nucleus of the thalamus in immature rats. *J Neurosci*. 2001;21:3593–9, <http://dx.doi.org/10.1523/JNEUROSCI.21-10-03593.2001>.
16. Fujikawa DG, Shinmei SS, Zhao S, Aviles ER. Caspase-dependent programmed cell death pathways are not activated in generalized seizure-induced neuronal death. *Brain Res*. 2007;1135:206–18, <http://dx.doi.org/10.1016/j.brainres.2006.12.029>.
17. El-Khayat HA, Aly GS, Tomoum HY, Mamdouh RM, Al-Badani AK, Mohamed EI. Growth hormone levels in children and adolescents with epilepsy. *Eur J Pediatr*. 2010;14:508–12, <http://dx.doi.org/10.1016/j.ejpn.2010.02.004>.
18. Lindbom T, Hulting AL, Tomson T. Paradoxical GH response to TRH during status epilepticus in man. *Eur J Neurosci*. 1999;140:307–14, <http://dx.doi.org/10.1530/eje.0.1400307>.
19. Kato K, Suzuki M, Kanno H, Sekino S, Kusakabe K, Okada T, et al. Distinct role of growth hormone on epilepsy progression in a model of temporal lobe epilepsy. *J Neurochem*. 2009;110:509–19, <http://dx.doi.org/10.1111/j.1471-4159.2009.06132.x>.
20. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol*. 2010;8:e1000412, <http://dx.doi.org/10.1371/journal.pbio.1000412>.
21. Paxinos G, Watson C. The rat brain in stereotaxic coordinates: Hard Cover Edition. 2007. USA: Academic Press. <https://www.elsevier.com/books/the-rat-brain-in-stereotaxic-coordinates/paxinos/978-0-12-391949-6>.
22. Racine RJ. Modification of seizure activity by electrical stimulation. II. Motor seizure. *Electroencephalogr Clin Neurophysiol*. 1972;32:295–9, [http://dx.doi.org/10.1016/0013-4694\(72\)90178-2](http://dx.doi.org/10.1016/0013-4694(72)90178-2).
23. Schmued LC, Hopkins KJ. Fluoro-Jade B: a high affinity fluorescent marker for the localization of neuronal degeneration. *Brain Res*. 2000;874:123–30, [http://dx.doi.org/10.1016/S0006-8993\(00\)02513-0](http://dx.doi.org/10.1016/S0006-8993(00)02513-0).
24. Barcia JA, Gallego JM. Intraventricular and intracerebral delivery of anti-epileptic drugs in the kindling model. *Neurotherapeutics*. 2009;6:337–43, <http://dx.doi.org/10.1016/j.nurt.2009.01.015>.
25. Walsler M, Schiöler L, Oscarsson J, Albert MAI, Sevansson J, Aberg ND, et al. Different modes of GH administration influence gene expression in the male rat brain. *J Endocrinol*. 2014;222:181–90, <http://dx.doi.org/10.1530/JOE-14-0223>.
26. Blackmore DG, Reynolds BA, Golmohammadi MG, Large B, Aguilar RM, Haro L, et al. Growth hormone responsive neural precursor cells reside within the adult mammalian brain. *Sci Rep*. 2012;2:250, <http://dx.doi.org/10.1038/srep00250>.
27. Postel-Vinay MC, de Mello Coelho V, Gagnerault MC, Dardenne M. Growth hormone stimulates the proliferation of activated mouse T lymphocytes. *Endocrinology*. 1997;138:1816–20, <http://dx.doi.org/10.1210/endo.138.5.5108>.
28. Zarkesh-Esfahani SH, Kolstad O, Metcalfe RA, Watson PF, von Laue S, Walters S, et al. High-dose growth hormone does not affect proinflammatory cytokine (tumor necrosis factor-alpha, interleukin-6, and interferon-gamma) release from activated peripheral blood mononuclear cells or after minimal to moderate surgical stress. *J Clin Endocrinol Metab*. 2000;85:3383–90, <http://dx.doi.org/10.1210/jcem.85.9.6823>.
29. Waters MJ, Brooks AJ. JAK2 activation by growth hormone and other cytokines. *Biochem J*. 2015;466:1–11, <http://dx.doi.org/10.1042/BJ20141293>.
30. Yang N, Langenheilm JF, Wang X, Jiang J, Chen WY, Frank SJ. Activation of growth hormone receptors by growth hormone and growth hormone antagonist dimers: insights into receptor triggering. *Mol Endocrinol*. 2008;22:978–88, <http://dx.doi.org/10.1210/me.2007-0424>.
31. Mahmoud GS, Grover LM. Growth hormone enhances excitatory synaptic transmission in area CA1 of rat hippocampus. *J Neurophysiol*. 2006;95:2962–74, <http://dx.doi.org/10.1152/jn.00947.2005>.
32. Zearfoss NR, Alarcon JM, Trifilieff P, Kandel E, Richter JD. A molecular circuit composed of CPEB-1 and c-Jun controls growth hormone-mediated synaptic plasticity in the mouse hippocampus. *J Neurosci*. 2008;28:8502–9, <http://dx.doi.org/10.1523/JNEUROSCI.1756-08.2008>.
33. Molina DP, Ariwodola OJ, Linville C, Sonntag WE, Weiner JL, Brunso-Bechtold JK, et al. Growth hormone modulates hippocampal excitatory synaptic transmission and plasticity in old rats. *Neurobiol Aging*. 2012;33:1938–49, <http://dx.doi.org/10.1016/j.neurobiolaging.2011.09.014>.
34. Molina DP, Ariwodola OJ, Weiner JL, Brunso-Bechtold JK, Adams MM. Growth hormone and insulin-like growth factor-I alter hippocampal excitatory synaptic transmission in young and old rats. *Age (Dordr)*. 2013;35:1575–87, <http://dx.doi.org/10.1007/s11357-012-9460-4>.
35. Le Grevès M, Steensland P, Le Grevès P, Nyberg F. Growth hormone induces age-dependent alteration in the expression of hippocampal growth hormone receptor and N-methyl-D-aspartate receptor subunits gene transcripts in male rats. *Proc Natl Acad Sci USA*. 2002;99:7119–23, <http://dx.doi.org/10.1073/pnas.092135399>.
36. Le Grevès M, Zhou Q, Berg M, Le Grevès P, Fhølenhag K, Meyerson B, Nyberg F. Growth hormone replacement in hypophysectomized rats affects spatial performance and hippocampal levels of NMDA receptor subunit and PSD-95 gene transcript levels. *Exp Brain Res*. 2006;173:267–73, <http://dx.doi.org/10.1007/s00221-006-0438-2>.
37. Dehkhoda F, Lee CMM, Medina J, Brooks AJ. The growth hormone receptor: mechanism of receptor activation cell signaling, and physiological aspects. *Front Endocrinol (Lausanne)*. 2018;9:35, <http://dx.doi.org/10.3389/fendo.2018.00035>.
38. Sims KD, Straff DJ, Robinson MB. Platelet-derived growth factor rapidly increases activity and cell surface expression of the EAAC1 subtype of glutamate transporter through activation of phosphatidylinositol 3-kinase. *J Biol Chem*. 2000;275:5228–37, <http://dx.doi.org/10.1074/jbc.275.7.5228>.
39. Krizman-Genda E, González MI, Zelenia O, Robinson MB. Evidence that Akt mediates platelet-derived growth factor-dependent increases in activity and surface expression of the neuronal glutamate transporter. EAAC1. *Neuropharmacology*. 2005;49:872–82, <http://dx.doi.org/10.1016/j.neuropharm.2005.07.014>.
40. Curia G, Longo D, Biagini G, Jones RS, Avoli M. The pilocarpine model of temporal lobe epilepsy. *J Neurosci Methods*. 2008;172:143–57, <http://dx.doi.org/10.1016/j.jneumeth.2008.04.019>.
41. Scorza FA, Arida RM, Naffah-Mazzacoratti M, da G, Scerni DA, Calderazzo L, Cavalheiro EA. The pilocarpine model of epilepsy: what have we learned? *An Acad Bras Cienc*. 2009;81:345–65, <http://dx.doi.org/10.1590/s0001-37652009000300003>.
42. Fujikawa DG, Shinmei SS, Cai B. Lithium–pilocarpine-induced status epilepticus produces necrotic neurons with internucleosomal DNA fragmentation in adult rats. *Eur J Neurosci*. 1999;11:1605–14, <http://dx.doi.org/10.1046/j.1460-9568.1999.00573.x>.
43. Wang L, Liu YH, Huang YG, Chen LW. Time-course of neuronal death in the mouse pilocarpine model of chronic epilepsy using Fluoro-Jade C staining. *Brain Res*. 2008;1241:157–67, <http://dx.doi.org/10.1016/j.brainres.2008.07.097>.
44. Lopez-Meraz ML, Wasterlain CG, Rocha LL, Allen S, Niquet J. Vulnerability of postnatal hippocampal neurons to seizures



- varies regionally with their maturational stage. *Neurobiol Dis.* 2010;37:394–402, <http://dx.doi.org/10.1016/j.nbd.2009.10.019>.
45. Isla A, Budke M, Cacicedo L, García-Grande A, Vázquez-Rodríguez I, de Miguel E, et al. Protective effects of growth hormone en cell cultures of the central nervous system. *Rev Neurol.* 2002;34:208–11. PMID: 12022066.
46. Svensson AL, Bucht N, Hallberg M, Nyberg F. Reversal of opiate-induced apoptosis by human recombinant growth hormone in murine foetus primary hippocampal neuronal cell cultures. *Proc Natl Acad Sci USA.* 2008;105:7304–8, <http://dx.doi.org/10.1073/pnas.0802531105>.
47. Alba-Betancourt C, Luna-Acosta JL, Ramírez-Martínez CE, Ávila-González D, Granados-Ávalos E, Carranza M, et al. Neuro-protective effects of growth hormone (GH) after hypoxia–ischemia injury in embryonic chicken cerebellum. *Gen Comp Endocrinol.* 2013;183:17–31, <http://dx.doi.org/10.1016/j.yggen.2012.12.004>.
48. Luna-Acosta JL, Alba-Betancourt C, Martínez-Moreno CG, Ramírez C, Carranza M, Luna M, et al. Direct antiapoptotic effects of growth hormone are mediated by PI3K/Akt pathway in the chicken bursa of Fabricius. *Gen Comp Endocrinol.* 2015;224:148–59, <http://dx.doi.org/10.1016/j.yggen.2015.07.010>.
49. Nylander E, Gronbladh A, Zelleroth S, Diwakarla S, Nyberg F, Hallberg M. Growth hormone is protective against acute methadone-induced toxicity by modulating the NMDA receptor complex. *Neuroscience.* 2016;339:538–47, <http://dx.doi.org/10.1016/j.neuroscience.2016.10.019>.