



ORIGINAL ARTICLE

Cognitive rehabilitation program in patients with multiple sclerosis: A pilot study



R.M. Jiménez-Morales^{a,b,*}, Y. Broche-Pérez^c, Y. Macías-Delgado^d, C. Sebrango^e, S. Díaz-Díaz^a, R. Castiñeira-Rodríguez^a, F.J. Pérez-González^b, C. Forn^f

^a Physical Medicine and Rehabilitation Department, Rehabilitation Hospital Dr. Faustino Pérez Hernández, Sancti Spiritus, Cuba

^b Center for Studies in Educational Sciences, José Martí University of Sancti Spiritus, Cuba

^c Psychology Department, Universidad Central "Marta Abreu" de Las Villas, Cuba

^d Psychology Department, University of Medical Sciences Dr. Fustino Pérez Hernández, de Sancti Spiritus, Cuba

^e Center for Studies Energy and Industrial, José Martí University of Sancti Spiritus, Cuba

^f Psicología Básica, Clínica i Psicobiología Department, Universitat Jaume I, Spain

Received 14 December 2020; accepted 24 March 2021

Available online 10 July 2021

KEYWORDS

Aerobic exercise;
Multiple sclerosis;
Cognitive function;
Cognitive
rehabilitation;
Cognitive reserve

Abstract

Introduction: In recent years, there has been an increase of studies dedicated to cognitive rehabilitation in patients with multiple sclerosis (MS); however, few of these analyze the impact on such variables as cognitive reserve. The study aims to explore the effects of a cognitive rehabilitation program comprising a combination of cognitive and physical exercises, as well as group sessions to improve cognitive performance, emotional state, and cognitive reserve index. **Method:** Fifty patients with MS were subdivided into 2 groups: the control group, which performed aerobic exercise ($n=25$), and the experimental group ($n=25$), which participated in the integrated cognitive rehabilitation program (ICRP). All participants were evaluated 3 times (baseline, post-treatment, and long-term) with the Brief Repeatable Battery of Neuropsychological Tests, Cognitive Reserve Scale, Beck Depression Inventory, and a scale evaluating trait and state anxiety.

Results: Compared with the control group, patients in the experimental group showed improvements in cognitive function, with significant changes in measures of information processing speed, attention, memory, cognitive reserve index, and long-term mood.

Conclusions: The ICRP was effective in improving cognitive and emotional function in MS, and increased the cognitive reserve index.

© 2021 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: mauricio770927@gmail.com (R.M. Jiménez-Morales).

PALABRAS CLAVE

Ejercicio aeróbico;
Esclerosis múltiple;
Función cognitiva;
Rehabilitación
cognitiva;
Reserva cognitiva

Programa de rehabilitación cognitiva para pacientes con esclerosis múltiple: estudio piloto**Resumen**

Introducción: En los últimos años se ha observado un interés creciente por la rehabilitación cognitiva en pacientes con esclerosis múltiple. Sin embargo, pocos estudios han analizado su impacto en variables como la reserva cognitiva. Analizamos el efecto de un programa de rehabilitación cognitiva que incluye ejercicios físicos y cognitivos, así como sesiones en grupo enfocadas a mejorar el rendimiento cognitivo, el estado emocional y el índice de reserva cognitiva.

Métodos: Nuestro estudio incluyó a 50 pacientes con esclerosis múltiple, divididos en 2 grupos: un grupo control (n = 25), en el que los pacientes realizaban ejercicio aeróbico, y un grupo experimental (n = 25), al que se administró un programa integral de rehabilitación cognitiva. Evaluamos a todos los pacientes en 3 momentos diferentes (al inicio, tras el tratamiento, y a largo plazo) con la Batería Neuropsicológica Breve, la Escala de Reserva Cognitiva, el Inventario de Depresión de Beck y una escala para medir la ansiedad rasgo y la ansiedad estado.

Resultados: Los pacientes del grupo experimental mostraron un mejor rendimiento cognitivo que los controles, con cambios significativos en medidas de velocidad de procesamiento de la información, atención, memoria, índice de reserva cognitiva y estado de ánimo a largo plazo.

Conclusión: Nuestros resultados demuestran la eficacia del programa de rehabilitación cognitiva para mejorar las funciones cognitivas y emocionales de los pacientes con esclerosis múltiple y aumentar el índice de reserva cognitiva.

© 2021 Sociedad Española de Neurología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Multiple Sclerosis (MS) is characterized, among other aspects, by the presence of cognitive deficits that affect approximately 65% of patients diagnosed with the disease.^{1,2} Among the most frequent cognitive disorders in patients with MS are, the decreasing in processing speed, attention deficits, memory disorders and executive dysfunctions.^{1,3} In patients with MS, these cognitive deficiencies can affect the performance of various daily activities.

Recently, several studies aimed at evaluating the efficacy of cognitive rehabilitation in MS patients. Interventions have included actions to improve attention, learning, memory and executive functions,^{4–7} while other researchers have used multimodal intervention strategies that combine cognitive stimulation and psychotherapeutic actions.⁸

However, the effectiveness of cognitive rehabilitation strategies in patients with MS is contradictory. This is due, among other factors, to the heterogeneity in the groups studied, the diversity of rehabilitation programs implemented and the measures used to assess outcomes.^{9,10}

Currently, rehabilitation studies in MS patients focus on verifying the effectiveness of multidisciplinary intervention programs, understanding the concept of neurorehabilitation from a broad perspective. This field includes cognitive rehabilitation programs that use a combination of different strategies, for example, the use of physical and cognitive exercises,^{11,12} group sessions to learn how to use compensatory strategies and home computer training,⁷ psychoeducation, self-regulation and compensatory training,¹³ and enrichment programs for promoting different cognitive leisure activities.^{14,15}

Programs focused on the enrichment of leisure activities are usually developed around the cognitive reserve construct. Cognitive Reserve (CR) The concept of CR refers to differences in cognitive processes that explain differential susceptibility to functional impairment in the presence of pathology. CR can therefore be considered an active model of reserve where the brain actively attempts to cope with brain damage by using pre-existing cognitive processes or by enlisting compensatory processes.¹⁶

From this perspective, it has been observed that MS patients with a higher level of cognitive reserve, compared to patients with a lower reserve, take longer to show cognitive decline.^{17,18} High levels of CR were associated with better cognitive task performance in verbal and spatial memory, attention, processing speed, verbal fluency, and inhibitory control. The results indirectly emphasize the value of early cognitive assessment of cognitive status and CR levels to enable timely initiation of cognitive interventions to increase cognitive reserve in MS in patients with low levels of CR.¹⁹

Currently, some authors conceive CR as a dynamic concept that can be modified and enriched throughout a lifespan. For this reason, CR construct has been incorporated from a multidisciplinary approach into rehabilitation programs.^{20–21}

However, there are still few studies of neurorehabilitation aimed at building cognitive reserve in patients with MS. In the present study, a cognitive rehabilitation program called the *Integrated Program of Cognitive Rehabilitation* (PIRICO) is proposed (PIRICO is the acronym for the name of the program in Spanish: *Programa Integrado de Rehabilitación Cognitiva*).

The objective of this study was to explore the effect of a multimodal program, in which cognitive exercises were combined with physical exercises; as well as group sessions aimed at improving cognitive performance, reserve-building activities and emotional states.

Methods

Participants

A total of 50 patients with Multiple Sclerosis (Relapsing-Remitting phenotype) who regularly attend to the Hospital de Rehabilitación “Dr. Faustino Pérez Hernández”, in Sancti Spiritus province, Cuba were evaluated. The patients had clinically defined MS according to the McDonald’s criteria²² and the criteria of the clinical forms division proposed by Lubling and Reingold.²³

All the patients were assessed at the neurological level, using the Expanded Disability Status Scale (EDSS). Additionally, the following inclusion criteria was considered: age between 18 years and 65 years, duration of the disease ≤ 20 years (ability to understand and comply with the cognitive rehabilitation program) and EDSS score ≤ 6 . Patients with a severe psychiatric disorder, acute relapses (outbreaks one month before the evaluation or during the investigation that are treated with steroids or immunosuppressive drugs and due to their functional impact limit participation in the study), and other neurological disorders that could affect cognition, were excluded.

A simple recruitment process was carried out where participants were randomly assigned (in a ratio of 1:1) into an experimental group (EG: $n=21$) and a control group (CG: $n=20$).

The reasons for exclusion of some patients in the research process are described in Fig. 1. The study was approved by the ethics committee of the Hospital Provincial de Rehabilitación. All patients were informed about the importance of the research and signed the informed consent in accordance with the ethical principles of the Helsinki declaration.

Neuropsychological assessment

The brief repeatable battery

The EG and CG participants completed the Brief Repeatable Battery, translated and adapted to the Spanish-speaking population by Sepulcre et al.,²⁴ The Brief Repeatable Battery (BRB) includes the Selective Reminding Test (SRT) (verbal long-term learning and memory), the Spatial Recall Test (SPART 10/36) (visuospatial long-term learning and memory), the Symbol Digit Modalities Test (SDMT) (attention and speed of information processing), the Paced Auditory Serial Addition Test (PASAT – 2 and 3) (speed of information processing and working memory), and the Word List Generation (WLG) (phonetic and semantic verbal fluency). The patients had never been evaluated using the Brief Repeatable Battery before. The results were considered by the individual measures of the BRB tests.

Cognitive reserve scale

This scale is made up of 25 items divided into four types of activities: basic daily activities, training and information activities, hobbies and activities of social life.^{25,26} The scale explores three periods of life using a five-option response scale (Likert) (0, never; 1, once or several times a year; 2, once or several times a month; 3, once or several times a week; 4, three times or more in the week). The scale score range is 0–96.

Emotional status

The emotional state of the patients was explored using the Spanish version of the Beck Depression Inventory (BDI)^{27,28} and the State-Trait Anxiety Inventory (STAI).²⁹

Integrated cognitive rehabilitation program (PIRCO)

General considerations

The PIRCO program was implemented in patients who conform the experimental group. The cognitive rehabilitation program is theoretically based on an individualized approach that is built on the basis of the individual’s strengths and works to compensate for deficit areas in order to increase the person’s ability to participate more fully in daily life activities.³⁰

The program implies three fundamental axes of intervention offered by a multidisciplinary team, where aerobics training, cognitive training, and group sessions from a cognitive and ecological perspective were combined. The training (physical and cognitive) involved the repeated practice of specific tasks designed to reflect the underlying cognitive processes. The main ingredient is repetitive work and explicit teaching of cognitive tasks.³⁰ Group sessions offer activities aimed at self-knowledge work, compensation to improve memory and executive functions, stress management, and promotion of cognitive leisure activities.

The program was implemented during six weeks, with a daily frequency. In the morning session, the treadmill aerobic exercises were performed on daily basis, followed by Dynamic board game of cubes and signs (TaDiCS®). In the afternoon session the ergometric bicycle was performed daily. After the aerobic exercises, the Modified PASAT tasks training (Monday, Tuesday and Friday) and group sessions (Tuesday and Thursday) were alternated. The CG patients practiced only the aerobic exercises, with the same duration, frequency and intensity as the EG (morning session, the treadmill aerobic exercises, and the ergometric bicycle in afternoon session). These patients did not receive any form of cognitive training.

Structure of individual sessions

Aerobics training

A graduated resistance program was implemented, by combining training on the WNQ-7000^a treadmill and the ergometric bicycle (ERGOCIT-AT), using as a reference the training protocol developed by Sandroff et al.,³¹ The dosage in both workouts was gradual, reaching a maximum time of




Figure 1 Flowchart of intervention.

30 min in the sixth week. Intensity (heart rate reserve) and exercise duration (time in minutes) were controlled. The increasing in intensity and duration of the exercise was carried out based on the physical conditions and the stability of the heart pulse of each patient. If the patients presented a greater motor impairment that limited the initial phase of training (EDSS score 5–6), the dose was adjusted according to their rehabilitation needs, and they were accompanied at all times to control gait dynamics (treadmill training). All sessions were conducted by licensed physical therapists.

Cognitive training

Dynamic board game of cubes and signs (TaDiCS®)

The TaDiCS® was used with the aim of training attention and the solution of practical-constructive problems. Its design is based on the clinical model of attention developed by Sohlberg and Mateer,³² as well as the problem solving model proposed by Luria.³³ The cognitive training was carried out from Monday to Friday, in the morning session, during 45 min. Tasks of visual tracking and visuomotor speed

(visual modality), sustained and selective attention (auditory and visual modality), inhibitory control (auditory and visual modality) and practical-constructive problem solving were used.¹¹ These activities were always performed after the aerobic exercises practiced in the morning.

The modified PASAT (Paced Auditory Serial Addition Task) and PVSAT (Paced Visual Serial Addition Task) tasks

The Modified PASAT tasks were implemented through the Computerized Cognitive Rehabilitation and Management System (GERCO®). The speed of information processing, sustained attention and working memory were trained.

Firstly, auditory tasks and then visual tasks were trained. The cognitive tasks had different indicators of complexity: type of tasks (auditory and visual), number of digits, complexity of the digits, duration of the stimulus, interval between stimuli and distractors. This training was carried out three times a week (Monday, Wednesday and Friday) in the afternoon session after the aerobic exercises and with a duration of 30 min. The modified PASAT and PVSAT tasks have shown effectiveness in the cognitive rehabilitation of MS patients.^{34,35}

In general, the cognitive training of attention tasks, problem solving, speed of information processing and working memory was used, because they are functions that are usually affected early in MS.

Group work sessions

A total of 10 group sessions were held with two weekly frequencies (Tuesday and Thursday). In each session, 60 min were dedicated to group work with the patients and 30 min to inform their caregivers on the topics discussed. Caregivers participated with prior informed consent. The topics of the sessions were the following: (1) how to improve cognitive performance through daily life activities; (2) what is cognitive reserve and how to enrich it; (3) how to improve learning and memory, (4) promotion of cognitive leisure activities and self-generation; (5) how to improve attention/concentration capacity; (6) successful behavior planning and self-regulation; (7) coping with stress and (8) development of leisure activities and cognitive health.

These group activities have been designed *ad-hoc* by program researchers with a psychoeducational and compensatory approach in the field of cognitive rehabilitation. The objective of the sessions with the caregivers was to provide information on the same content that was discussed with the patients; as well as to inform about the compensation strategies used during the work session. All sessions were coordinated by an experienced therapist and co-therapist.

Implementation of the intervention program: General procedure

During the first week of treatment, a one-hour briefing session was held. During the session, the characteristics of

the intervention program were explained to the patients and caregivers of the EG and CG (number and distribution of sessions, type of activities, and evaluation periods). In the case of the EG participants, the characteristics of the multimodal cognitive rehabilitation activities were presented individually and in groups. Both groups (EG and CG) were evaluated three times: baseline, post-intervention (6 weeks), and long-term follow-up (6 months). For the evaluations, the same brief battery of cognitive tests was used, but with different versions (version A, B and A) to control the effect of learning. It should be noted that the patients that participated in the program had not previously carried out cognitive assessments that included PASAT or other cognitive tests.

The post-intervention assessment was performed at the end of the sixth week and the follow-up stage (long-term assessment) was performed 30 weeks later (6 months after completing the intervention). The research process was developed in a double-blind manner. Both PIRCO participants, as well as the evaluators and analysts remained blind to the other activities of the program.

Analysis of data

The data were processed using SPSS/Windows, version 21. Descriptive statistics were used to explore the demographic and clinical characteristics of the participants. The Chi-square test was used to explore the distribution between genders. Before selecting the *t*-test or ANOVA method, we explored the distribution of the data and checked that the homogeneity of variances was not violated (Levene's test). An independent *t*-test sample was performed to compare demographic and clinical variables between groups. A repeated measures ANOVA (2 (GE and GC) \times 3 (baseline assessment (T1), post-intervention (T2) and long-term evaluation (T3)) was performed to check the effect of the program in EG patients in comparison to CG patients. A two-way ANOVA with repeated measures on one factor was conducted. The Mauchly's test of sphericity was used to assess whether or not the assumption of sphericity was met. The Greenhouse–Geisser effect correction was employed when sphericity was violated.³⁶ The effect sizes were calculated using omega-squared (ω^2). The reference values for omega-squared were .01, .06, .14 (small, medium and large effect size respectively).³⁷ An additional independent-samples *t*-test was conducted to compare the differences in the scores recorded at the baseline and time 3 (*change score*). Effect sizes were calculated using Cohen's *d*. Cohen classifies .2 as a small effect, .5 as a medium effect and .8 as a large effect.³⁸ Values of $p < .05$ were considered significant.

Results

Characteristics of the sample

The demographic and clinical characteristics of the study groups are shown in Table 1. As shown in Table 1, no significant differences were observed in the different variables

Table 1 Demographical and clinical characteristics of the experimental and control group.

Characteristic	Experimental group (<i>n</i> = 21)	Control group (<i>n</i> = 20)	<i>t</i>	<i>p</i> Value
<i>Age_{years} Mean (SD)</i>	46.00 (10.363)	41.75 (10.968)	1.276	.210
20–30 (%)	2 (9.5)	4 (20)		
31–40 (%)	4 (19)	3 (15)		
41–50 (%)	8 (38.1)	10 (50)		
51–65 (%)	7 (33.3)	3 (15)		
<i>Sex (%)</i>				
Female	17 (81)	11 (55)		
Male	4 (19)	9 (45)	χ^2 : 3.186	.100
<i>Education_{years} Mean (SD)</i>	15.81 (3.907)	14.15 (3.731)	1.390	.173
9th grade (%)	—	—		
12th grade (%)	2 (9.5)	5 (25)		
Middle Technician (%)	10 (47.6)	9 (45)		
Academic (%)	9 (42.9)	6 (30)		
<i>Disease duration_{years} Mean (SD)</i>	9.71 (5.217)	7.15 (4.837)	1.633	.111
<i>Median EDSS (SD)</i>	4.57 (0.939)	4.37 (0.809)	0.716	.478

SD: standard deviation; EDSS: Expanded Disability Status Scale.

Table 2 Cognitive tests and emotional states in the experimental and control group.

Measure	EG Mean \pm SD	CG Mean \pm SD	<i>p</i> Value
SRT-S	41.43 \pm 8.44	40.50 \pm 10.26	.753
SRT-R	27.76 \pm 8.99	30.45 \pm 12.28	.427
SRT-D	6.00 \pm 1.92	6.90 \pm 1.83	.133
SPART 10/36	11.71 \pm 5.11	11.75 \pm 6.25	.984
SPART 10/36 D	4.67 \pm 1.74	4.50 \pm 1.79	.764
SDMT	24.67 \pm 13.17	28.85 \pm 13.60	.323
PASAT 3 seg.	30.29 \pm 8.93	29.20 \pm 12.43	.749
PASAT 2 seg.	18.52 \pm 11.31	22.70 \pm 8.39	.189
WLG Phonetic	14.52 \pm 7.05	17.50 \pm 6.68	.180
WLG Semantic	20.81 \pm 8.61	22.95 \pm 6.24	.370
STAI – State	34.67 \pm 7.61	39.75 \pm 12.17	.115
STAI – trait	35.67 \pm 8.55	41.15 \pm 9.80	.063
BDI (total score)	14.71 \pm 5.51	13.00 \pm 6.62	.372

SD: Standard deviation; SRT-S: Selective Reminding Test (storage), SRT-R: Selective Reminding Test (recovery); SRT-D: Selective Reminding Test (Delayed Recall); SPART 10/36: Spatial Recall Test; SPART 10/36D: Spatial Recall Test (Delayed Recall); SDMT: Symbol Digit Modalities Test; PASAT 2–3: Paced Auditory Serial Addition Task 2 and 3 segundos; WLG: Word List Generation; STAI: State-Trait Anxiety Inventory; BDI: The Beck Depression Inventory.

explored. An equivalent cognitive and emotional profile was observed between EG and CG before starting the cognitive rehabilitation program (Table 2).

Cognitive functioning

The ANOVA results showed significant changes over time in nine of the ten neuropsychological tests (Table 3). Compared to the baseline (T1), a significant difference was observed in the experimental group according to the scores obtained in T3. Significant differences were observed in verbal memory (SRT) and visuospatial memory (SPART), processing speed (SDMT), attention and working

memory (PASAT-3), and verbal fluency (WLG). No significant differences were observed in the control group.

On the other hand, Time \times Group interactions showed significant differences. The effect of the iterations was significant in the measures of verbal memory (SRT) and visuospatial (SPART), speed of processing (SDMT), attention and working memory (PASAT-3). The analysis of the calculation of effect on changes in cognitive performance (effect \times time), showed a small effect in all the cognitive variables studied ($\omega^2 < 0.06$) except for SPART 10/36 where the effect size was medium ($\omega^2 = 0.06–0.14$) (Table 3). In the case of time \times group interactions showed a small effect in all the cognitive variables studied. For SRT-S values the effect sizes were medium.

Table 3 Evaluation of cognitive functions by repeated measures analysis (ANOVA) between the experimental group (EG) and the control group (CG).

Neuropsychological test	Group	T1	T2	T3	F (df)				F (df)			
	Baseline	Mean ± SD	Mean ± SD	time effect	Group by time effect	p	ω ²	1 − β	F	p	ω ²	1 − β
SRT-S	EG	41.43 ± 8.43	45.10 ± 6.83	42.95 ± 7.95	4.97	.012	.01	.80	6.47	.004	.13	.93
	CG	40.50 ± 10.25	40.05 ± 11.46	39.50 ± 10.13								
SRT-R	EG	27.76 ± 8.99	31.29 ± 8.80	29.95 ± 9.41	5.44	.008	.03	.66	10.54	<.0001	.01	.86
	CG	30.45 ± 12.28	30.10 ± 12.76	29.85 ± 11.90								
SRT-D	EG	6.00 ± 1.92	7.05 ± 1.43	7.29 ± 1.61	8.49	.001	.04	.99	4.22	.022	.02	.83
	CG	6.90 ± 1.83	7.30 ± 1.13	7.05 ± 1.05								
SPART 10/36	EG	11.71 ± 5.11	15.81 ± 4.99	14.19 ± 4.11	14.36	<.0001	.08	.98	10.02	<.0001	.02	.92
	CG	11.75 ± 6.24	12.30 ± 5.54	11.45 ± 4.65								
SPART 10/36 D	EG	4.67 ± 1.74	6.33 ± 1.15	5.67 ± 1.46	16.11	<.0001	.02	.99	4.91	.013	.03	.87
	CG	4.50 ± 1.792	5.05 ± 1.23	4.40 ± 1.35								
SDMT	EG	24.67 ± 13.17	32.38 ± 10.95	30.71 ± 10.24	27.71	<.0001	.01	.99	15.61	<.0001	.01	.99
	CG	28.85 ± 13.59	29.95 ± 12.33	29.45 ± 10.94								
PASAT 3 seg.	EG	30.29 ± 8.93	37.48 ± 9.57	35.67 ± 9.04	23.42	<.0001	.01	.99	17.32	<.0001	.01	.99
	CG	29.20 ± 12.43	30.50 ± 11.65	28.35 ± 10.58								
PASAT 2 seg.	EG	18.52 ± 11.31	20.05 ± 8.63	19.62 ± 8.98	1.77	.18	0	.98	0.61	.55	.01	.96
	CG	22.70 ± 8.38	23.45 ± 7.49	22.55 ± 8.19								
WLG Phonetic	EG	14.57 ± 7.04	15.81 ± 5.47	15.90 ± 5.95	3.28	.049	0	.51	2.48	.10	.01	.61
	CG	17.50 ± 6.67	18.05 ± 6.13	16.85 ± 6.11								
WLG Semantic	EG	20.81 ± 8.61	25.10 ± 5.73	22.62 ± 6.53	6.97	.003	.01	.63	1.84	.17	.01	.76
	CG	22.95 ± 6.24	21.45 ± 5.83	21.90 ± 6.39								

SD: Standard deviation; *F*-tests: Wilks' lambda statistic; SRT-S: Selective Reminding Test (storage), SRT-R: Selective Reminding Test (recovery); SRT-D: Selective Reminding Test (Delayed Recall); SPART 10/36: Spatial Recall Test; SPART 10/36D: Spatial Recall Test (Delayed Recall); SDMT: Symbol Digit Modalities Test; PASAT 2–3: Paced Auditory Serial Addition Task 2 and 3 s; WLG: Word List Generation; ω^2 : omega-squared; 1 – β : observed power.

Table 4 Evaluation of the cognitive reserve index, cognitively stimulating activities and emotional states through repeated measures between the experimental group (EG) and the control group (CG).

Measure	Groups	T1	T2	T3	F(df)			F (df)				
		Baseline	Mean ± SD	Mean ± SD	time effect	Group by time effect	p	ω ²	1 − β	F	p	ω ²
Basic daily activities	EG	8.67 ± 1.93	9.81 ± 2.44	9.24 ± 1.81	9.05	.001	.02	.94	2.49	.096	.01	.55
	CG	9.25 ± 1.92	9.55 ± 2.48	9.70 ± 2.13								
Training and information activities	EG	8.62 ± 2.04	9.19 ± 1.81	9.95 ± 2.04	7.38	.002	.02	.68	4.50	.016	.04	.91
	CG	9.05 ± 1.23	9.70 ± 1.59	8.85 ± 1.72								
Hobbies	EG	26.14 ± 6.50	31.29 ± 7.58	32.05 ± 4.64	9.70	<.0001	.28	.96	3.82	.031	.03	.66
	CG	26.55 ± 3.87	28.65 ± 5.67	27.30 ± 5.09								
Activities of social life	EG	5.10 ± 2.07	8.38 ± 1.66	7.76 ± 2.03	19.37	<.0001	.07	.99	1.08	.351	.01	.18
	CG	5.80 ± 2.46	8.55 ± 2.06	7.40 ± 1.73								
Cognitive reserve index	EG	48.52 ± 7.24	58.67 ± 7.91	59.00 ± 5.86	26.67	<.0001	.13	.99	7.05	.002	.06	.96
	CG	50.65 ± 5.06	56.45 ± 7.70	53.25 ± 6.34								
STAI — State	EG	34.67 ± 7.61	32.52 ± 7.17	31.14 ± 4.85	4.91	.013	.01	.82	3.25	.050	.01	.63
	CG	39.75 ± 12.17	37.60 ± 8.83	39.50 ± 8.75								
STAI — trait	EG	35.67 ± 8.55	32.76 ± 7.13	33.33 ± 6.20	9.32	.001	.01	.94	1.99	0.15	0	.45
	CG	41.15 ± 9.80	39.85 ± 8.39	41.05 ± 8.35								
BDI (total score)	EG	14.71 ± 5.50	12.14 ± 3.95	11.71 ± 3.85	14.85	<.0001	.01	.65	7.17	.002	.02	.92
	CG	13.00 ± 6.62	12.45 ± 6.04	13.55 ± 5.09								

SD: Standard deviation; F-tests: Wilks' lambda statistic; STAI: State-Trait Anxiety Inventory; BDI: The Beck Depression Inventory; ω^2 : omega-squared; 1 – β : observed power.

Table 5 Change score (baseline to time 3) in cognitive tests, cognitive reserve and emotional states in both groups.

Measure	EG Mean \pm SD	CG Mean \pm SD	<i>p</i> -Value	<i>d</i>
SRT-S	1.52 \pm 3.06	−1.00 \pm 3.91	.026	.72
SRT-R	2.19 \pm 3.59	−0.60 \pm 1.76	.003	.98
SRT-D	1.29 \pm 1.19	0.15 \pm 1.31	.006	.91
SPART 10/36	2.48 \pm 2.71	−0.30 \pm 3.28	.005	.93
SPART 10/36 D	1.00 \pm 1.34	−0.10 \pm 1.02	.005	.92
SDMT	6.05 \pm 5.91	0.60 \pm 4.20	.002	1.06
PASAT 3 seg.	5.38 \pm 5.14	−0.85 \pm 3.47	<.0001	1.41
PASAT 2 seg.	1.09 \pm 3.97	−0.15 \pm 3.17	.27	.34
WLG Phonetic	1.33 \pm 3.01	−0.65 \pm 2.06	.019	.76
WLG Semantic	1.81 \pm 4.28	−1.05 \pm 2.82	.016	.79
Basic daily activities	0.57 \pm 1.08	0.45 \pm 1.05	.717	.11
Training and information activities	1.33 \pm 1.35	−0.20 \pm 1.88	.005	.94
Hobbies	5.90 \pm 6.66	0.75 \pm 5.23	.009	.86
Activities of social life	2.67 \pm 2.46	1.60 \pm 2.60	.185	.42
Cognitive reserve index	10.48 \pm 6.48	2.60 \pm 6.89	.001	1.18
STAI – State	−3.52 \pm 4.51	−0.25 \pm 5.35	.040	.66
STAI – trait	−2.33 \pm 4.53	−0.10 \pm 2.94	.07	.20
BDI (total score)	−3.00 \pm 3.50	0.55 \pm 2.95	.001	1.09

SD: Standard deviation; *d*: Effect sizes were calculated using Cohen's *d*: SRT-S: Selective Reminding Test (storage), SRT-R: Selective Reminding Test (recovery); SRT-D: Selective Reminding Test (Delayed Recall); SPART 10/36: Spatial Recall Test; SPART 10/36D: Spatial Recall Test (Delayed Recall); SDMT: Symbol Digit Modalities Test; PASAT 2–3: Paced Auditory Serial Addition Task 2 and 3 s; WLG: Word List Generation; STAI: State-Trait Anxiety Inventory; BDI: The Beck Depression Inventory.

Cognitive reserve

The analysis of the effect \times time interactions (Table 4), revealed significant changes in the cognitive reserve index. It was found that the training and information activities, hobbies, and social life showed significant changes after the 6-month follow-up. The time \times group interaction was also observed. A significant effect in the cognitive reserve index was verified; as well as self-report measures of basic daily activities, activities that involved training and information, and hobbies. The effect size of the effect \times time interactions was small ($\omega^2 < 0.06$) except for cognitive reserve index where the effect size was medium ($\omega^2 = 0.13$) and hobbies where the effect size was large ($\omega^2 = 0.28$). In the time \times group interaction, analysis showed a small effect in all the reserve variables studied ($\omega^2 < 0.06$).

After 6 months, significant changes were also found in the effect \times time interactions in the levels of trait and state anxiety, as well as in depression in both groups (Table 4). In the time \times group interaction analysis, it was found that most of the emotional variables changed significantly, except for anxiety levels as a trait. In the time \times group interaction, analysis revealed a small effect in all the emotional variables studied ($\omega^2 < 0.06$).

Analysis of change score from T1 to T3

The data showed an increase in the performance of all neuropsychological tests in the EG, indicating values (+1); unlike the CG where a decline was observed in the execution of most of these tests (−0) (Table 5). The difference between the two groups was significant. In a special way, in the SRT-R tests ($p = .003$; $d = .98$); SRT-D ($p = .006$; $d = .91$); SPART

10/36 ($p = .005$; $d = .93$); SPART 10/36D ($p = .005$; $d = .92$); SDMT ($p = .002$; $d = 1.06$); PASAT-3 ($p < .001$; $d = 1.41$).

Training and information activities ($p = .005$; $d = .94$), hobbies – hobbies ($p = .009$; $d = .86$) and the cognitive reserve index ($p < .001$; $d = 1.18$) showed a significant difference between the groups on increasing frequency after the end of the program. No significant changes were observed in basic daily activities and social life activities. On the other hand, significant differences were found in the values of IDARE (State) ($p = .040$; $d = .66$) and IDB ($p = .001$; $d = 1.09$) reflecting a decrease in depression in EG (−3); however, an increase in this variable was observed in the CG during follow-up (+0.55) (Table 5).

The calculation of the magnitude of the differences between the score change (T1 to T3) is generally high ($d > .80$) between the EG and the CG.

Discussion

This study provides evidence on the efficacy of PIRCO to help improving long-term cognitive reserve and functioning in MS patients by generalize cognitive leisure activities.

Previous works have described that physical exercises^{31,39} as well as cognitive rehabilitation programs^{40,41} are profitable in optimizing cognitive functions in patients with MS. Nevertheless, other authors state that combining this two type of intervention (physical and cognitive) could be applied in the context of a cognitive reserve bettering, in a global form, the ill effects on motor and cognitive disability in MS.^{12,31} Our studies are in line with previous one, because we have observed that the combined training was effective, improving some diverse functions, such as verbal and visu-

ospatial memory, information processing speed, attention, and working memory. Recent studies on MS agree on the positive effect of the motor-cognitive approach on visuospatial memory performance and the speed to process information, attention, and working memory compared to the group that received physical training only.^{11,12}

Another important variable that can influence the effectiveness of the PIRCO rehabilitation program was group work. During these activities, the EG participants had the opportunity to reinforce their behavior; as well as the development of knowledge and skills necessary to configure and transfer cognitive and leisure activities into daily life one. These group interventions could contribute to increase up the cognitive reserve index once the rehabilitation program is concluded (T2), as well as the benefits observed in the long term (T3), and to reinforce the importance of transferring the benefits that education provides for cognitive health and compensation strategies to activities of daily living. These results coincide with few studies that examine the importance of group sessions included in cognitive rehabilitation programs in MS, which seem to reinforce the learning of new compensation strategies taught by therapists^{7,42} and also help their successful experience in MS. daily life.⁷

The effect observed through PIRCO on the increase of cognitive reserve and leisure activities, in the participants, was a key factor in reinforcing the hypothesis that premorbid physical and intellectual activities not only act as a buffer for limiting the MS-related damage but also as functional reserve that can be retrieved by task-oriented training to promote recovery through rehabilitation.⁴³ These results are consistent to a study to identify patients with MS, a relationship among leisure activities generating new learnings (Ex. Reading and writing) with changes been related to brain reserves (larger volume in the hippocampus), as well as bettering memory functions up.¹⁴ Schwartz et al.⁴⁴ in a longitudinal study proved that leisure and cognitive activities been more profitable in patients having larger active cognitive reserves, were: self-willing activities, attending religious or spiritual organizations and social participation. As some authors suggest¹⁵ these results are an evidence of the continuous participation in cognitive stimulating activities are capable of creating a cognitive reserve and acting as a protector against worsening disability related to MS, been summed up to previous cognitive reserve in patients.

It is also important to point out the effect caused by training in such a relevant aspect as the well affair. These results suggest that this multimodal rehabilitation approach, creates a notorious effect on the depressive symptoms present in MS patients. These results are in consonance to previous studies carried out by our research group,¹¹ as well as studies carried out by other authors.¹²

To conclude, it is necessary, to point out an important limitation in this study. The inclusion of a control group made up of MS patients treated with cognitive treatment but not with group therapy; in addition, a group with equivalent demographic characteristics but without MS would have yielded more consistent data of short- and long-term effectiveness in improving cognitive reserve in daily live activities. Hence, although the results are conclusive, it is not possible to conclude, in a certain way, the value of combination therapy. However, it is possible to conclude that this study provides evidence of the adequate benefits of a

rehabilitation program, in which cognitive exercises were combined with physical exercises, as well as group sessions; aimed at improving cognitive performance, reserve-building activities, and emotional states.

Regarding the control of confounding variables, there is no significant difference in education between EG and CG. However, the EG group shows a greater number of participants with academic studies and fewer participants with low education (in our case, 9th grade) than the CG group. Due to the relationship between cognitive reserve and education level, the random block design could be useful in future studies.

Finally, although it was not the objective of our study, the analysis of the caregivers' work could be of vital importance in the effect of the intervention. Therefore, the control of some family factors in the rehabilitation process could validate the approach of the program in the future.

Conflict of interests

The authors declare that they have no conflict of interest.

References

1. Chiaravalloti ND, DeLuca J. Cognitive impairment in multiple sclerosis. *Lancet Neurol.* 2006;7:1139–51, [http://dx.doi.org/10.1016/S1474-4422\(08\)70259-X](http://dx.doi.org/10.1016/S1474-4422(08)70259-X).
2. Rocca MA, Amato MP, De Stefano N, Enzinger CH, Geurts JJ, Penner I, et al. Clinical and imaging assessment of cognitive dysfunction in multiple sclerosis. *Lancet Neurol.* 2015;14:302–17, [http://dx.doi.org/10.1016/S1474-4422\(14\)70250-9](http://dx.doi.org/10.1016/S1474-4422(14)70250-9).
3. Arnett P, Forn C. Evaluación neuropsicológica en la esclerosis múltiple. *Rev Neurol.* 2007;44:166–72, <http://dx.doi.org/10.33588/rn.6411.2016312>.
4. Chiaravalloti ND, DeLuca J. The influence of cognitive dysfunction on benefit from learning and memory rehabilitation in MS: a sub-analysis of the MEMREHAB trial. *Mult Scler.* 2015;21:1575–82, <http://dx.doi.org/10.1177/1352458514567726>.
5. Perez-Martin MY, Gonzalez-Platas M, Eguia-Del Rio P, Croissier-Elias C, Jimenez Sosa A. Efficacy of a short cognitive training program in patients with multiple sclerosis. *Neuropsychiatric disease and treatment.* *Neuropsychiatr Dis Treat.* 2017;13:245–52, <http://dx.doi.org/10.2147/NDT.S124448>.
6. Chiaravalloti ND, Goverover Y, Costa SL, DeLuca J. A pilot study examining Speed of Processing Training (SPT) to improve processing speed in persons with multiple sclerosis. *Front Neurol.* 2018;9:685, <http://dx.doi.org/10.3389/fneur.2018.00685>.
7. Stuifbergen AK, Becker H, Pérez F, Morison J, Kullberg V, Todd A. A randomized controlled trial of a cognitive rehabilitation intervention for persons with multiple sclerosis. *Clin Rehabil.* 2012;26:882–93, <http://dx.doi.org/10.1016/j.dhjo.2018.02.001>.
8. O'Brien AR, Chiaravalloti N, Goverover Y, Deluca J. Evidenced-based cognitive rehabilitation for persons with multiple sclerosis: a review of the literature. *Arch Phys Med Rehabil.* 2008;89:761–9, <http://dx.doi.org/10.1016/j.apmr.2007.10.019>.
9. Mitolo M, Venneri A, Wilkinson ID, Sharrack B. Cognitive rehabilitation in multiple sclerosis: a systematic review. *J Neurol Sci.* 2005;354:1–9, <http://dx.doi.org/10.1016/j.jns.2015.05.004>.

10. Lampit A, Heine J, Finke C, Barnett MH, Valenzuela M, Wolf A, et al. Computerized cognitive training in multiple sclerosis: a systematic review and meta-analysis. *Neurorehabil Neural Repair*. 2019;33:695–706, <http://dx.doi.org/10.1177/1545968319860490>.
11. Jiménez-Morales RM, Herrera-Jiménez LF, Macías-Delgado Y, Pérez-Medinilla YT, Díaz-Díaz SM, Forn C. Entrenamiento cognitivo combinado con ejercicios aeróbicos en pacientes con esclerosis múltiple: estudio piloto. *Rev Neurol*. 2017;64:489–95, <http://dx.doi.org/10.33588/rn.6411.2016312>.
12. Barbarulo AM, Lus G, Signoriello E, Trojano L, Grossi D, Esposito M, et al. Integrated cognitive and neuromotor rehabilitation in multiple sclerosis: a pragmatic study. *Front Behav Neurosci*. 2018;12:196, <http://dx.doi.org/10.3389/fnbeh.2018.00196>.
13. Mani A, Chohedri E, Ravanfar P, Mowla A, Nikseresht A. Efficacy of group cognitive rehabilitation therapy in multiple sclerosis. *Acta Neurol Scand*. 2018;137:589–97, <http://dx.doi.org/10.1111/ane.12904>.
14. Sumowski JF, Rocca MA, Leavitt VM, Riccitelli G, Meani A, Comi G, et al. Reading, writing, and reserve: literacy activities are linked to hippocampal volume and memory in multiple sclerosis. *Mult Scler*. 2016;22:1621–5, <http://dx.doi.org/10.1177/1352458516630822>.
15. Sandroff BM, Schwartz CE, DeLuca J. Measurement and maintenance of reserve in multiple sclerosis. *J Neurol*. 2016;263:2158–69, <http://dx.doi.org/10.1007/s00415-016-8104-5>.
16. Stern Y. What is cognitive reserve? Theory and research application of the reserve concept. *J Int Neuropsychol Soc*. 2002;8:448–60, <http://dx.doi.org/10.1017/S1355617702813248>.
17. Sumowski JF, Rocca MA, Leavitt VM, Riccitelli G, Comi G, Deluca J, et al. Brain reserve and cognitive reserve in multiple sclerosis. Brain reserve and cognitive reserve in multiple sclerosis: What you've got and how you use it. *Neurology*. 2013;80:2186–93, <http://dx.doi.org/10.1212/WNL.0b013e318296e98b>.
18. Sumowski JF, Glenn RW, Deluca J, Chiaravalloti N. Intellectual enrichment is linked to cerebral efficiency in multiple sclerosis: functional magnetic resonance imaging evidence for cognitive reserve. *Brain*. 2010;33(Pt 2):362–74, <http://dx.doi.org/10.1093/brain/awp307>.
19. Santangelo G, Altieri M, Gallo A, Trojano L. Does cognitive reserve play any role in multiple sclerosis? A meta-analytic study. *Mult Scler Relat Disord*. 2019;30:265–76, <http://dx.doi.org/10.1016/j.msard.2019.02.017>.
20. Sumowski JF. Cognitive reserve as a useful concept for early intervention research in multiple sclerosis. *Front Neurol*. 2015;6:176, <http://dx.doi.org/10.3389/fneur.2015.00176>.
21. Schwartz CE, Rapkin BD, Healy BC. Reserve and reserve-building activities research: key challenges and future directions. *BMC Neurosci*. 2016;17:62, <http://dx.doi.org/10.1186/s12868-016-0297-0>.
22. Polman CH, Reingold SC, Banwell B, Clanet M, Cohen JA, Filippi M, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol*. 2011;69:292–302, <http://dx.doi.org/10.1002/ana.22366>.
23. Lubling FD, Reingold SC. Defining the clinical course of multiple sclerosis: results of an international survey. *Neurology*. 1996;46:907–11, <http://dx.doi.org/10.1212/wnl.46.4.907>.
24. Sepulcre J, Vanotti S, Hernández R, Sandoval G, Cáceres F, Garcea O, et al. Cognitive impairment in patients with multiple sclerosis using the Brief Repeatable Battery-Neuropsychology test. *Mult Scler*. 2006;12:187–95, <http://dx.doi.org/10.1191/1352458506ms1258oa>.
25. León-Estrada I, García-García J, Roldán-Tapia L. Escala de reserva cognitiva: ajuste del modelo teórico y baremación. *Rev Neurol*. 2017;64:7–16, <http://dx.doi.org/10.33588/rn.6401.2016295>.
26. León I, García J, Roldán-Tapia L, León I, García-García J, Roldán-Tapia L. Estimating cognitive reserve in healthy adults using the cognitive reserve scale. *PLOS ONE*. 2014;9:e102632, <http://dx.doi.org/10.1371/journal.pone.0102632>.
27. Beck AT, Steer RA. *BDI: Beck Depression Inventory Manual*. New York: Psychological Corporation; 1987.
28. Sanz J, Vázquez C. Fiabilidad, validez y datos normativos del inventario para la depresión de Beck. *Psycothema*. 1998;10:303–18.
29. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. *Manual for the state-trait anxiety inventory*. Palo Alto CA: Consulting Psychologists Press; 1983.
30. Krch D, Diaz-Orueta U, Santana E, Vasquez D. Rehabilitación de la Memoria: historia, factores implicados y enfoques de tratamiento. *Revista de Neuropsicología, Neuropsiquiatría y Neurociencias*. 2016;1:91–121.
31. Sandroff BM, Diggs MD, Bamman MB, Cuttred GR, Baird JF, Jones CD, et al. Protocol for a systematically-developed, phase I/II, single-blind randomized controlled trial of treadmill walking exercise training effects on cognition and brain function in persons with multiple sclerosis. *Contemp Clin Trials*. 2019;87:105878, <http://dx.doi.org/10.1016/j.cct.2019.105878>.
32. Sohlberg MM, Mateer CA. Effectiveness of an attention training program. *J Clin Exp Neuropsychol*. 1987;9:117–30, <http://dx.doi.org/10.1080/01688638708405352>.
33. Luria AR. *El cerebro y acción*. La Habana: Editorial Pueblo y Educación; 1989.
34. Filippi M, Riccitelli G, Mattioli F, Capra R, Stampatori CH, Pagani E, et al. Multiple sclerosis: effects of cognitive rehabilitation on structural and functional MR imaging measures – an explorative study. *Radiology*. 2012;262:932–40, <http://dx.doi.org/10.1148/radiol.11111299>.
35. Mattioli F, Bellomi F, Stampatori C, Capra R, Miniussi C. Neuroenhancement through cognitive training and anodal tDCS in multiple sclerosis. *Mult Scler*. 2015;22:222–30, <http://dx.doi.org/10.1177/1352458515587597>.
36. Field A. *Discovering statistics using IBM SPSS statistics*. 4 ed. London, England: SAGE Publications; 2013.
37. Olejnik S, Algina J. Generalized eta and omega squared statistics: measures of effect size for some common research designs. *Psychol Methods*. 2003;8:434–47, <http://dx.doi.org/10.1037/1082-989X.8.4.434>.
38. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. New York: Lawrence Erlbaum Associates Publishers; 1988.
39. Sandroff GR, Wylie BP, Sutton BP, Johnson CL, DeLuca J, Motl RW. Treadmill walking exercise training and brain function in multiple sclerosis: preliminary evidence setting the stage for a network-based approach to rehabilitation. *Mult Scler J Exp Transl Clin*. 2018;4, <http://dx.doi.org/10.1177/2055217318760641>, 055217318760641.
40. Gromisch ES, Fiszdon JM, Kurtz MM. The effects of cognitive-focused interventions on cognition and psychological well-being in persons with multiple sclerosis: a meta-analysis. *Neuropsychol Rehabil*. 2018;30:767–86, <http://dx.doi.org/10.1080/09602011.2018.1491408>.
41. Lincoln NB, Bradshaw LE, Constantinescu CS, Day F, Drummond AER, Fitzsimmons D, et al. Cognitive rehabilitation for attention and memory in people with multiple sclerosis: a randomized controlled trial (CRAMMS). *Clin Rehabil*. 2010;34:229–41, <http://dx.doi.org/10.1177/0269215519890378>.
42. Brissart H, Leroy M, Morel E, Baumann C, Spitz E, Debouverie M. Cognitive rehabilitation

- in multiple sclerosis. *Neurocase*. 2013;19:553–65, <http://dx.doi.org/10.1080/13554794.2012.701644>.
43. Castelli L, De Giglio L, Haggiag S, Traini A, De Luca F, Ruggieri S, et al. Premorbid functional reserve modulates the effect of rehabilitation in multiple sclerosis. *Neurol Sci*. 2020;41:1251–7, <http://dx.doi.org/10.1007/s10072-019-04237-z>.
44. Schwartz CE, Quaranto BR, Healy BC, Benedict RH, Vollmer TL. Cognitive reserve and symptom experience in multiple sclerosis: a buffer to disability progression over time? *Arch Phys Med Rehabil*. 2013;94:1971–81, <http://dx.doi.org/10.1016/j.apmr.2013.05.009>.