




ELSEVIER

## Allergología et immunopathología

Sociedad Española de Inmunología Clínica,  
Alergología y Asma Pediátrica

[www.elsevier.es/ai](http://www.elsevier.es/ai)



### ORIGINAL ARTICLE

# Level of serum neutrophil gelatinase-associated lipocalin in childhood asthma



A.Z. Akelma<sup>a,\*</sup>, M.K. Kanburoglu<sup>b</sup>, M.N. Cizmeci<sup>b</sup>, E. Mete<sup>c</sup>, F. Catal<sup>d</sup>, N. Tufan<sup>b</sup>

<sup>a</sup> Division of Pediatric Allergy, Department of Pediatrics, Turgut Ozal University Medical School, Ankara, Turkey

<sup>b</sup> Department of Pediatrics, Turgut Ozal University Medical School, Ankara, Turkey

<sup>c</sup> Division of Pediatric Allergy, Department of Pediatrics, Pamukkale University Medical School, Ankara, Turkey

<sup>d</sup> Division of Pediatric Allergy, Department of Pediatrics, Inonu University Medical School, Malatya, Turkey

Received 30 July 2013; accepted 14 November 2013

Available online 15 March 2014

#### KEYWORDS

Asthma;  
Children;  
Extracellular matrix  
protein;  
Lipocalin-2;  
Remodeling

#### Abstract

**Background:** The role of neutrophil gelatinase-associated lipocalin (NGAL) in childhood asthma remains unknown. This study aimed to measure the serum levels of NGAL in children with asthma and to investigate the correlation between NGAL and transforming growth factor beta 1 (TGF-β1), a good indicator of airway remodeling in children with asthma.

**Methods:** This prospective, cross-sectional study was conducted on 75 children. Serum NGAL and TGF-β1 concentrations were measured by the ELISA method. Complete blood count, high sensitive C reactive protein (hsCRP), eosinophil cationic protein (ECP), and total serum IgE were investigated in the study population. Atopy in the asthma group was investigated using a skin prick test and specific IgE measurements.

**Results:** Forty-three asthmatic children and 32 healthy children were enrolled in the study. Total eosinophil numbers, white blood cell count, total serum IgE levels and ECP levels were significantly higher in the asthma group than in the control group ( $p < 0.05$ ). Similarly, serum TGF-β1 levels were significantly higher in children with asthma ( $p = 0.012$ ). The difference in NGAL levels between the groups was insignificant ( $p = 0.268$ ). NGAL levels did not show a significant correlation with total IgE, ECP, eosinophil numbers and TGF-β1 levels ( $p > 0.05$ ).

**Conclusion:** As a conclusion, while elevated TGF-β1 levels in children with asthma might be regarded as an indicator of airway remodeling, we did not find a similar prediction strength for NGAL. Further studies are required to better identify the role of NGAL in childhood asthma and to determine its potential use as a clinical marker.

© 2013 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

\* Corresponding author.

E-mail addresses: [akelma@gmail.com](mailto:akelma@gmail.com) (A.Z. Akelma), [madduckk@gmail.com](mailto:madduckk@gmail.com) (M.K. Kanburoglu), [nevzatcizmeci@hotmail.com](mailto:nevzatcizmeci@hotmail.com) (M.N. Cizmeci), [eminmete@gmail.com](mailto:eminmete@gmail.com) (E. Mete), [ferhatcatal@gmail.com](mailto:ferhatcatal@gmail.com) (F. Catal), [nailetufan@gmail.com](mailto:nailetufan@gmail.com) (N. Tufan).

## Introduction

Asthma is one of the most common chronic diseases, estimated to be affecting 300 million people worldwide, with an increasing prevalence especially among children. Asthma is a clinical diagnosis based on episodic symptoms and variable airways obstruction. It is also characterized by variable degrees of chronic inflammation and structural alterations in the airways. These structural alterations, generally called airway remodeling, encompass complex changes in composition, content, and organization of the various cellular and molecular constituents of the airway wall. The most important abnormalities are epithelial detachment, goblet cell hyperplasia, subepithelial thickening, hyperplasia and hypertrophy of airway smooth muscle, bronchial gland enlargement, angiogenesis, and alterations in the extracellular matrix component.<sup>1-3</sup>

Neutrophil gelatinase-associated lipocalin (NGAL) is a 25 kDa glycoprotein first identified as a matrix protein of specific granules of human neutrophils.<sup>4</sup> NGAL is secreted by neutrophils<sup>4</sup> and other cells such as respiratory<sup>5</sup> and intestinal<sup>6</sup> epithelial cells, vascular endothelial cells,<sup>7</sup> adipose tissue,<sup>8</sup> macrophages<sup>9</sup> and tubuli cells in the kidneys.<sup>10</sup> The role of extracellular matrix proteins in the process of airway remodeling is well known. Matrix metalloproteinase-9 (MMP-9) and NGAL have been shown to be particularly associated with lung capacity and clinical severity. Several studies performed on patients with asthma, pulmonary emphysema, chronic obstructive pulmonary disease (COPD) revealed increased MMP-9 and NGAL levels in bronchoalveolar lavage (BAL) fluid samples, probably as a result of structural alterations in the airways.<sup>11-15</sup> Additionally, a recent study reported significantly higher plasma NGAL levels in COPD patients than in control patients.<sup>16</sup>

The role of NGAL as a potential marker of disease severity in childhood asthma remains unknown. Research involving invasive bronchoscopy techniques in young children is limited; hence studies have focused on systemic markers indicating the possible structural alterations in the airways. In the present study we investigated the association between serum NGAL levels and clinical and laboratory parameters in children presenting with asthma. Additionally, we investigated the correlation between NGAL and transforming growth factor beta 1 (TGF- $\beta$ 1), a good indicator of airway remodeling in children with asthma.<sup>17,18</sup>

## Materials and methods

### Study subjects

This prospective, cross-sectional study was conducted on 75 children, 43 of whom suffered from asthma presenting with wheezing to the Fatih University pediatric allergy outpatient clinic and 32 of whom were healthy controls. The asthma group included children of ages  $\leq 6$  years, with at least four wheezing attacks during the previous year, and ages  $> 6$  years who were diagnosed clinically and functionally according to GINA criteria.<sup>2</sup> Children of similar age and gender with no history of allergic disease or wheezing were selected as the control group. Healthy children were chosen among those being brought for routine controls. Those with

chronic diseases (e.g. malnutrition, anatomic malformation of the respiratory system, chronic lung disease, heart disease, gastro-esophageal reflux disease, cystic fibrosis) were excluded from the study.

Venous fasting blood samples were collected into Vacutte tubes (Greiner Bio-One, Monroe, NC) and centrifuged at  $3000 \times g$  for 15 min at  $4^{\circ}\text{C}$ . Serum samples were stored at  $-80^{\circ}\text{C}$  for not more than 6 months. Levels serum of TGF- $\beta$ 1 was measured by the ELISA (enzyme linked immunosorbent assay) method using a ELX-800 system (RayBiotech, Norcross, GA, US). Serum NGAL concentrations were determined with BioVendor Human Lipocalin-2/NGAL ELISA kit (BioVendor GmbH, Heidelberg, Germany). The limit of detection for this assay was 0.02 ng/mL with total imprecision (as CV) <8% at concentrations of 23.63–68.19 ng/mL. The antibodies used in this ELISA are specific for human lipocalin-2. Complete blood count was investigated with the LH-780 system (Beckman Coulter Diagnostics, Image 8000, Brea, CA, USA). Levels of high sensitive C reactive protein (hsCRP) were measured by turbidimetric assay method using a Roche P 800 modular system (Hitachi, Tokyo, Japan). Levels of eosinophil cationic protein (ECP) were measured by the chemiluminescent assay method using an Immulite 2000 systems (Simens, Llanberis, Gwynedd, UK). The eosinophil counts were measured by LH-780 system (Beckman Coulter, Mervue, Galway, Ireland). Levels of total serum IgE were measured by the ECLIA (electrochemiluminescence) method using a ELX-800 system (DIAsource, Nivelles, Belgium).

Atopy in the asthma group was investigated using a skin prick test (SPT) and specific IgE (sIgE) measurements. A test was considered positive if the SPT results demonstrated a wheal with a mean diameter of at least 3 mm greater than that of a saline control. Each child was tested with a core battery of allergens (e.g. dust mite, cockroach, cat, dog, mold, grass, tree, weed, milk, egg, peanut) and a clinic-specific battery of locally relevant allergens (ALK Abelló, Hørsholm, Denmark). Spirometry (Vmax encore; VIASYS Healthcare Inc., Conshohocken, PA) and bronchodilator reversibility were defined as greater than a 12% or 200 ml change from baseline FEV<sub>1</sub>. These parameters were measured according to the GINA criteria.<sup>2</sup>

### Statistical analysis

Data analysis was performed using Statistical Package for the Social Sciences (SPSS) for Windows, version 16.0 (SPSS Inc., Chicago, IL, United States). All the continuous values were presented as mean  $\pm$  standard deviation (SD). The categorical values were presented as n (%). The homogeneous distribution of the data was evaluated using the Kolmogorov-Smirnov test. Homogeneity of variance was evaluated by the Levene test. In the comparisons of the groups, Student *t*-test was used. The chi-square test was used for categorical variables. Correlation between the independent parameters was investigated by bivariate (Pearson) correlation analysis. A *p*-value of less than 0.05 was considered as statistically significant.

The study was initiated upon approval by the Local Ethics Committee of Fatih University in accordance with the Helsinki Declaration. The written informed consent of the parent(s) of each subject was also obtained before the study.

**Table 1** Demographic and laboratory characteristics of study population.

	Asthma group (n=43)	Control group (n=32)	p
<i>Age, months</i>			
Mean ± SD	83.58 ± 50.12	68.40 ± 50.00	0.199 <sup>a</sup>
Minimum–maximum	17–181	9–179	
<i>Gender</i>			
Male, n (%)	27 (62.8)	17 (53.1)	0.40 <sup>b</sup>
<i>Atopic dermatitis, n (%)</i>	8 (18.6)	–	
<i>Allergic rhinitis, n (%)</i>	12 (17.9)	–	
<i>Exposure to smoking, n (%)</i>	13 (30.2)	2 (6.2)	0.01 <sup>b</sup>
<i>Atopy, n (%)</i>	35 (85.4)	–	
<i>Familial history of atopy, n (%)</i>	16 (37.2)	1 (3.7)	0.001 <sup>b</sup>

<sup>a</sup> Student t-test.<sup>b</sup> Chi-square test.

## Results

Forty-three asthmatic children (mean age:  $83.58 \pm 50.12$  months; 27 male) and 32 healthy children (control group) (mean age:  $68.40 \pm 50.00$  months; 17 male) were enrolled in the study. Mean ages and gender distribution were similar between the groups ( $p > 0.05$ ). The demographic characteristics of study population are presented in Table 1.

Total eosinophil numbers, white blood cell (WBC) numbers, total serum IgE levels and ECP levels were significantly higher in the asthma group than in the control group ( $p = 0.004$ ,  $p = 0.029$ ,  $p = 0.014$ , and  $p < 0.001$ ; respectively). Similarly, serum TGF-β1 (ng/mL) levels were significantly higher in children with asthma ( $1.48 \pm 1.07$ ,  $0.90 \pm 0.86$ ;  $p = 0.012$ ). Serum hsCRP levels, measured to exclude possible co-morbid infections, were similar between the groups ( $p > 0.05$ ). The difference in NGAL (ng/mL) levels between the groups was insignificant ( $109.69 \pm 85.00$ ,  $89.90 \pm 68.48$ ;  $p = 0.268$ ) (Table 2).

Children from the asthma group were classified into subgroups based on disease severity (12 mild intermittent, 26 mild persistent, 5 moderate persistent). None of these children presented with severe asthma. We found similar NGAL levels between the subgroups. While NGAL levels did not show a significant correlation with total IgE, ECP, eosinophil numbers and TGF-β1 levels ( $p > 0.05$ ), a weak

correlation was noted between NGAL and hsCRP levels ( $p = 0.015$ ,  $r = 0.292$ ).

## Discussion

Asthma is a disease of airway inflammation characterized by hyper-responsiveness and airway obstruction that lead to symptoms such as cough and wheezing. The remodeling process observed following chronic inflammation and airway obstruction in asthmatic patients may lead to structural alterations in the airways which in turn may result in a decrease in the lung functions and resistance to medical therapy.<sup>2</sup> Data from the Tucson Children's Respiratory Study suggest that pulmonary function in children who ultimately have persistent asthma is normal at birth, with deficits acquired by as early as 6 years of age.<sup>19</sup> Lung function data from a large cohort study, with follow-up from early childhood through age 35 years revealed that children with asthma had lower lung function during childhood and at age 35 years compared with that seen in cohort members without a history of childhood asthma.<sup>20</sup> This remodeling process has been demonstrated in children with asthma by means of endobronchial biopsy samples.<sup>21,22</sup> In addition to these data suggest that structural changes in the asthmatic children airway that might result in permanent lung function deficits occur early in the natural history

**Table 2** Comparison of laboratory parameters between asthma and control groups.

	Asthma group (n=43)	Control group (n=32)	*p
Eosinophils (count/mm <sup>3</sup> )	313.00 ± 292.64	141.55 ± 128.02	0.004
WBC (count/mm <sup>3</sup> )	9109 ± 3108	7680 ± 2077	0.029
Total IgE (IU/mL)	235.99 ± 489.17	36.10 ± 24.09	0.014
ECP (ng/mL)	45.37 ± 31.98	19.77 ± 18.91	<0.001
TGF-β1 (ng/mL)	1.48 ± 1.07	0.90 ± 0.86	0.012
hsCRP (mg/dL)	3.68 ± 5.96	3.26 ± 10.57	0.845
NGAL (ng/mL)	109.69 ± 85.00	89.90 ± 68.48	0.268

ECP, eosinophil cationic protein; hsCRP, high sensitive C reactive protein; NGAL, neutrophil gelatinase-associated lipocalin; TGF-β1, transforming growth factor beta 1; WBC, white blood cells.

Date are given mean ± SD.

\* Student t-test.

of asthma. A good understanding of the remodeling process and structural alterations observed in children with asthma is important for the follow up and management of the patients. Since research involving invasive bronchoscopy techniques in young children is limited, studies have focused on non-invasive methods. NGAL, which is an extracellular matrix protein, has emerged as a potential marker of airway remodeling in children with asthma.

NGAL is characterized as a specific granule of human neutrophils and is known to be secreted from a number of tissues including the respiratory epithelial cells<sup>5</sup> and macrophages.<sup>9</sup> Current research is focused on the role of NGAL on various diseases including acute kidney injury, obesity, cancer, etc.<sup>23–25</sup> Previous studies reported an increase in NGAL and MMP-9 levels in asthmatic patients with airway structural alterations.<sup>12–14</sup> In a recent study, NGAL levels, when compared to matched controls, were shown to be higher in COPD patients presenting with notable airway structural alterations and with a significant increase in their extracellular matrix protein levels. Taken together, one may suggest NGAL as a potential predictor of airway remodeling.<sup>16</sup> However, in the present study, we found similar serum NGAL levels in children with asthma and control subjects.

TGF-β1 is a profibrotic cytokine that is associated with airway fibrosis. TGF-β1 is believed to play an important role in most of the cellular biological processes leading to airway remodeling.<sup>18,26</sup> Another study reported increased TGF-β1 concentrations and expression in children with asthma.<sup>17</sup> Thus, TGF-β1 might be a key mediator of the airway remodeling in children with asthma. In this study although we found significantly higher serum TGF-β1 levels in children with asthma as compared to control subjects, the difference between NGAL levels were insignificant. This finding implies that NGAL is not a strong predictor of airway remodeling in children with asthma.

Our study has several limitations. First, the sample size was small. Second, the study group lacked of patients with severe asthma and third, concurrent bronchoalveolar examinations were missing. Higher serum TGF-β1 levels found in children with asthma as compared to control subjects indicates, however, an active airway remodeling and structural alteration process in these patients. Hence the absence of subjects with severe asthma in the study group may not actually be regarded as a substantial limitation.

To the best of our knowledge, this is the first study to investigate the role of NGAL in asthmatic children. As a conclusion, while elevated TGF-β1 levels in children with asthma might be regarded as an indicator of airway remodeling, we did not find similar prediction strength for NGAL. Further studies conducted on larger groups, preferably examined by means of bronchoscopic methods, are required to better identify the role of NGAL in childhood asthma and to determine its potential use as a clinical marker.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this investigation.

**Confidentiality of data.** The authors declare that they have followed the protocols of their work center on the publication of patient data and that all the patients included in the study have received sufficient information and have given their informed consent in writing to participate in that study.

**Right to privacy and informed consent.** The authors have obtained the informed consent of the patients and/or subjects mentioned in the article. The author for correspondence is in possession of this document.

## Conflict of interest

The authors have no conflict of interest to declare.

## Acknowledgment

**Funding/support:** This study is supported by the Scientific Research Fund of Fatih University.

## References

- Davies DE, Wicks J, Powell RM, Puddicombe SM, Holgate ST. Airway remodeling in asthma: new insights. *J Allergy Clin Immunol*. 2003;111:215–25, quiz 226.
- From the Global Strategy for Asthma Management and Prevention, Global Initiative for Asthma (GINA); 2011. Available from <http://www.ginasthma.org/>
- Mauad T, Bel EH, Sterk PJ. Asthma therapy and airway remodeling. *J Allergy Clin Immunol*. 2007;120:997–1009, quiz 1010–1.
- Kjeldsen L, Bainton DF, Sengelov H, Borregaard N. Identification of neutrophil gelatinase-associated lipocalin as a novel matrix protein of specific granules in human neutrophils. *Blood*. 1994;83:799–807.
- Cowland JB, Sorensen OE, Sehested M, Borregaard N. Neutrophil gelatinase-associated lipocalin is up-regulated in human epithelial cells by IL-1 beta, but not by TNF-alpha. *J Immunol*. 2003;171:6630–9.
- Nielsen BS, Borregaard N, Bundgaard JR, Timshel S, Sehested M, Kjeldsen L. Induction of NGAL synthesis in epithelial cells of human colorectal neoplasia and inflammatory bowel diseases. *Gut*. 1996;38:414–20.
- Bu DX, Hemdahl AL, Gabrielsen A, Fuxe J, Zhu C, Eriksson P, et al. Induction of neutrophil gelatinase-associated lipocalin in vascular injury via activation of nuclear factor-kappaB. *Am J Pathol*. 2006;169:2245–53.
- Kratchmarova I, Kalume DE, Blagoev B, Scherer PE, Podtelezhnikov AV, Molina H, et al. A proteomic approach for identification of secreted proteins during the differentiation of 3T3-L1 preadipocytes to adipocytes. *Mol Cell Proteomics*. 2002;1:213–22.
- Meheus LA, Fransen LM, Raymackers JG, Blockx HA, Van Beeumen JJ, Van Bun SM, et al. Identification by microsequencing of lipopolysaccharide-induced proteins secreted by mouse macrophages. *J Immunol*. 1993;151:1535–47.
- Mishra J, Ma Q, Prada A, Mitsnefes M, Zahedi K, Yang J, et al. Identification of neutrophil gelatinase-associated lipocalin as a novel early urinary biomarker for ischemic renal injury. *J Am Soc Nephrol*. 2003;14:2534–43.
- Betsuyaku T, Nishimura M, Takeyabu K, Tanino M, Venge P, Xu S, et al. Neutrophil granule proteins in bronchoalveolar lavage fluid from subjects with subclinical emphysema. *Am J Respir Crit Care Med*. 1999;159:1985–91.

12. Cundall M, Sun Y, Miranda C, Trudeau JB, Barnes S, Wenzel SE. Neutrophil-derived matrix metalloproteinase-9 is increased in severe asthma and poorly inhibited by glucocorticoids. *J Allergy Clin Immunol.* 2003;112:1064–71.
13. Metso T, Rytila P, Peterson C, Haahtela T. Granulocyte markers in induced sputum in patients with respiratory disorders and healthy persons obtained by two sputum-processing methods. *Respir Med.* 2001;95:48–55.
14. Karakoc GB, Yukselen A, Yilmaz M, Altintas DU, Kendirli SG. Exhaled breath condensate MMP-9 level and its relationship with asthma severity and interleukin-4/10 levels in children. *Ann Allergy Asthma Immunol.* 2012;108:300–4.
15. Klein T, Bischoff R. Physiology and pathophysiology of matrix metalloproteases. *Amino Acids.* 2011;41:271–90.
16. Eagan TM, Damas JK, Ueland T, Voll-Aanerud M, Mollnes TE, Hardie JA, et al. Neutrophil gelatinase-associated lipocalin: a biomarker in COPD. *Chest.* 2010;138:888–95.
17. Brown SD, Baxter KM, Stephenson ST, Esper AM, Brown LA, Fitzpatrick AM. Airway TGF-beta1 and oxidant stress in children with severe asthma: association with air-flow limitation. *J Allergy Clin Immunol.* 2012;129:388–96, 396.e381–8.
18. Yang YC, Zhang N, Van Crombruggen K, Hu GH, Hong SL, Bachert C. Transforming growth factor-beta1 in inflammatory airway disease: a key for understanding inflammation and remodeling. *Allergy.* 2012;67:1193–202.
19. Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. The Group Health Medical Associates. *N Engl J Med.* 1995;332:133–8.
20. Kelly WJ, Hudson I, Raven J, Phelan PD, Pain MC, Olinsky A. Childhood asthma and adult lung function. *Am Rev Respir Dis.* 1988;138:26–30.
21. Barbato A, Turato G, Baraldo S, Bazzan E, Calabrese F, Panizzi C, et al. Epithelial damage and angiogenesis in the airways of children with asthma. *Am J Respir Crit Care Med.* 2006;174:975–81.
22. Payne DN, Rogers AV, Adelroth E, Bandi V, Guntupalli KK, Bush A, et al. Early thickening of the reticular basement membrane in children with difficult asthma. *Am J Respir Crit Care Med.* 2003;167:78–82.
23. Akelma AZ, Abaci A, Ozdemir O, Celik A, Avci Z, Razi CH, et al. The association of serum lipocalin-2 levels with metabolic and clinical parameters in obese children: a pilot study. *J Pediatr Endocrinol Metab.* 2012;25:525–8.
24. Smetka M, Chudek J. Using NGAL as an early diagnostic test of acute kidney injury. *Ren Fail.* 2012;34:130–3.
25. Li C, Chan YR. Lipocalin 2 regulation and its complex role in inflammation and cancer. *Cytokine.* 2011;56:435–41.
26. Halwani R, Al-Muhsen S, Al-Jahdali H, Hamid Q. Role of transforming growth factor-beta in airway remodeling in asthma. *Am J Respir Cell Mol Biol.* 2011;44:127–33.