




ELSEVIER

Allergología et immunopathología

Sociedad Española de Inmunología Clínica,
Alergología y Asma Pediátrica

www.elsevier.es/ai



ORIGINAL ARTICLE

Real life study of three years omalizumab in patients with difficult-to-control asthma

J. Jesús López Tiro^{a,*}, E. Angélica Contreras Contreras^a, M. Elena Ramírez del Pozo^a, J. Gómez Vera^a, D. Larenas Linnemann^b



^a Department of Allergy and Clinical Immunology, H.R. Lic. Adolfo López Mateos, ISSSTE, Mexico City, Mexico

^b Allergy Staff, Hospital Medica Sur, Mexico City, Mexico

Received 9 September 2013; accepted 14 November 2013

Available online 26 April 2014

KEYWORDS

Severe asthma;
Omalizumab;
Biological;
Anti-IgE;
Difficult-to-control
asthma;
Hospitalisation;
Emergency room;
Asthmatic crisis;
Asthma exacerbation

Abstract

Background: Even though there are multiple options for the treatment of asthma, there still exists a fair group of patients with difficult-to-control asthma. We describe for the first time the real-world effects of three-year omalizumab treatment on patients with difficult-to-control asthma, seen in a social security hospital in a Latin American country.

Methods: Difficult-to-control asthmatic patients from the out-patient clinic of a regional hospital were recruited to receive a three-year omalizumab course. Efficacy parameters were asthma control test (ACT) score; FEV1; daily beclomethasone maintenance dose; and unplanned visits for asthma exacerbations (emergency room (ER), hospitalisations, intensive care).

Results: 52 patients were recruited, 47 completed the three-year treatment (42 female, 15–67 years, mean age 43.5). Comparing efficacy parameters of the year before omalizumab with the 3rd year of omalizumab: mean ACT improved from 12.4 to 20.5, mean FEV1 from 66.3% (standard deviation (SD) 19.1%) to 88.4% (SD 16.2%) of predicted, while mean beclomethasone dose reduced from 1750 to 766 mcg/day and there was a significant reduction in patients experiencing ER visits (from 95% to 1%, $p < 0.0001$), hospitalisation (38% to 2%, $p < 0.0001$) and intensive care (4% to 0, NS). Five patients discontinued omalizumab, two because of an adverse event (anaphylaxis, severe headache, both resolved without sequelae).

Conclusion: Omalizumab improved most clinical parameters of Mexican patients with difficult-to-control asthma. Especially the rates of ER visits and hospitalisation were significantly reduced, thus reducing costs. Omalizumab was generally well tolerated.

© 2013 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

Introduction

The diagnosis of difficult-to-control asthma is considered in those patients in whom asthma is not well controlled in spite of an adequate therapeutic strategy applied by

* Corresponding author.

E-mail addresses: drlopeztiro@gmail.com, marlar1@prodigy.net.mx (J.J. López Tiro).

a specialist for the duration of at least six months.¹ This variety and severity of asthma is complex in nature, due to the multiple phenotypes and endotypes that might lead to the persistence of the asthma symptoms, the poor response of the patient to conventional pharmacotherapy, the non-satisfactory evolution of the disease and the susceptibility of the patient to co-morbidities.² Apart from the patients with true difficult-to-control asthma there is a considerable group of patients with seemingly difficult-to-control asthma in which the uncontrolled state of their disease is due to other factors, generally poor compliance.

Approximately 5% of all asthmatic patients develop difficult-to-control asthma, resulting in an elevated rate of complications, emergency visits, hospitalisations and even admissions to the ICU.³ That is why almost half of the total annual cost of asthma treatment corresponds to the severe asthmatic patients.⁴ It is important to make an effort to determine the cause of the absence of asthma control. Predisposing factors include an adverse environment, poor compliance, inadequate dosing or a level of asthma maintenance treatment unbalanced in relation to the degree of asthma severity.

Of equal importance is the detection of co-morbidities, e.g. gastro-oesophageal reflux-disease, or the differential diagnosis with other pulmonary disorders, such as chronic obstructive pulmonary disease (COPD), allergic bronchopulmonary aspergillosis, pleuropulmonary eosinophilic syndromes, Churg Strauss syndrome and occupational asthma, among others.⁵⁻⁷ Several treatment options have been explored to gain control in severe asthmatic patients. Inhaled and systemic corticosteroids have been the cornerstone of most treatment variants and in some cases clinically satisfactory results have been obtained as for the symptomatic and spirometric control of asthma, but the side-effects of long-term high-dose corticosteroid treatment are well-known to the medical community and close monitoring of these patients is mandatory. A small proportion of difficult-to-control asthmatic patients develop corticosteroid resistance, defined as a less than 15% improvement of the forced expiratory volume in the first second (FEV₁) after a 14-day course of 40 mg prednisolone. Asthma control is generally limited in these patients.⁵⁻⁷

Omalizumab is a humanised monoclonal antibody that binds to the constant region (Fc) of immunoglobulin E (IgE). Omalizumab blocks the binding of IgE to its high-affinity receptor Fc_εR1, situated on the surface of the mast cells (and other cells). Consequently, activation of the mast cells and liberation of its pro-inflammatory mediators is inhibited. The efficacy and safety of omalizumab in patients – six years and above – with severe asthma and difficult-to-control asthma have been documented in multiple clinical trials.⁸ As a consequence, omalizumab has been integrated in major clinical guidelines on the treatment of asthma, as a treatment option in the last step of therapy, in those patients not well controlled on high dose inhaled corticosteroids in combination with long-acting beta-agonists.^{5,9-11} Therefore, the objective of this study is to describe the results obtained after three years of treatment with omalizumab in patients with moderate to severe difficult-to-control asthma in a tertiary hospital in Mexico City.

Materials and methods

We conducted an open prospective, observational, cohort study, describing the evolution of difficult-to-control moderate to severe asthmatic patients, recruited from the out-patients clinic of the Allergy and Immunology Department of the Regional Hospital *Licenciado Adolfo López Mateos* (ISSSTE), in Mexico City. Patients fulfilling the inclusion criteria were invited to a thirty-six months' course of omalizumab. The study was implemented from January 2009 through December 2012. Patients from 12 to 75 years of age, of either sex, fulfilling the American Thoracic Society (ATS) criteria for difficult-to-control asthma⁶ were invited to participate and recruited into the study once an informed consent form had been signed. The following were exclusion criteria: COPD, occupational asthma, cystic fibrosis, cardiac insufficiency grade 3 or higher according to the New York Heart Association classification,¹² severe pulmonary emphysema, disorders that cause laryngeal or trachea obstruction, carcinoid syndromes, hyperthyroidism, sleep apnoea and psychiatric disorders. An unacceptable dosing interval between two doses of omalizumab or a no-show of the patient for the next injection were elimination criteria.

Omalizumab (XOLAIR[®]) was administered subcutaneously to all patients. The omalizumab dose was calculated according to the guidelines of the manufacturer, taking into account total IgE serum levels and the patient's weight.

Depending on the administration schedule of omalizumab the patients were evaluated every 15 days or every month during the three-year surveillance. The Asthma Control TestTM (ACT) was taken at the beginning and at the end of the study.¹³ The ACTTM is a simple five-point questionnaire, which is self-completed by patients. Each item is scored from one (poor control) to five (good control), related to the frequency and severity of the symptoms and their impact on the patient's quality of life; the scores of each of each of the five questions are added up to give a final score with a maximum of 25. The total score of the ACTTM should be interpreted in the following way: 14 or less is considered as very poor control of asthma, 15–19 total score is un-controlled asthma, 20–24 total score reflects well controlled asthma and finally 25 points is completely controlled asthma. We compared initial ACTTM values of our patients with ACTTM scores at the end of three years of omalizumab administration to evaluate the impact of the monoclonal antibody on asthma control. Moreover, pulmonary function tests were carried out (World Spirometer EasyOne 2001) and pre-bronchodilator FEV₁ was recorded in percent of the predicted value, according to anthropometric characteristics, sex, race and a history of tobacco smoke exposure.

Before the study started and during the surveillance period unscheduled visits to the clinic because of asthma exacerbations were recorded. These unscheduled visits were classified according to their nature (emergency department, hospitalisation, intensive care unit (ICU)) and expressed as unscheduled visits per year. Additionally, the start dose and the final dose of beclomethasone were documented, as well as the intake of any systemic glucocorticosteroids. No detailed record was kept on other asthma medication taken, so only a general picture can be given on this issue. A log of adverse events possibly or




Figure 1 Mean Asthma Control Test scores, forced expiratory volume (% predicted) values and hospitalisations per year (nr) pre- and post-three-year omalizumab treatment. $N=47$ patients.

probably related to the administration of omalizumab was carefully updated at each clinic visit.

All included patients continued normally with their regular asthma maintenance treatment in a step-wise approach as indicated by the 2010 Global INitiative on Asthma (GINA) guidelines.⁵ All records were downloaded into an electronic database and we used the 19th version of the SPSS program for statistical analysis.

Results

Fifty-six patients were pre-selected as possible candidates for our study, but four patients were excluded from the trial (3 COPD, 1 sleep apnoea), leaving 52 patients who started omalizumab administration (42 females, 15–67 years, mean age 43.5 years).

The mean omalizumab dose was 279.2 mg monthly. Five patients discontinued the study because of the following reasons: pulmonary tuberculosis, mild anaphylaxis, severe headache, poor compliance with the administration of the study medication and weight loss of more than 20 kg. A thorough clinical history showed that the weight loss of the patient had already started before the administration of omalizumab; the patient was referred to the department of internal medicine for further investigation. The event was judged as not related to the administration of omalizumab by the investigator.

Forty-seven patients finished the three year surveillance period. The results of the per protocol analysis of the ACT™ showed a mean score of 12.4 points before the first administration of omalizumab (=very poorly controlled asthma). At the end of the three years of omalizumab administration the mean group score was 20.47 points (=controlled asthma). At patient level, the mean personal ACT™ improvement was 8.06 points (median 8, SD 3.57). Thirty-five patients (74.4%) reached the 20–25 interval of well-controlled asthma, twelve patients maintained a score between 15 and 19, thus in the range of uncontrolled asthma (Fig. 1).

Before the administration of omalizumab, at study-start, pulmonary function testing showed a general group mean




Figure 2 Mean daily dose of beclomethasone before starting omalizumab and after three years of administration, bars indicate one standard deviation. $N=47$ patients.

FEV₁ of 66.3% of the predicted values (SD 19.1%). After the three years of omalizumab the group mean FEV₁ of the per protocol population had risen to 88.4% of predicted values (SD 16.2%) (Fig. 1). For the intention-to-treat analysis the mean FEV₁ values changed from 66.7% pre-omalizumab to 86.1% after 36 months of omalizumab.

At study-start 45 patients (95.7%) had had at least one emergency visit over the past year (mean 1.94 visits/year, SD 1.3), 18 patients (38.2%), reported at least one hospitalisation due to an asthma exacerbation the year before study-start and two patients (4.2%) had been admitted to the intensive care unit (ICU) because of severe, near fatal asthma exacerbations (Table 1).

After three years of omalizumab administration a clear reduction in all types of unscheduled visits was seen. During the three years of the trial, only nine (19.1%) of the 47 patients that concluded the surveillance period had an emergency room visit because of an asthma attack, only one patient (2.1%) was hospitalised and there was not one ICU admission (Fig. 1).

As for the inhaled corticosteroid dosing, all patients used high-dose therapy before starting omalizumab (mean 1750 mcg/day, SD 345 mcg). At the end of our three-year study, five patients (10.7%) were completely weaned off of their inhaled corticosteroids, while the mean beclomethasone dose for the remaining 42 patients was of 765 mcg/day (SD 654 mcg) (Fig. 2).

The year before being put on omalizumab all patients had received continuous- or intermittently oral corticosteroids (OCS, deflazacort ranging from 24 to 60 mg daily, prednisone 15 to 75 mg daily and some received intravenous methylprednisolone 125 to 250 mg during exacerbations. At the end of the trial, no patient was receiving daily OCS anymore; some did however still need short-time pulses for exacerbations. At study start almost all patients used LABA, specifically salmeterol 25/50 mcg, MDI, at doses ranging from 50 up until 100 mcg daily. However, the use of this treatment was intermittent, as this drug was often not available to our patients. In the majority of the subjects this dose was reduced or even suspended in the course of the trial, in line with the ICS dose reduction. At the end of the trial, there were still five patients receiving 25/50 mcg once daily.

Table 1 Mean number of yearly^a unscheduled clinic visits for asthma exacerbations, before and at the end of three years omalizumab ($N=47$ pts).

Nature of the visit	At study-start	During the three years of omalizumab	<i>p</i>
Emergency room visits	45 patients ^b (95.7%)	9 (19.1%)	<0.0001
Hospitalisations	18 patients (38.2%)	1 (2.1%)	<0.0001 ^{**}
Intensive care admissions	2 patients (4.2%)	0	NS

NS: not significant.

^a Mean number of visits/year over the last 2–3 years before starting omalizumab (recall data) as compared to registered visits during the third year of omalizumab treatment.

^b Mean 1.94 visits per year.

** With Yates' correction.

As for Montelukast: almost all patients have been on Montelukast 10 mg *ante nocte* throughout the study, as this is standard treatment in our hospital. A minority of patients received theophylline at the start of the study and some continued it during the trial.

During the three years of the study four adverse events were documented (see Fig. 3): one case of osteo-articular pain, two cases of mild headache and one case of a vasovagal syncope. All four cases received specific treatment and all patients concluded the study without interrupting the administration of omalizumab. In conclusion, we registered four adverse events over the three-year period of administration of omalizumab in the 49 patients; none of the events were considered life-threatening and none of these four events led to further drop-outs.

Discussion

The present study shows that in difficult-to-control, asthmatic patients in a specialised hospital in Mexico City, three

years of omalizumab treatment resulted in an important improvement in asthma control in most patients, as well as an improved lung function and a significant reduction in unscheduled visits to the emergency room, hospitalisations and admissions to ICU due to asthma exacerbations. At the end of the study the patients showed notable clinical improvement compared to the pre-study condition of their asthma as we demonstrated with the clear rise in ACT score. At study-start all patients were classified as poorly and very poorly controlled, while at the end of three years of omalizumab two thirds had reached the level of well-controlled asthma, as expressed by the group mean ACT score of 20.47 points.

Lung function testing is one of the prime objective measurements for follow-up of asthmatic patients, especially those patients with difficult control. The three years' treatment with omalizumab significantly improved the mean FEV₁ from 66.3% to 88.4% of predicted. As such the results of the ACT and of lung-function testing are in the same line.

Finally, the most important parameter of asthma control is the rate of emergency visits and hospital/ICU admissions because of asthma exacerbations. Before study-start practically all patients reported several unscheduled visits, because of their asthma per year and even two patients had a history of ICU admission because of a near-fatal asthma attack. These numbers improved impressively with the anti-IgE antibody administration, reducing unscheduled emergency room visits to below ten percent and only one hospital admission over the last year of treatment. The impact of such a reduction is not to be under-estimated as it does not only significantly improve the patient's quality of life, but it also reduces work-loss and it reduces – in a much greater extend – direct and in-direct health care costs of asthma.

A post-marketing surveillance trial of 280 German asthmatic patients¹⁴ showed similar results to our surveillance trial: after six month omalizumab there was a reduction in unscheduled health care contacts and hospitalisations of around 80%. Twelve months omalizumab in 12–20-year-old asthmatics reduced the need for systemic corticosteroids almost to zero,¹⁵ just as in our population. Compared to these and other data published, concerning the impact of omalizumab on emergency visits and hospitalisations in real-life situations¹⁶ several factors might influence the even better results we found: the above mentioned post-marketing surveillance trials report on the effect of omalizumab administration over a shorter period of time. Post-marketing results on three years are scarce. Another




Figure 3 Number of patients with adverse events over the three years of administration of omalizumab, judged by the investigator(s) as probably related to the administration of the monoclonal. $N=47$ patients.

factor that might play a role is the fact that our hospital is a reference hospital for patients of the social security system. It might be possible that for some patients optimal asthma management was only installed after they were referred to our clinic, months before trial recruitment, thus improving their asthma control even more than it would have improved with only omalizumab.

Over the past years asthma still remains a frequent cause of hospitalisation. Demographic and health statistics show it continues to cause remarkable morbidity among the population of both high-level and low-level income countries.^{17,18}

With the venue of the era of biologics for the control of diseases, these new treatment options have been shown to be effective in controlling severe forms of several disorders, uncontrolled until that moment with even the best practice. Omalizumab is a monoclonal antibody that has nowadays a well-accepted place in the management of severe asthma, as numerous clinical trials have shown its efficacy in severe cases of asthmatic airway inflammation in difficult-to-control patients. At present many guidelines have given omalizumab a place in the last step of the escalated treatment recommendations.^{5,10,11}

Investigators had been looking for alternative treatment options that might reduce the adverse effects of prolonged high-dose corticosteroid use or, even worse, of frequent systemic gluco-corticosteroid (GCS) bursts. As such, the GCS reducing capacity of several immunosuppressive drugs was explored, e.g. chloroquine, dapsone, azathioprine, salts of gold, cyclosporine, methotrexate, to mention some. However, the clinical trials with these agents, used as GCS saver medication in severe asthma, have not been able to show consistent results. Based on the results of Cochrane meta-analysis these agents have received the level of evidence B and C, because of some contradictory outcomes. Although their use has shown some benefit in severe asthmatic patients, frequently observed safety issues and other potentially harmful effects limit their usefulness in clinical practice. As a consequence their indication in severe, difficult-to-control asthmatic patients is still a matter of debate and much room is left to the judgement of the treating physician as for the decision to prescribe or refrain from prescription in this small subgroup of asthmatics.¹⁹⁻²⁴ Intravenous gamma globulin offers some benefit to GCS-dependent asthmatic patients, but here again studies show contradictory results. Additionally, the adverse events that have been described with IVIG cannot be neglected (e.g. fever, aseptic meningitis, urticaria).²⁴

Over the past years several biological treatments have been developed, with monoclonal antibodies directed against various pro-inflammatory molecules. Tumour necrosis factor alpha (TNF-alpha) is one of the molecules involved in the pathophysiology of severe asthma.²⁵ Trials have been designed to test anti-TNF-alpha antibodies, such as the monoclonal chimeric antibody infliximab that binds TNF-alpha and inactivates its inflammatory capacity. Preliminary results have shown an improvement in the quality of life of severe asthmatic patients, a mild improvement in lung function and a reduction in the hospitalisation rate. However, not all trials have shown positive results, some of them recruited a very limited number of patients and the reported side-effects (tuberculosis, infections, and others), have made

the acceptance of infliximab as regular treatment for severe asthma poor.²⁶

Following the same line of thinking industry has sought other monoclonal antibodies to block different molecules of inflammatory pathways in asthma. Some examples are mepolizumab, that blocks interleukin 5; lumiliximab, an anti-CD23 antibody, pascolizumab, a monoclonal antibody that interacts with interleukin 4; lebrikizumab, an anti-IL-13 antibody and inhibitor of periostin. Thus far, results have been promising, but these molecules are still in the experimental phase and consequently they have no official place in the asthma treatment yet.²⁶⁻²⁸

As such omalizumab is to date the sole treatment accepted in official guidelines in the last step of asthma treatment as corticosteroid safer, without being an immunosuppressant. In our study the mean dose of inhaled corticosteroid was significantly reduced. Those patients who completed the three years of omalizumab administration diminished the inhaled beclomethasone dose from the high to the low-middle range, without losing control of their asthma. 10.5% were even able to withdraw inhaled corticosteroids completely. As such we showed that omalizumab has an important corticoid sparing effect, reducing concomitantly the possible side-effects of prolonged ICS use.

Another prime concern in our cohort study was the safety of the administered biological substance, omalizumab; we observed our patients closely for possible side-effects. We reported one case of anaphylaxis after the second administration, a serious adverse event that has been documented in rare cases after the first several administrations of this medication.²⁹ As more than one organ system was involved in the symptoms of our patient after contact with a substance that might cause anaphylaxis, according to international guidelines we had to catalogue the reaction as anaphylaxis,³⁰ even though the symptoms were not of severe intensity. For safety reasons it was decided to exclude the patient from further omalizumab treatment. A recent review on this subject concluded that the data document low risk of anaphylaxis with omalizumab use. Moreover, the authors concluded that additional research is needed to elucidate the mechanism by which omalizumab causes anaphylaxis,³¹ as at first sight it seems a paradox that an anti-IgE antibody can induce massive mastcell degranulation.²⁶

In general, over the three years of omalizumab treatment there were few adverse events and they were generally mild in intensity, including headache, myalgia, and arthralgia as the most frequent ones. In only one of these patients omalizumab was discontinued (severe headache).

In this respect, our real-world data are in agreement with those previously reported in a randomised, placebo controlled clinical trial with omalizumab (the INNOVATE study), in which omalizumab was administered to 419 severe asthmatic patients for 28 weeks.²⁷ At study-start these patients were also on high dose ICS, mostly combined with long acting beta-agonists and courses of oral corticosteroids and they had FEV₁ values ranging from 40 to 80% of predicted. Favourable and significant efficacy data obtained with this group of patients were similar to those reported in our cohort.

Another post-marketing study with omalizumab was conducted in Israel. However, the 47 patients included in that

study were notably older than our patients. Although significant improvement was also shown in that cohort, the results were less impressive than ours.³²

In conclusion, after three years of omalizumab treatment a significant reduction in the asthma severity (ACT) could be documented in patients with difficult-to-control asthma, combined with an improved lung function, a reduction in the maintenance dose of ICS. However, our prime finding is the significant reduction in the frequency of asthma exacerbations and emergency room visits and hospitalisations caused by these asthma exacerbations. This finally translates into a positive pharmaco-economic balance, even with this high-cost treatment.

Finally, it has not yet been established clearly for how long omalizumab treatment should be continued to change a difficult-to-control asthmatic patient into a well-controlled patient and to avoid relapses after discontinuation.³³ Long-term studies shall have to be conducted to clarify this issue.

Ethical disclosures

Protection of human subjects. The authors declare that no experiments were performed on humans or animals for this investigation, as it is a post-marketing trial.

Confidentiality of data. The authors declare that they have followed the protocols of their work centre on the publication of patient data and that all the patients included in the study have received sufficient information and have given their informed consent in writing to participate in that study.

Right to privacy and informed consent. The authors have obtained the informed consent of the patients and/or subjects mentioned in the article. The author for correspondence is in possession of this document.

Conflict of interest

The authors have no conflict of interest to declare.

Acknowledgement

Investigators did not receive any specific funding for the study.

References

1. Lotvall J, Akdis CA, Bacharier LB, Bjermer L, Casale TB, Custovic A, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol.* 2011;127:355–60.
2. Neffen H. Consenso Latinoamericano sobre el Asma de difícil control, actualización 2008. *Drugs Today.* 2008;44:1–43.
3. Humbert M, Beasley R, Ayres J, Slavin R, Hebert J, Bousquet J, et al. Benefits of omalizumab as add-on therapy in patients with severe persistent asthma who are inadequately controlled despite best available therapy (GINA 2002 step 4 treatment): INNOVATE. *Allergy.* 2005;60:309–16.
4. Godard P, Chanez P, Siraudin L, Nicoloyannis N, Duru G. Costs of asthma are correlated with severity: a 1-yr prospective study. *Eur Respir J.* 2002;19:61–7.
5. Becerril-Ángeles M, Pérez-Chavira R, Martínez-Aguilar N. Treatment of persistent and difficult-to-control asthma: updating of anti IgE therapy. *Rev Alergia Mex.* 2011;58 Suppl. 1:1–18.
6. Proceedings of the ATS workshop on refractory asthma: current understanding, recommendations, and unanswered questions American Thoracic Society. *Am J Respir Crit Care Med.* 2000;162:2341–51.
7. Navarro Merino M, Martín AA, Asensio de la Cruz O. Guía de diagnóstico y tratamiento del asma de control difícil en el niño. *An Pediatr (Barc).* 2009;71:548–67.
8. Lanier B, Bridges T, Kulus M, Taylor AF, Berhane I, Vidaurre CF. Omalizumab for the treatment of exacerbations in children with inadequately controlled allergic (IgE-mediated) asthma. *J Allergy Clin Immunol.* 2009;124:1210–6.
9. Turner S, Paton J, Higgins B, Douglas G. British Guidelines on the Management of A. British guidelines on the management of asthma: what's new for 2011? *Thorax.* 2011;66:1104–5.
10. British Guideline on the Management of Asthma. *Thorax.* 2008;63 Suppl. 4:iv1–121.
11. www.gemasma.com/presentacion/index.asp,. Ultimo accessed 13.08.09.
12. Hurst JW, Morris DC, Alexander RW. The use of the New York Heart Association's classification of cardiovascular disease as part of the patient's complete Problem List. *Clin Cardiol.* 1999;22:385–90.
13. Thomas DM, Navarro-Barrientos JE, Rivera DE, Heymsfield SB, Bredlau C, Redman LM, et al. Dynamic energy-balance model predicting gestational weight gain. *Am J Clin Nutr.* 2012;95:115–22.
14. Korn S, Thielen A, Seyfried S, Taube C, Kornmann O, Buhl R. Omalizumab in patients with severe persistent allergic asthma in a real-life setting in Germany. *Respir Med.* 2009;103: 1725–31.
15. Klyucheva M, von Berg A, Gappa M, Suerbaum C, Berdel D. Omalizumab therapy in adolescents with severe allergic asthma – results of a post-marketing surveillance. *Pneumologie.* 2013;67:233–7.
16. Lafeuille MH, Dean J, Zhang J, Duh MS, Gorsh B, Lefebvre P. Impact of omalizumab on emergency-department visits, hospitalizations, and corticosteroid use among patients with uncontrolled asthma. *Ann Allergy Asthma Immunol.* 2012;109:59–64.
17. To T, Shen S, Atenafu EG, Guan J, McLimont S, Stocks B, et al. The air quality health index and asthma morbidity: a population-based study. *Environ Health Perspect.* 2013;121:46–52.
18. Perry TT, Rettiganti M, Brown RH, Nick TG, Jones SM. Uncontrolled asthma and factors related to morbidity in an impoverished, rural environment. *Ann Allergy Asthma Immunol.* 2012;108:254–9.
19. Dewey A, Dean T, Bara A, Lasserson TJ, Walters EH. Colchicine as an oral corticosteroid sparing agent for asthma. *Cochrane Database Syst Rev.* 2003;CD730032.
20. Dean T, Dewey A, Bara A, Lasserson TJ, Walters EH. Chloroquine as a steroid sparing agent for asthma. *Cochrane Database Syst Rev.* 2003;CD003275.
21. Richeldi L, Ferrara G, Fabbri LM, Lasserson TJ, Gibson PG. Macrolides for chronic asthma. *Cochrane Database Syst Rev.* 2005;CD970029.
22. Evans DJ, Cullinan P, Geddes DM. Cyclosporin as an oral corticosteroid sparing agent in stable asthma. *Cochrane Database Syst Rev.* 2001;CD930029.
23. Evans DJ, Cullinan P, Geddes DM. Gold as an oral corticosteroid sparing agent in stable asthma. *Cochrane Database Syst Rev.* 2001;CD850029.

24. Davies H, Olson L, Gibson P. Methotrexate as a steroid sparing agent for asthma in adults. *Cochrane Database Syst Rev*. 2000;CD000391.
25. Babu KS, Davies DE, Holgate ST. Role of tumor necrosis factor alpha in asthma. *Immunol Allergy Clin North Am*. 2004;24:583–97, v–vi.
26. Rubio-Padilla M, del Rio-Navarro BE, Segura NH, Sienra-Monge JJ. Difficult-to-control asthma. A bibliographical review. *Rev Alerg Mex*. 2009;56:115–23.
27. Poole JA, Meng J, Reff M, Spellman MC, Rosenwasser LJ. Anti-CD23 monoclonal antibody, lumiliximab, inhibited allergen-induced responses in antigen-presenting cells and T cells from atopic subjects. *J Allergy Clin Immunol*. 2005;116:780–8.
28. Hart TK, Blackburn MN, Brigham-Burke M, Dede K, Al-Mahdi N, Zia-Amirhosseini P, et al. Preclinical efficacy and safety of pascolizumab (SB 240683): a humanized anti-interleukin-4 antibody with therapeutic potential in asthma. *Clin Exp Immunol*. 2002;130:93–100.
29. Velarde-Dominguez TS, Martinez SENP, Pantoja-Alc Ntar JM. Acute urticaria associated to omalizumab treatment in a child with asthma. *Rev Alerg Mex*. 2012;59:209–13.
30. Simons FE, Arduoso LR, Bilo MB, Dimov V, Ebisawa M, El-Gamal YM, et al. 2012 Update: World Allergy Organization Guidelines for the assessment and management of anaphylaxis. *Curr Opin Allergy Clin Immunol*. 2012;12:389–99.
31. Shankar T, Petrov AA. Omalizumab and hypersensitivity reactions. *Curr Opin Allergy Clin Immunol*. 2013;13:19–24.
32. Shitrit D, Talker O, Metabichek A, Yaakov I. Omalizumab treatment in patients with asthma: summary of Meir Medical Center experience with 47 patients. *Harefuah*. 2012;151:216–9, 54, 53.
33. Nopp A, Johansson SG, Adedoyin J, Ankerst J, Palmqvist M, Oman H. After 6 years with Xolair; a 3-year withdrawal follow-up. *Allergy*. 2010;65:56–60.