



Allergologia et immunopathologia

Sociedad Española de Inmunología Clínica,
Alergología y Asma Pediátrica

www.elsevier.es/ai



ORIGINAL ARTICLE

Exposure to dogs but not cats is associated to a decrease in the prevalence in atopic dermatitis amongst school-children



M. Bedolla-Barajas^{a,*}, J. Morales-Romero^b, T.I. Bedolla-Pulido^c, T.R. Bedolla-Pulido^a,
C. Meza-López^d, N.A. Pulido-Guillén^e

^a Servicio de Alergia e Inmunología Clínica, Hospital Civil de Guadalajara "Dr. Juan I. Menchaca", Salvador de Quevedo y Zubieta No. 750, Colonia La Perla, C.P. 44340, Guadalajara, Jalisco, Mexico

^b Instituto de Salud Pública, Universidad Veracruzana, Av. Luis Castelazo Ayala s/n., Colonia Industrial Ánimas, C.P. 91190, Xalapa, Veracruz, Mexico

^c Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara. Sierra Mojada No. 950, Colonia Independencia Oriente, C.P. 44340, Guadalajara, Jalisco, Mexico

^d División de Pediatría, Hospital Civil de Guadalajara "Dr. Juan I. Menchaca", Salvador de Quevedo y Zubieta No. 750, Colonia La Perla, C.P. 44340, Guadalajara, Jalisco, Mexico

^e Psicología Clínica, Eulogio Parra No. 2330-301, Col. Las Américas, C.P. 44650, Guadalajara, Jalisco, Mexico

Received 17 August 2017; accepted 14 September 2017

Available online 15 February 2018

KEYWORDS

Pets;
Atopic dermatitis;
Asthma;
Allergic rhinitis;
Dogs;
Cats

Abstract

Introduction: The association regarding the exposure to pets, especially cats and dogs, and the prevalence of allergic diseases is inconsistent.

Objective: We analyzed the role played by early exposure to dogs or cats in the prevalence of allergic diseases amongst school-aged children.

Method: Through a cross-sectional study, we examined 756 children, aged 6–7; these candidates were selected through cluster sampling. We inquired about the exposure that these children had had to dogs and cats, and whether these pets spent most of their time indoors or outdoors during the first year of the child's life. In order to identify the prevalence of allergic diseases and their symptoms, each child's parent completed the *International Study of Asthma and Allergies in Childhood* questionnaire.

Results: Exposure to outdoor dogs was associated to nocturnal coughing, odds ratio (OR) 0.64, with a confidence interval of 95% (95% CI) 0.43–0.95 and with atopic dermatitis (OR: 0.39; 95% CI: 0.20–0.76). Interestingly, exposure to outdoor cats was associated to nocturnal coughing (OR: 0.51; 95% CI: 0.32–0.83) and current rhinitis symptoms (OR: 0.59; 95% CI 0.36–0.97). After carrying out the multivariate analyses, only exposure to dogs, both indoor and outdoor, was significantly associated to a decrease in the prevalence of atopic dermatitis OR 0.40 (95% CI: 0.20–0.79) and OR 0.38 (95% CI: 0.18–0.83), respectively.

* Corresponding author.

E-mail address: drmbedbar@gmail.com (M. Bedolla-Barajas).

Conclusion: Our findings suggest that exposure to dogs, whether they be indoor or outdoor pets, is associated to a decreased prevalence in atopic dermatitis.

© 2018 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

Introduction

On a global level, there is controversy regarding the role that interaction with dogs or cats plays in the development of allergic diseases. Through epidemiological studies have proven that pets are a risk factor when it comes to the development of asthma, allergic rhinitis and atopic dermatitis^{1,2}; on the other hand, there is also evidence that has demonstrated that pets actually act as protective agents³⁻⁵; moreover, there are additional studies that have found pets to be irrelevant factors in the matter.^{6,7}

There is also discrepancy concerning the type of pet that could be associated to the development of these allergic diseases, since there are places in the world where farm animals are considered children's pets,⁸ in some cases fish, birds and rodents act as childhood companions^{2,7}; yet, only cats and dogs have been studied consistently. In Mexico, the role that cats and dogs play in the prevalence of allergic diseases has only been studied a handful of times.

In Mexico, the role that the exposure to dogs and cats plays in the development of the prevalence of allergic diseases has rarely been studied.^{2,8-11} In our country, the most common household pets are cats and dogs, notably the interaction with these pets can vary, as some Mexican families tend to allow their pets to live indoors. The study at hand aims to evaluate how early exposure to dogs or cats affects the prevalence of allergic diseases amongst school-aged children, and whether it makes a difference if these pets reside indoors or outdoors.

Methods

Design and sample size

For this cross-sectional study, we included boys and girls, aged 6–7, all of whom attended primary schools in the city of Guadalajara, Mexico. The study universe consisted of 30,234 children from the 2013 to 2014 school year, distributed throughout 705 public and private schools.

Sampling procedure

For a detailed description of the sampling procedure, read above.¹² In brief, children were incorporated into our study through stratified conglomerated sampling, from April to December of 2014, Fig. 1. The city of Guadalajara is divided into seven administrative districts, each of these regions was contemplated as a stratum; through the proportional assignment technique we obtained a sub-sample from each area, we then multiplied the total sample size by the proportion of

registered students in each district. Through random selection, we chose at least one school (conglomerate) within each district (stratum), and when it was necessary, we selected more schools in order to obtain the required sub-sample size.

Questionnaire

Our inquiry form was structured, and it included questions regarding each family's medical history of allergic diseases for both parents, as well as any previous medical diagnoses of atopic diseases for each child. In order to identify children with allergic diseases, we applied *The International Study of Asthma and Allergies in Childhood (ISAAC)* questionnaire to parents of the children. We also asked whether or not each child had been exposed to dogs or cats during the first year of life, and whether this pet had been a predominantly indoor or outdoor pet.

Ethics

This study was approved by the Ethics and Research Committees at The Hospital Civil de Guadalajara Dr. Juan I. Menchaca; furthermore, this study received support from the Secretaría de Educación del Estado de Jalisco. Most importantly, each parent or guardian, along with his or her child, signed a written consent form in order to participate.

Analysis

The prevalence of allergic diseases and their symptoms was estimated by dividing the number of subjects that answered each question affirmatively in The International Study of Asthma and Allergies in Childhood questionnaire, by the total number of subjects in the study group, additionally, we estimated confidence intervals for proportions. In order to identify a possible association between allergic diseases and the exposure to dogs and cats, we calculated the odds ratio (OR) and their respective Confidence Intervals to measure the magnitude of this association. We employed logistic binary regression to estimate the adjusted OR (OR_a), in which the dependent variable was atopic dermatitis and the independent covariates were: exposure to dogs (indoor or outdoor), sex, family history of atopy, personal history of atopy, a cesarean birth, and breastfeeding. The values $p \leq 0.05$ served as a means to establish statistical significance. Analysis of the data was performed by using the IBM SPSS program, version 20.0 for Windows (Armonk, NY, USA).




Figure 1 Selection of study subjects and distribution of the groups to be compared.

Table 1 Characteristics of study population.

	Total sample	
	n = 756	%
Gender		
Male	356	47.1
Familiar history of allergic disease	148	19.6
Prevalence of allergic diseases		
Asthma	35	4.6
Allergic rhinitis	61	8.1
Atopic dermatitis	37	4.9
Breastfeeding 4 months	460	60.8
Cesarean delivery	427	56.5
Exposure to outdoor dogs	544	72.0
Exposure to outdoor cats	209	27.6
Exposure to indoor dogs	348	46.0
Exposure to indoor cats	81	10.7

Results

We included 756 children in our study sample, of whom, 356 were male (47.1%), Table 1. The prevalence for asthma, allergic rhinitis and atopic dermatitis were: 4.6%, (95% CI: 3.3–6.4), 8.1% (95% CI: 6.3–10.2) and 4.9% (95% CI: 3.5–6.7), respectively. The frequency of previous maternal breastfeeding for at least four months was 61%, for cesarean section deliveries it was 56%. Children interacted with outdoor dogs and cats in 72% and 27.6% of the cases, they were

exposed to indoor pets at a frequency of 46% and 10.7%, respectively.

Children with exposure to outdoor dogs had a significantly decreased prevalence of nocturnal coughing that was unassociated to respiratory infection, one year prior to our study (OR = 0.64; 95% CI: 0.43–0.95, p = 0.028), Table 2. Similarly, this was the case regarding the prevalence of atopic dermatitis (OR = 0.39; 95% CI: 0.20–0.76, p = 0.006). Notably, contact with outdoor cats also presented a statistically significant negative association to nocturnal coughing and rhinitis, both during the year prior to our study, (OR = 0.51; 95% CI: 0.32–0.83, p = 0.006 and OR = 0.59; 95% CI: 0.36–0.97, p = 0.037, respectively).

When we analyzed exposure to indoor pets, contact with dogs was also significantly associated to a decreased prevalence of atopic dermatitis (OR = 0.36; 95% CI: 0.17–0.77, p = 0.009), Table 3. However, there was no observed association of this occurrence with indoor cats.

Multivariate analyses showed that atopic dermatitis was significantly linked to contact with outdoor dogs (ORa = 0.40; 95% CI: 0.20–0.79, p = 0.008) and to a personal history of atopy (ORa = 4.84; 95% CI: 2.35–9.99, p < 0.001), Table 4. This behavior was constant when atopic dermatitis was associated to contact with indoor dogs and a personal history of atopy (ORa = 0.38, p = 0.015 and OR = 4.67, p < 0.0001), Table 5.

Discussion

This study shows that exposure to dogs, but not cats, during the first year of life is associated with a decreased prevalence of atopic dermatitis in children aged 6–7, this was more notable when the dog remained indoors. It also

Table 2 Association between outdoor cats and dogs to the prevalence of allergic diseases and their symptoms.

	Exposure to outdoor dogs				Exposure to outdoor cats			
	Yes (n=544)	No (n=212)	OR (95% CI)	p	Yes (n=209)	No (n=547)	OR (95% CI)	p
<i>Wheezing, ever</i>	85 (15.6)	32 (15.1)	1.04 (0.67–1.62)	0.856	31 (14.8)	86 (15.7)	0.93 (0.59–1.45)	0.762
<i>Prevalence in last year</i>								
Wheezing	31 (5.7)	16 (7.5)	0.74 (0.39–1.38)	0.345	13 (6.2)	34 (6.2)	1.00 (0.51–1.94)	0.998
Exercise wheezing	13 (2.4)	7 (3.3)	0.71 (0.28–1.82)	0.484	4 (1.9)	16 (2.9)	0.65 (0.21–1.96)	0.442
Nocturnal coughing	86 (15.8)	48 (22.6)	0.64 (0.43–0.95)	0.028	24 (11.5)	110 (20.1)	0.51 (0.32–0.83)	0.006
<i>Asthma prevalence</i>	23 (4.2)	12 (5.7)	0.74 (0.36–1.51)	0.401	11 (5.3)	24 (4.4)	1.21 (0.58–2.52)	0.609
<i>Rhinitis, ever</i>	104 (19.1)	54 (25.5)	0.69 (0.47–1.01)	0.054	36 (17.2)	122 (22.3)	0.72 (0.48–1.09)	0.126
<i>Prevalence in last year</i>								
Rhinitis	74 (13.6)	39 (18.4)	0.69 (0.45–1.09)	0.098	22 (10.5)	91 (16.6)	0.59 (0.36–0.97)	0.037
Rhinoconjunctivitis	41 (7.5)	12 (5.7)	1.36 (0.69–2.64)	0.365	12 (5.7)	41 (7.5)	0.75 (0.39–1.46)	0.399
<i>Allergic rhinitis prevalence</i>	41 (7.5)	20 (9.4)	0.78 (0.45–1.37)	0.390	16 (7.7)	45 (8.2)	0.92 (0.51–1.67)	0.796
<i>Rash, ever</i>	33 (6.1)	13 (6.1)	0.98 (0.51–1.92)	0.972	10 (4.8)	36 (6.6)	0.71 (0.35–1.46)	0.357
<i>Prevalence in last year</i>								
Itchy rash	24 (4.4)	10 (4.7)	0.93 (0.44–1.98)	0.855	8 (3.8)	26 (4.8)	0.79 (0.35–1.79)	0.583
Flexural rash	19 (3.5)	6 (2.8)	1.24 (0.49–3.15)	0.647	8 (3.8)	17 (3.1)	0.43 (0.18–1.01)	0.053
Rash cleared	13 (2.4)	8 (3.8)	0.62 (0.25–1.53)	0.302	5 (2.4)	16 (2.9)	0.81 (0.29–2.25)	0.691
<i>Atopic dermatitis prevalence</i>	19 (3.5)	18 (8.5)	0.39 (0.20–0.76)	0.006	7 (3.3)	30 (5.5)	0.59 (0.26–1.38)	0.228

Values in parentheses represent %.

OR: odds ratio; CI: confidence interval.

Table 3 Association between indoor cats and dogs to the prevalence of allergic diseases and their symptoms.

	Exposure to indoor dogs				Exposure to indoor cats			
	Yes (n=348)	No (n=408)	OR (95% CI)	p	Yes (n=81)	No (n=675)	OR (95% CI)	p
<i>Wheezing, ever</i>	60 (17.2)	57 (14.0)	1.28 (0.86–1.90)	0.216	16 (19.8)	101 (15.0)	1.65 (0.91–2.99)	0.098
<i>Prevalence in last year</i>								
Wheezing	19 (5.5)	28 (6.9)	0.78 (0.42–1.43)	0.427	5 (6.2)	42 (6.2)	0.99 (0.38–2.58)	0.986
Exercise wheezing	6 (1.7)	14 (3.4)	0.49 (0.18–1.28)	0.146	0 (0)	20 (3.0)	0.65 (0.21–1.96)	0.442
Nocturnal coughing	59 (17.0)	75 (18.4)	0.90 (0.62–1.32)	0.608	12 (14.8)	122 (18.1)	0.79 (0.41–1.50)	0.469
<i>Asthma prevalence</i>	14 (4.0)	21 (5.1)	0.77 (0.39–1.54)	0.465	4 (4.9)	31 (4.6)	1.08 (0.37–3.14)	0.889
<i>Rhinitis, ever</i>	73 (21.0)	85 (20.8)	1.01 (0.71–1.43)	0.961	20 (24.7)	138 (20.4)	1.27 (0.74–2.18)	0.375
<i>Prevalence in last year</i>								
Rhinitis	51 (14.7)	62 (15.2)	0.96 (0.64–1.43)	0.835	12 (14.8)	101 (15.0)	0.99 (0.52–1.89)	0.971
Rhinoconjunctivitis	26 (7.5)	27 (6.6)	1.17 (0.67–2.06)	0.569	7 (8.6)	46 (6.8)	1.29 (0.56–2.97)	0.544
<i>Allergic rhinitis prevalence</i>	24 (6.9)	37 (9.1)	0.74 (0.43–1.27)	0.276	7 (8.6)	64 (8.0)	0.90 (0.39–2.04)	0.807
<i>Rash, ever</i>	16 (4.6)	30 (7.4)	0.61 (0.32–1.13)	0.117	3 (3.7)	43 (6.4)	0.71 (0.35–1.46)	0.357
<i>Prevalence in last year</i>								
Itchy rash	10 (2.9)	24 (5.9)	0.47 (0.22–1.00)	0.051	3 (3.7)	31 (4.6)	0.79 (0.24–2.67)	0.716
Flexural rash	9 (2.6)	16 (3.9)	0.65 (0.28–1.49)	0.309	2 (2.5)	23 (3.4)	0.72 (0.17–3.10)	0.657
Rash cleared	5 (1.4)	16 (3.9)	0.36 (0.13–0.98)	0.302	3 (3.7)	18 (2.7)	1.40 (0.40–4.87)	0.593
<i>Atopic dermatitis prevalence</i>	9 (2.6)	28 (6.9)	0.36 (0.17–0.77)	0.009	3 (3.7)	34 (5.0)	0.72 (0.22–2.41)	0.601

Values in parentheses represent %.

OR: odds ratio; CI: confidence interval.

demonstrates that a family history of atopy is a factor that is significantly associated to the prevalence of atopic dermatitis. To our knowledge, thus far, no other studies have evaluated whether the child interacts with indoor or outdoor pets and how this affects allergic diseases.

The attempt to quantify the association between pets and allergic diseases has produced contradictory findings, furthermore, there is no consistency regarding the types of pets that might be associated to such diseases. For example, one study carried out in Iran, factored in farm animals such

as cows, pigs, goats and sheep as pets: in their study the children presented a reduced frequency of wheezing fits, coughing, sneezing, rhinorrhea or nasal obstruction and epidermal itchiness during the year prior to their study,⁸ and while those animals lived with the children.

In our country, given the structural characteristics of our homes, the majority of cats and dogs, tend to live outdoors, that is to say, they live in open garages, patios or roof tops, yet, they can also reside indoors, sometimes they even sleep inside their owners' bedrooms. Thus, the

Table 4 Association between exposure to an outdoor dog during the first year of life and the prevalence of atopic dermatitis.

	Multivariate analysis					
	Unadjusted model			Adjusted model		
	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
Outdoor dog	0.42	0.21–0.83	0.012	0.40	0.20–0.79	0.008
Sex, male	1.31	0.66–2.61	0.440	–	–	0.457
Personal history of allergic diseases	1.38	0.63–2.99	0.422	–	–	0.421
Familiar history of allergic disease	4.59	2.11–9.96	<0.0001	4.84	2.34–9.99	<0.001
Cesarean delivery	0.72	0.36–1.43	0.348	–	–	0.365
Breastfeeding 4 months	1.06	0.53–2.13	0.876	–	–	0.889

OR: odds ratio; CI: confidence interval;

OR obtained by logistic regression.

OR for variables that were used to adjust the model were not contemplated.

Table 5 Association between exposure to an indoor dog during the first year of life and the prevalence of atopic dermatitis.

	Multivariate analysis					
	Unadjusted model			Adjusted model		
	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
Indoor dog	0.39	0.18–0.87	0.021	0.38	0.18–0.83	0.015
Sex, male	1.31	0.66–2.61	0.442	–	–	0.452
Personal history of allergic diseases	1.41	0.65–3.05	0.385	–	–	0.386
Familiar history of allergic disease	4.44	2.05–9.62	<0.0001	4.67	2.26–9.65	<0.0001
Cesarean delivery	0.72	0.36–1.44	0.353	–	–	0.369
Breastfeeding 4 mo.	1.03	0.51–2.06	0.943	–	–	0.947

OR: odds ratio; CI: confidence interval.

OR obtained by logistic regression.

OR for variables that were used to adjust the model were not contemplated.

level of exposure to pets is a variable factor, such variances can directly influence the development of allergic diseases.

In regard to asthma, a recent study showed that exposure to dogs or pets (cattle or sheep), during the first year of life, was associated to a reduced rate of asthma incidences in pre-school to school-aged children.⁴ After analyzing the data obtained in exclusively European cohort studies (11 studies), there was no documented association between exposure to dogs or cats and asthma amongst school-aged children.⁷ In China, a study that included more than 13 thousand children, aged 4–6, demonstrated that interaction with pets acted as a risk factor for asthma prevalence.² In our study, exposure to dogs and cats was associated to a reduced frequency in coughing, during the 12 months prior to our study, however, this was only the case when pets remained outside the home, nevertheless, this effect disappeared when the interaction with dogs or cats was indoor. The rest of the asthma symptoms, or those previously associated to asthma, were no longer documented as a factor.

When it comes to allergic rhinitis, and its symptoms, our analysis of exposure to outdoor cats showed that these pets acted as protective agents against the presence of rhinitis during the year prior to our study; however, this agent disappeared when the family cat lived indoors. Phase III of the

ISAAC study, which included data from various international research centers, provided contradicting results when it came to the association between dogs or cats and the symptoms of allergic diseases; this was the case for children aged 6–7 and adolescents, allergic rhinitis and allergic rhinoconjunctivitis appeared to be more prevalent as a result of exposure to cats during the first year of life; however, dogs did not have this effect on the adolescent group.¹ In Latin America, the rhinoconjunctivitis symptoms amongst Bolivian schoolchildren, did not increase in relation to their exposure to dogs or cats, but it did rise with exposure to farm animals.⁶ On the other hand, Chinese children that were exposed to dogs during birth were at greater risk of being diagnosed with allergic rhinitis, when compared to children that were not exposed to dogs during birth.² Recently, an independent cohort study observed that indoor exposure to dogs was strongly associated to a decreased risk of developing atopic dermatitis at the age of three, notably, this decrease was dose-dependent.⁵ In a meta-analysis that only included cohort studies, a 25% decreased risk of atopic dermatitis was observed in children that were exposed to dogs; contrastingly, cats offered no observable protection.³ In our study, which is not a cohort study, the findings were similar to those in the meta-analysis. We observed a 60% decrease

in the prevalence of atopic dermatitis when children maintained contact with a dog during the first year of their lives, regardless of whether the dog lived indoors or outdoors.

In households where fish are favored as indoor pets, these appear to act as a risk factor for asthma, allergic rhinitis or allergic dermatitis symptoms; however, these symptoms varied based on the stage of life during which the child was exposed to fish.² Birds or rodents are other pets that families choose to adopt; interestingly, according to the meta-analysis, exposure to these animals appear to neither increase nor decrease the risk of developing asthma or allergic rhinitis amongst schoolchildren.⁷ Our study did not encompass exposure to pets such as fish, birds or rodents and their possible association to allergic diseases, thus, we are unaware of how these animals might affect our population.

Various explanations have been proposed when exposure to dogs or cats has acted as a protective agent against the development of allergic diseases; of these explanations, the hygiene hypothesis is one of the most well-known, it is said that fewer family members, better domestic services, and hygiene, have decreased the number of contagions, consequently, the expression of atopic diseases has increased¹³; nevertheless, this hypothesis seems to leave out the exposure to animals that have fur, which is why this would not be the most comprehensive justification when it comes to the roles that pets might play in the prevalence of allergic diseases. Furthermore, it has been observed that children who live in rural areas or farms, have a lower chance of manifesting allergic diseases, when compared to children that live in an urban setting.^{14–17} In our country, it appears that living in a rural area or being exposed to farm animals does not offer any allergic protection.¹⁸ When researchers have attempted to explain a rural environment as a protective mechanism, endotoxins are usually credited for this phenomenon. In fact, there is evidence that has proven that exposure to endotoxin reduces the risk of atopic dermatitis.¹⁹ Interestingly, the presence of dogs, but not cats, has been associated to an increased variety of microbiome bacteria where these animals live²⁰; however, researchers have also detected a higher amount of endotoxin bacteria in household dust;²¹ thus, early exposure to dogs might influence the child's immune response, and consequently, reduce the frequency of atopic diseases.

An area that has seldom been explored concerns the role that the attachment theory plays in the development of allergic diseases. According to Bowlby, attachment is a child's willingness to seek proximity and contact with an individual²²; from this perspective, children are likely to form attachments to their pets, especially to a dog, and when this companionship is established during the first years of life; thus, the coexistence with a pet would allow a child to obtain psychological and social benefits.²³ Similarly, an overall sensation of wellbeing might be attained by having a pet, this could stimulate the production of oxytocin, a neuropeptide that has been associated to social behavior in human beings.²⁴ Recently, it has been proven that oxytocin is not only produced in the pituitary gland, but that it can also be produced and released through keratinocyte and fibroblasts in the skin; which is why the lack of expression from the oxytocin receptor, on the keratinocyte surface, has been linked to the release of IL-6, CCL5 and CXCL10; all of these cytokines have pro-inflammatory activity.²⁵ Thus,

close contact with dogs, especially during the early stages of life, regulates an immunological response in the skin through oxytocin, resulting in a reduced prevalence of atopic dermatitis.

One of the factors that has been consistently associated to allergic diseases regards the genetic predisposition of an individual.^{9,26–28} In our study, a family history of atopy was notably associated to the prevalence of atopic dermatitis. Notably, in our population, factors such as breastfeeding or a cesarean delivery, do not seem to be associated to allergic diseases.^{27,29}

In our study there were additional limitations to those that are inherent to cross sectional studies. It is necessary to make note of the fact that we did not analyze the number of dogs in each household; as previously noted, when there are more dogs, there is a reduced likelihood of atopic dermatitis. As far as each family's dogs, we are unaware of what their characteristics were; it is likely that a dog's size, fur, or sex could influence the development of allergic diseases. Furthermore, we did not inquire about the amount of time that each child might have spent directly with their dogs or cats, or whether this contact had been established during gestation, nor did we ask if they had maintained contact with a pet until the time of our investigation. Likewise, we did not ask about exposure to farm animals, as we assumed that children in an urban setting are unlikely to be in contact with them. Conversely, one of this study's strengths lies in the fact that we did give importance to where these pets spent most of their time; the latter is a relevant matter when studying how pets can affect the development of allergic diseases. An additional strength in our study regards the sampling procedure, which allowed us to incorporate children from all socioeconomic strata, different school schedules and every geographic area within the city of Guadalajara.

Conclusions

In conclusion, this study proves that exposure to dogs during the first year of life, whether they are indoor or outdoor pets, is associated to a reduced prevalence of atopic dermatitis in schoolchildren.

To date, there is no uniformity in the results that have been obtained from the various approaches to the research projects that have evaluated the association between pets and allergic diseases. Moreover, there is no regularity when it comes to the types of pets that might be acting as protective agents against these diseases.

Lastly, it is important to note that a personal history of the allergic disease favored its manifestation.

Funding

None.

Conflict of interest

The authors have no conflict of interest to declare.

Acknowledgements

As authors of this study, we wish to express our gratitude to the parents and children that kindly chose to partake in our Project; as well as the teachers at all participating schools.

We would also like to thank Pharmacist Sandra Vega Villamar for her help with the design of the flowchart.

References

1. Brunekreef B, Von Mutius E, Wong G, Odhiambo J, García-Marcos L, Foliaki S, ISAAC Phase Three Study Group. Exposure to cats and dogs, and symptoms of asthma, rhinoconjunctivitis, and eczema. *Epidemiology*. 2012;23:742–50.
2. Huang C, Hu Y, Liu W, Sundell Je. Pet-keeping and its impact on asthma and allergies among preschool children in Shanghai, China. *Chin Sci Bull*. 2013;58:4203–10, <http://dx.doi.org/10.1007/s11434-013-5679-4>.
3. Pelucchi C, Galeone C, Bach JF, La Vecchia C, Chatenoud L. Pet exposure and risk of atopic dermatitis at the pediatric age: a meta-analysis of birth cohort studies. *J Allergy Clin Immunol*. 2013;132:616–22.
4. Fall T, Lundholm C, Örtqvist AK, Fall K, Fang F, Hedhammar Å, et al. Early exposure to dogs and farm animals and the risk of childhood asthma. *JAMA Pediatr*. 2015;169:e153219.
5. Thorsteinsdóttir S, Thyssen JP, Stokholm J, Vissing NH, Waage J, Bisgaard H. Domestic dog exposure at birth reduces the incidence of atopic dermatitis. *Allergy*. 2016;71:1736–44.
6. Solis-Soto MT, Patiño A, Nowak D, Radon K. Association between environmental factors and current asthma, rhinoconjunctivitis and eczema symptoms in school-aged children from Oropeza Province-Bolivia: a cross-sectional study. *Environ Health*. 2013;12:95.
7. Lødrup Carlsen KC, Roll S, Carlsen KH, Mowinckel P, Wijga AH, Brunekreef B, et al., G ALEN WP 1.5 'Birth Cohorts' working group. Does pet ownership in infancy lead to asthma or allergy at school age? Pooled analysis of individual participant data from 11 European birth cohorts. *PLoS ONE*. 2012;7:e43214.
8. Karimi M, Mirzaei M, Baghiani Moghadam B, Fotouhi E, Zare Mehrjardi A. Pet exposure and the symptoms of asthma, allergic rhinitis and eczema in 6-7 years old children. *Iran J Allergy Asthma Immunol*. 2011;10:123–7.
9. Dong GH, Ding HL, Ma YN, Jin J, Cao Y, Zhao YD, et al. Asthma and asthma-related symptoms in 16,789 Chinese children in relation to pet keeping and parental atopy. *J Investig Allergol Clin Immunol*. 2008;18:207–13.
10. Dong GH, Ma YN, Ding HL, Jin J, Cao Y, Zhao YD, et al. Pets keeping in home, parental atopy, asthma, and asthma-related symptoms in 12,910 elementary school children from northeast China. *Indoor Air*. 2009;19:166–73.
11. Lombardi E, Simoni M, La Grutta S, Viegi G, Bisanti L, Chellini E, et al., SIDRIA-2 Collaborative Group. Effects of pet exposure in the first year of life on respiratory and allergic symptoms in 7-yr-old children. The SIDRIA-2 study. *Pediatr Allergy Immunol*. 2010;21:268–76.
12. Bedolla-Barajas M, Valdez-López F, Alcalá-Padilla G, Bedolla-Pulido TI, Rivera-Mejía V, Morales-Romero J. Prevalence and factors associated to peanut allergy in Mexican school children. *Allergol Immunopathol (Madr)*. 2017;45:69–76.
13. Strachan DP. Hay fever, hygiene, and household size. *BMJ*. 1989;299:1259–60.
14. Solé D, Cassol VE, Silva AR, Teche SP, Rizzato TM, Bandim LC, et al. Prevalence of symptoms of asthma, rhinitis, and atopic eczema among adolescents living in urban and rural areas in different regions of Brazil. *Allergol Immunopathol (Madr)*. 2007;35:248–53.
15. Bäcker C, Barraza-Villarreal A, Moreno-Macías H, Escamilla-Núñez C, Romieu I. The effects of a rural environment on the prevalence of allergic rhinitis among schoolchildren in Mexicali, Baja California, Mexico. *Rev Panam Salud Publica*. 2009;25:431–7 [in Spanish].
16. Kausel L, Boneberger A, Calvo M, Radon K. Childhood asthma and allergies in urban, semiurban, and rural residential sectors in Chile. *ScientificWorldJournal*. 2013;2013:937935.
17. Horak E, Morass B, Ulmer H, Genuneit J, Braun-Fahrlander C, von Mutius E, GABRIEL Study Group. Prevalence of wheezing and atopic diseases in Austrian schoolchildren in conjunction with urban, rural or farm residence. *Wien Klin Wochenschr*. 2014;126:532–6.
18. Bedolla-Barajas M, Ramírez-Cervantes FJ, Morales-Romero J, Pérez-Molina JJ, Meza-López C, Delgado-Figueroa N. A rural environment does not protect against asthma or other allergic diseases amongst Mexican children. *Allergol Immunopathol (Madr)*. 2018;46:31–8.
19. Gehring U, Bolte G, Borte M, Bischof W, Fahlbusch B, Wichmann HE, et al., LISA study group. Lifestyle-related factors on the immune system and the development of allergies in childhood. Exposure to endotoxin decreases the risk of atopic eczema in infancy: a cohort study. *J Allergy Clin Immunol*. 2001;108:847–54.
20. Kettleson EM, Adhikari A, Vesper S, Coombs K, Indugula R, Reponen T. Key determinants of the fungal and bacterial microbiomes in homes. *Environ Res*. 2015;138:130–5.
21. Heinrich J, Gehring U, Douwes J, Koch A, Fahlbusch B, Bischof W, et al., INGA-Study Group. Pets and vermin are associated with high endotoxin levels in house dust. *Clin Exp Allergy*. 2001;31:1839–45.
22. Repeteur-Safrany K, Quezada-Len A. Vínculo y desarrollo psicológico: la importancia de las relaciones tempranas. *Revista Digital Universitaria*. 2005;6:1–15.
23. Rodríguez-Niño MM. Relación entre presencia de mascotas (caninas) co nestilos de apego en niños entre 6 a 13 años de edad procedentes de Bogotá, Colombia. Available from: [Internet] http://www.academia.edu/15980641/TEORIA.DEL.APEGO_NINO_MASCOTA [accessed 19.2.17].
24. Carter CS. Oxytocin pathways and the evolution of human behavior. *Annu Rev Psychol*. 2014;65:17–39.
25. Deing V, Roggenkamp D, Kühnl J, Gruschka A, Stäb F, Wenck H, et al. Oxytocin modulates proliferation and stress responses of human skin cells: implications for atopic dermatitis. *Exp Dermatol*. 2013;22:399–405.
26. Baek JO, Hong S, Son DK, Lee JR, Roh JY, Kwon HJ. Analysis of the prevalence of and risk factors for atopic dermatitis using an ISAAC questionnaire in 8,750 Korean children. *Int Arch Allergy Immunol*. 2013;162:79–85.
27. Pohlabein H, Jacobs S, Böhm J. Exposure to pets and the risk of allergic symptoms during the first 2 years of life. *J Investig Allergol Clin Immunol*. 2007;17:302–8.
28. Morales-Romero J, Bedolla-Barajas M, López-Vargas L, Romero-Velarde E. Prevalence of allergic diseases and their association with breastfeeding and initiation of complementary feeding in school-age children of Ciudad Guzmán, Mexico. *Arch Argent Pediatr*. 2015;113:324–30.
29. Bedolla-Barajas M, Barrera-Zepeda AT, López-Zaldo JB, Morales-Romero J. Asthma in Mexican school-age children is not associated with passive smoking or obesity. *Asia Pac Allergy*. 2013;3:42–9.