



ORIGINAL ARTICLE

Multispecies probiotics combination prevents ovalbumin-induced airway hyperreactivity in mice

Jui-Cheng Chen^a, Cheng-Chih Tsai^a, Chang-Chi Hsieh^b, Alice Lan^c, Chun-Chih Huang^c, Sew-Fen Leu^{a,*}

^a Department of Food Science and Technology, HungKuang University, Taichung, Taiwan

^b Department of Animal Science and Biotechnology, Tunghai University, Taichung, Taiwan

^c New Bellus Enterprises Co., Ltd., 48 Industrial Road, Tainan City 72042, Taiwan

Received 5 September 2017; accepted 22 February 2018

Available online 5 May 2018

KEYWORDS

Lactobacillus plantarum;
Pediococcus acidilactici;
Anti-allergy;
Airway hyperreactivity

Abstract

Background: Probiotics could be beneficial to health and some of them have shown to modulate immune responses.

Aim: The aim of this study is to investigate if the probiotic strains including *Lactobacillus* and *Pediococcus* strains are able to alleviate allergic reactions in an ovalbumin-induced airway allergy model.

Methods: *Lactobacillus* multi-species preparation (LMP) was gavaged to BALB/c for total six weeks and BALB/c was challenged with ovalbumin in the last two weeks. A barometric whole-body plethysmography was used to assess enhanced pause (Penh) of airway hyperreactivity (AHR). Immunoglobulins (Ig) such as IgE, IgG1, IgG2a and cytokines such as IL-12, IFN- γ , IL-4, IL-5, TNF- α and IL-13 in bronchoalveolar lavage fluid were assayed using ELISA kits.

Results: The results showed this LMP significantly reduced Th2 cytokines and enhanced Th1 cytokines production. OVA-specific IgE and IgG1 was lower in the probiotics-treated mice whereas IgG2a was increased. Most importantly, this murine model showed LMP supplementation significantly reduced AHR.

Conclusions: Overall, this *Lactobacillus* multi-species preparation seemed to suppress OVA-sensitized airway hyperreactivity, thus serving as a possible candidate for therapeutic uses for allergic airway symptoms.

© 2018 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: sfleu@hk.edu.tw (S.-F. Leu).

Introduction

Asthma is a severe health issue worldwide. One of the etiologic agents for asthma is the exposure to allergens, which cause excess mucus secretion, airway obstruction, inflammation and airway hyperreactivity (AHR).¹ The pathogenesis of asthma is largely mediated by Th2 responses, which consist of a complex network of cytokines interaction. The Th-2 type cytokines such as interleukin (IL)-4, -5 and -13 are involved in pathogenesis of allergic response and eosinophil infiltration in AHR.²⁻⁴ Antigen presenting cells (APCs) such as dendritic cells recognize allergens such as dusts, viral particles and then activate Th2 response. The activated Th2 cells facilitate production of IgE, IgG1 by B cells, degranulation of mast cells and eosinophilia via cytokine secretion. The ovalbumin-induced acute inflammation on airway mucosa gradually goes to a chronic inflammation and leads to airway remodeling. Thus, dampening excess Th2 responses in asthmatic subjects is considered as a plausible approach in ameliorating AHR. Th1 responses on the other hand are important in regulating Th2 responses. APCs such as dendritic cells secret IL-12 to drive Th1 cell differentiation, which induce IFN- γ and IgG2a production. IFN- γ has both anti-asthmatic and pro-inflammatory effects. It tends to inhibit the proliferation of Th2 cells, stimulate preferential expansion of Th1 cells and promotes macrophage killing in response to intracellular microbes and parasites. TNF- α also has the potential to stimulate induction of cytokine, therefore it could play an important role in the pathogenesis of airway remodeling. As a conclusion from the above, enhancing Th1 responses seems to be effective in counter-acting excess Th2 responses in alleviating AHR.

Probiotics refer as microorganisms that confer health benefits to the host.⁵ It is known from mouse studies that intestinal microflora are necessary to shape our immune system as germ-free mice do not have a well-functioning immune system.⁶ The surface molecules on the bacteria are recognized by APCs and stimulate T cell development. *Bifidobacterium* and *Lactobacillus* probiotics have been shown to alleviate various allergic conditions. For example, *B. bifidum* BGN4 has been shown to alleviate food allergy by inhibiting IgE and IgG production.⁷ *Lactobacillus* spp. have been widely studied in food and airway allergy models.⁸⁻¹¹ *Lactobacillus rhamnosus* GG was shown in lowering Th2 cytokines and IgE production in an ovalbumin-induced air inflammation mouse model.¹¹

Another example, *Lactobacillus casei* shiota cannot only lower the production of Th2 cytokines such as IL-4, IL-5, but also increase production of Th1-associated cytokines including IFN- γ and IL-12 in a food allergy model.¹² Overall, the restoration of Th1 and Th2 balance seems the most sensible approach to alleviate allergic conditions.

In this study, the potential probiotic preparation contains two *Lactobacillus plantarum* strains LP109 and LP110 and *Pediococcus acidilactici* PA320. We screened several probiotic strains which demonstrated the potential to modulate AHR and immunoglobulin production in a mouse model (data not shown) and chose the potential ones for this experiment. A previous study by Nawaz demonstrated a multi-species preparation including *Bifidobacterium breve* M-16V and *L. plantarum* NumRes8 reduced IL-4 and IL-5 production.¹³ To

the best of our knowledge, no study has reported the anti-allergy effect of *P. acidilactici* in the airway allergy model before. The aim of this study is to investigate if the probiotic preparation including *Lactobacillus* and *Pediococcus* strains is able to alleviate allergic reactions in an ovalbumin-induced airway allergy model and whether the anti-allergy effect are exerted by either or both Th1 and Th2 responses.

Materials and methods

Animals

BALB/c mice were purchased from BioLasco, Taiwan Co., Ltd. (Taipei, Taiwan). Six-week old female mice were first acclimatized in an animal room for a week at 20–23 °C with 12-hour light/dark cycles. The animals were provided with Lab Diet® 5010 Rodent Diet (PMI Nutrition International, USA) and sterile reverse osmosis water ad libitum. The study was approved by the Institutional Animal Care and Use Committee (IACUC) in Hungkuang University (Taichung, Taiwan).

LMP preparation

The lactobacillus multi-species preparation (abbreviated as LMP) containing *L. plantarum* LP109, *L. plantarum* LP110 and *P. acidilactici* PA320 was fermented and freeze-dried by New Bellus Enterprises Co. Ltd. (Tainan, Taiwan), the powder (3.5×10^{11} CFU/g) was stored at –20 °C. Daily intake of LMP was 1.8 g or 7.2 g per day for adult. The animal equivalent dose using body surface area was 0.39 g/kg BW (low dose) or 1.56 g/kg BW (high dose).¹⁴ The probiotic powder was suspended in sterile phosphate-buffer saline (PBS) for gavage.

Experimental regime


Forty animals were divided into four groups, include naïve group, OVA control group (OVA challenge, PBS gavage), lactobacillus multi-species preparation (abbreviated as LMP) low-dose group (L.L) (OVA challenge, LMP at 0.39 g/kg BW) and LMP high-dose group (L.H) (OVA challenge, LMP at 1.56 g/kg BW), with ten mice in each group. The LMP was orally gavaged to animals daily for the first four weeks.

After that, animals were intraperitoneal sensitized using OVA (0.5 mg OVA mixed with 2 mg Al(OH)₃, 0.1 ml/10 g BW per mouse) for ten days, then intranasally challenged with OVA (2 mg/ml, 75 μ l per mouse) for five days¹⁵; airway hyperreactivity was then assessed using barometric whole-body plethysmography. After animal sacrifice, bronchoalveolar lavage fluid was collected for further analysis.

The experimental timeline is as follows:

Assessment of airway hyperreactivity by barometric whole-body plethysmography

At the end of the last intranasal challenge, the animals were placed in a chamber for the whole-body plethysmograph (Buxco, Troy, NY, USA) for 3 min, and the pressure differences between the chamber and a reference chamber were measured using a differential pressure transducer connected



to an amplifier for recording. The respiratory cycle of the mouse would cause the change of the box pressure and the signal. The enhanced pause (Penh) reflects changes in the wave form of the box pressure signal during both inspiration and expiration. Penh was calculated as $(Te-Tr)/Tr$ (PEP/PIP), where Te is expiratory time (s); Tr is relaxation time (s), defined as the time of pressure decay to 30% of the total expiratory pressure signal (area under the box pressure signal at expiration); PEP is peak expiratory pressure (ml/s); and PIP is peak inspiratory pressure (ml/s). The Penh values were recorded.¹⁶

Harvest of bronchoalveolar lavage fluid and cell count

The trachea was cannulated after mouse was interperitoneally injected with pentobarbital (1.5 g/kg bw) and the cannula was clamped. One ml of Hank's balanced salt solution (HBSS) was lavaged into the lung and retrieved from the lung after gentle message. The whole process was repeated three times. The retrieved bronchoalveolar lavage fluid (BALF) was centrifuged and stored frozen (-70°C) for further analysis. Cell preparations were stained with Liu's stain method and a differential cell count was performed on 300 cells or more. The cell proportions were determined by cell counting using a hemocytometer under an optical microscope.

Evaluation of immunoglobulins and cytokines in BALF

The OVA-specific antibodies IgE, IgG1 and IgG2a in BALF were assayed using the commercial ELISA kits (Bethyl Laboratories, Inc. Montgomery, TX, USA).¹⁷ Cytokines such as IL-4, IL-5, TNF- α , IL-13, IL-12 and IFN- γ were also assayed using the commercial ELISA kits (eBioscience, San Diego, CA, USA).^{18,19}

Statistical analysis

Means and standard deviations were calculated for the collected data. The data were analyzed by one-way analysis of variance (one-way ANOVA), followed by post hoc Duncan's multiple range testing. A $P < 0.05$ value was considered as the minimum level for significance.




Figure 1 Effect of oral administration of LMP on the airway response measured at different time points after the last OVA challenge as expressed by Penh. The Penh value of mice was measured using a non-invasive whole-body plethysmography at 0, 1, 6, 12, 24 and 48 h after the last OVA challenge. (A) Penh values were expressed as means, $n = 10$. (B) Penh value, each value represents the mean \pm SD, $n = 10$. Groups are normal control (naïve), OVA-challenged control (OVA ctrl), low dose LMP (L.L) and high dose LMP (L.H). A difference between the LMP groups and OVA ctrl group at the same time point was considered statistically significant when $P < 0.05$ (*).

Results

Effect of LMP administration on AHR in OVA-sensitized mice

AHR was measured using a non-invasive whole-body plethysmography. The airway resistance was measured in animals after five days of intranasal OVA-challenge. The measurement was expressed as Penh (Enhanced Pause). The higher the Penh value the higher the airway resistance. In Fig. 1, Penh values for naïve animals over the time course showed minimal fluctuations. At 1, 6, 12 h, the naïve group showed lower Penh values ($P < 0.05$) compared with OVA control




Figure 2 Effect of oral administration of LMP on lung tissue inflammatory cell infiltration in OVA-sensitized mice. Cell preparations were stained with Liu's stain method and a differential cell count was performed using a hemocytometer. Each value represents the mean \pm SD, $n=7$. Groups are normal control (naive), OVA-challenged control (OVA ctrl), low dose LMP (L.L) and high dose LMP (L.H). Mono: monocyte, LYMPH: lymphocyte, NEUT: neutrophil, EOSIN: eosinophil. A difference between the LMP groups and OVA ctrl group was considered statistically significant when $P < 0.05$ (*).

group (Fig. 1A and B). Low dose LMP and high dose LMP group showed lower Penh values at 1, 6, 12 h ($P < 0.05$) compared with the OVA control group (Fig. 1).

Effect of LMP on cellular distribution in BALF of OVA-sensitized mice

Animal showed normal body weight increase and food intake during the experiment period (data not shown). No significant difference ($P > 0.05$) on numbers of monocytes and neutrophils were observed between the OVA control group and both

LMP dose groups (Fig. 2). A lower lymphocyte count ($P < 0.05$) was found in both LMP dose groups compared with the OVA control group. The number of eosinophils was significantly lower ($P < 0.05$) in the low dose LMP-treated group than the OVA control group (Fig. 2).

Effect of LMP on immunoglobulins in BALF of OVA-sensitized mice

In BALF, the high dose LMP-treated group showed lower levels of OVA-specific IgE and IgG1 ($P < 0.05$) compared with the OVA control group (Fig. 3A and B). The OVA-specific IgG2a level was significantly higher in both LMP dose groups ($P < 0.05$) than the OVA control group (Fig. 3C).

Effect of LMP on cytokines in BALF of OVA-sensitized mice

The high dose LMP-treated group showed higher IL-12 and IFN- γ secretion ($P < 0.05$) compared with the OVA control group (Fig. 4A and B). While IL-4 was lower in the high dose LMP-treated group ($P < 0.05$) and IL-5 was lower in both LMP dose groups ($P < 0.05$) (Fig. 5A and B). The level of TNF- α and IL-13 in BALF of both LMP dose groups were lower than the OVA control group ($P < 0.05$) (Fig. 6A and B).




Figure 3 Effect of oral administration of LMP on immunoglobulins in BALF of OVA-sensitized mice. (A) OVA-specific IgE (B) OVA-specific IgG1 and (C) OVA-specific IgG2a in BALF of mice. Each value represents the mean \pm SD, $n=7$. Groups are normal control (naive), OVA-challenged control (OVA ctrl), low dose LMP (L.L) and high dose LMP (L.H). A difference between the LMP groups and OVA ctrl group was considered statistically significant when $P < 0.05$ (*).

Discussion

The BALB/c strain has been commonly used for allergy model.²⁰ Upon OVA sensitization, these animals displayed airway hyperreactivity, increased IgE levels and elevated numbers of eosinophils in the BALF. Th2 cytokines such as IL-4, IL-5, IL-13 are known for promoting allergic responses including eosinophil infiltration, airway hyperreactivity, IgE production and airway remodeling. All of these features resemble that in humans.²¹ It is clear that the asthmatic allergy is a pathogenic Th2-driven response.

After all six weeks of probiotic feeding, both doses of LMP showed a similar trend on alleviating airway hyperreactivity and regulating productions of OVA-specific antibodies and cytokines in this OVA-sensitized mouse model. Considering all the data, it seemed that both doses were promising on alleviating AHR in this model. The Penh value was reduced in both LMP dose groups. The level of OVA-specific IgE and IgG1 was reduced in high does LMP-treated group, whereas IgG2a level was enhanced in both does LMP dose groups. The above results indicated LMP suppressed allergic reactions related




Figure 4 Effect of oral administration of LMP on IL-12 and IFN- γ production in BALF of OVA-sensitized mice. The concentration of (A) IL-12 and (B) IFN- γ in BALF of mice were assayed using ELISA. Each value represents the mean \pm SD, $n=7$. Groups are normal control (naive), OVA-challenged control (OVA ctrl), low dose LMP (L.L) and high dose LMP (L.H). A difference between the LMP groups and OVA ctrl group was considered statistically significant when $P < 0.05$ (*).

to airway hyperreactivity, production of OVA-specific IgE, IgG1 and IgG2a. Th1 associated cytokines IL-12 and IFN- γ were enhanced in high dose LMP-treated group, while Th2 associated cytokines IL-4, IL-5, IL-13 were reduced in both dose groups. The level of TNF- α were reduced in both dose groups. The data suggested LMP may drive T cell toward Th1 pathway and counter-acting excess Th2 responses. The eosinophils in the low dose LMP-treated group were reduced but this was not observed in the high dose LMP group. It may be due to the anti-asthmatic and pro-inflammatory effects of IFN- γ in the high dose LMP-treated group. The presence of IFN- γ may modulate the effects of IL-13 in the lungs.²² In Ford's mouse model,²² the blockade of IL-13 partially inhibited airway hyperreactivity and goblet cell hyperplasia but not inflammation. The IFN- γ in high dose LMP-treated group may play double-sides effects (inhibiting some, potentiating others) on IL-13-induced changes in the lung. However, this speculation may need further investigation.

In summary, we demonstrated that a lactobacillus multi-species preparation including two strains of *L. plantarum* and *P. acidilactici* exerted an anti-allergy effect by reducing AHR, decreasing OVA-specific IgE production and Th2 cytokines production (IL-4, IL-5, and IL-13).

Lactobacillus species have been widely evaluated for their efficacy to reduce AHR in an OVA-induced allergy model. However, their efficacy in anti-allergy is both species- and strain-dependent. Forsythe et al. (2007)




Figure 5 Effect of oral administration of LMP on IL-4 and IL-5 production in BALF of OVA-sensitized mice. The concentration of (A) IL-4 and (B) IL-5 in BALF of mice were assayed using ELISA. Each value represents the mean \pm SD, $n=7$. Groups are normal control (naive), OVA-challenged control (OVA ctrl), low dose LMP (L.L) and high dose LMP (L.H). A difference between the LMP groups and OVA ctrl group was considered statistically significant when $P < 0.05$ (*).

showed that *Lactobacillus reuteri* but not *L. salivarius* reduced AHR.²³ In the study, *L. reuteri* supplementation reduced influx of inflammatory cells in the BALF and decreased levels of MCP-1, TNF- α and IL-13, but *L. reuteri* supplementation did not affect Th1 responses. Another study showed *Lactobacillus acidophilus* supplementation effectively reduced eosinophil infiltration in BALF and lowered serum IgE production. IL-4 and IL-5 production was reduced whereas IFN- γ levels were enhanced.²⁴ However, *L. acidophilus* did not significantly reduce AHR in the OVA-sensitized mice.²⁴ Previous studies also reported that *L. plantarum* effectively reduced allergic responses in OVA-sensitized mice. For example, heat-killed *L. plantarum* KTCT 3014 reduced IgE production, Th2 cytokines production (IL-4 and IL-5) and enhanced Th1 cytokine IFN- γ levels, however, the Th1 associated IgG2a secretion was lowered.²⁵ In another study, *L. plantarum* K37 effectively reduced AHR and lowered production of serum and OVA-specific IgE production.⁶ Th2 cytokines including IL-4, IL-5 and IL-13 were reduced in the *L. plantarum* K37 supplemented group. IgG2a production as in the present study was enhanced. From the above studies, it is clear that the anti-allergy effects of *Lactobacilli* are both species- and strain-dependent.

P. acidilactici has not been reported for anti-allergy effects in an OVA-induced airway allergy model before. A potential role of *P. acidilactici* for anti-allergy effects is that two strains *P. acidilactici* H0709 and *P. acidilactici* NHNM1 were able to suppress histamine release from




Figure 6 Effect of oral administration of LMP on TNF- α and IL-13 production in BALF of OVA-sensitized mice. The concentration of (A) TNF- α and (B) IL-13 in BALF of mice were assayed using ELISA. Each value represents the mean \pm SD, $n=7$. Groups are normal control (naive), OVA-challenged control (OVA ctrl), low dose LMP (L.L) and high dose LMP (L.H). A difference between the LMP groups and OVA ctrl group was considered statistically significant when $P<0.05$ (*).

ionomycin-stimulated basophils.²⁶ A *P. acidilactici* strain R037 suppressed experimental autoimmune encephalomyelitis by increasing CD4 $^{+}$ Interleukin (IL)-10-producing cells.²⁷ IL-10 has shown to down-regulate excessive Th2 responses in airway inflammation.^{28,29} To the best of our knowledge, this is the first study to report potential anti-allergy effects of *P. acidilactici* in an asthmatic allergy model. This *P. acidilactici* strain was previously selected in a preliminary in vitro and in vivo screen for anti-allergy effects (data not shown). It is possible that *P. acidilactici* PA320 in the lactobacillus multi-species preparation exert the anti-allergy effects via IL-10 or IL-10 producing regulatory cells. Further investigation for its regulation on the specific population of regulatory cells could be performed for greater understanding.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgments

We thank New Bellus Enterprises Co. Ltd. for the production of the test sample LMP.

References

- Djukanović R, Roch WR, Wilson JW, Beasley CRW, Twentyman OP, Howarth PH, Holgate ST. Mucosal inflammation in asthma. Am Rev Respir Dis. 1990;142:434–57.
- Dabbagh K, Takeyama K, Lee HM, Ueki IF, Lausier JA, Nadel JA. IL-4 induces mucin gene expression and goblet cell metaplasia in vitro and in vivo. J Immunol. 1999;162:6233–7.
- Tavernier J, Plaetinck G, Guisez Y. The role of IL-5 in the production and function of eosinophils. In: Whetton AD, Gordon JR, editors. Cell biochemistry. Vol. 7. Hematopoietic cell growth factors and their receptors. New York: Plenum Press; 2000. p. 321–61.
- Huang SK, Xiao HQ, Kleine-Tebbe J. IL-13 expression at the sites of allergen challenge in patients with asthma. J Immunol. 1995;155:2688–94.
- Guarner F, Schaafsma GJ. Probiotics. Int J Food Microbiol. 1998;39:237–8.
- Hooper LV, Littman DR, Macpherson AJ. Interactions between the microbiota and the immune system. Science. 2012;336:1268–73.
- Kim H, Kwack K, Kim DY, Ji GE. Oral probiotic bacterial administration suppressed allergic responses in an ovalbumin-induced allergy mouse model. FEMS Immunol Med Microbiol. 2005;45:259–67.
- Jan RL, Yeh KC, Hsieh MH, Lin YL, Kao HF, Li PH, Chang YS, Wang JY. *Lactobacillus gasseri* suppresses Th17 pro-inflammatory response and attenuates allergen-induced airway inflammation in a mouse model of allergic asthma. Br J Nutr. 2012;108:130–9.
- Liu YW, Liao TW, Chen YH, Chiang YC, Tsai YC. Oral administration of heat-inactivated *Lactobacillus plantarum* K37 modulated airway hyperresponsiveness in ovalbumin-sensitized BALB/c mice. PLOS ONE. 2014;9:e100105.
- Sagar S, Morgan ME, Chen S, Vos AP, Garsen J, van Bergenhenegouwen J, Folkerts G. *Bifidobacterium breve* and *Lactobacillus rhamnosus* treatment is as effective as budesonide at reducing inflammation in a murine model for chronic asthma. Respir Res. 2014;15:46.
- Wu CT, Chen PJ, Lee YT, Ko JL, Lue KH. Effects of immunomodulatory supplementation with *Lactobacillus rhamnosus* on airway inflammation in a mouse asthma model. J Microbiol Immunol. 2015;49:625–35.
- Shida K, Takahashi R, Iwadate E, Takamizawa K, Yasui H, Sato T, Kaminogawa S. *Lactobacillus casei* strain Shirota suppresses serum immunoglobulin E and immunoglobulin G1 responses and systemic anaphylaxis in a food allergy model. Clin Exp Allergy. 2002;32:563–70.
- Nawaz M, Ma C, Basra MAR, Wang J, Xu J. Amelioration of ovalbumin induced allergic symptoms in Balb/c mice by potentially probiotic strains of lactobacilli. Benef Microbes. 2015;6:669–78.
- Nair AB, Jacob S. A simple practice guide for dose conversion between animals and human. J Basic Clin Pharma. 2016;7:27–31.
- Hsieh CC, Hsiao HB, Lin WC. A standardized aqueous extract of *Anoectochilus formosanus* modulated airway hyperresponsiveness in an OVA-inhaled murine model. Phytomedicine. 2010;17:557–62.
- Hamelmann E, Schwarze J, Takeda K, Oshiba A, Larsen GL, Irvin CG. Noninvasive measurement of airway responsiveness in allergic mice using barometric plethysmography. Am J Respir Crit Care Med. 1997;156:766–75.
- Hornbeck P. Assays of antibody production. Current Protocols in Immunology, Vol. Supplement 1. John Wiley & Sons; 1992, 2.1.1–22.
- Mosmann T. Measurement of mouse and human interleukin 10. Curr Protoc Immunol. 2001. Chapter 6: 6.14.

19. Schreiber RD. Measurement of mouse and human interferon gamma. *Curr Protoc Immunol*. 2001; Chapter 6: 6.8.
20. Gueders M, Paulissen G, Crahay C, Quesada-Calvo F, Hacha J, Van Hove C, Cataldo D. Mouse models of asthma: a comparison between C57BL/6 and BALB/c strains regarding bronchial responsiveness, inflammation, and cytokine production. *Inflamm Res*. 2009;58:845–54.
21. Erle DJ, Sheppard D. The cell biology of asthma. *J Cell Biol*. 2014;205:621–31.
22. Ford JG, Rennick D, Donaldson DD, Venkayya R, McArthur C, Hansell E, et al. IL-13 and IFN- γ : Interactions in lung inflammation. *J Immunol*. 2001;167:1769–77.
23. Forsythe P, Inman MD, Bienenstock J. Oral treatment with live *Lactobacillus reuteri* inhibits the allergic airway response in mice. *Am J Respir Crit Care Med*. 2007;175:561–9.
24. Choi I, Lim Y, Na H, Lee H, Choung J, Kook H. The effect of *Lactobacillus acidophilus* on the primary prevention of asthma in a murine asthmatic model. *Pediatr Allergy Respir Dis*. 2008;18:208–18.
25. Hong HJ, Kim E, Cho D, Kim T. Differential suppression of heat-killed lactobacilli isolated from kimchi, a Korean traditional food, on airway hyper-responsiveness in mice. *J Clin Immunol*. 2010;30:449–58.
26. Fujii K, Fujiki T, Koiso A, Hirakawa K, Yamashita M, Matsumoto T, Kataoka Y. Identification of anti-allergic lactic acid bacteria that suppress Ca²⁺ influx and histamine release in human basophilic cells. *J Funct Foods*. 2014;10:370–6.
27. Takata K, Kinoshita M, Okuno T, Moriya M, Kohda T, Honarat JA, Nakatsuji Y. The lactic acid bacterium *Pediococcus acidilactici* suppresses autoimmune encephalomyelitis by inducing IL-10-producing regulatory T cells. *PLoS ONE*. 2011;6:e27644.
28. Oh JW, Seroogy CM, Meyer EH, Akbari O, Berry G, Fathman CG, et al. CD4 T-helper cells engineered to produce IL-10 prevent allergen-induced airway hyperreactivity and inflammation. *J Allergy Clin Immunol*. 2015;110:460–8.
29. Tournoy KG, Kips JC, Pauwels RA. Endogenous interleukin-10 suppresses allergen-induced airway inflammation and non-specific airway responsiveness. *Clin Exp Allergy*. 2000;30:775–83.