

# REVISTA ARGENTINA DE MICROBIOLOGÍA

www.elsevier.es/ram



## INFORME BREVE

### Enhancing adherence of *Arcobacter butzleri* after serial intraperitoneal passages in mice

Heriberto Fernández\*, Sandra Flores, María P. Villanueva, Gustavo Medina, and Macarena Carrizo

Instituto de Microbiología Clínica, Facultad de Medicina, Universidad Austral de Chile, Valdivia, Chile

Received 18 September 2012; accepted 23 April 2013

#### KEYWORDS

*Arcobacter butzleri*;  
Virulence;  
Adhesion;  
Intraperitoneal passage

#### Abstract

We investigated the possibility of enhancing the adherence capacity of four low-adherent *Arcobacter butzleri* strains after serial intraperitoneal passage (i.p.) in mice. All the strains enhanced their adherence capacity after the first passage, increasing their adhesion rates after each passage. These results suggest that i.p. passage enhances the expression of adherence in *A. butzleri* strains.

© 2013 Asociación Argentina de Microbiología. Published by Elsevier España, S.L. All rights reserved.

#### PALABRAS CLAVE

*Arcobacter butzleri*;  
Virulencia;  
Adherencia;  
Pasaje intraperitoneal

#### Incremento de la adherencia de *Arcobacter butzleri* por sucesivos pasajes intraperitoneales en ratón

#### Resumen

Se investigó la posibilidad de incrementar la capacidad de adherencia de cuatro cepas de *Arcobacter butzleri* de baja adherencia por sucesivos pasajes por peritoneo de ratón. Todas las cepas aumentaron su capacidad de adherencia después del primer pasaje, e incrementaron las tasas de adherencia después de cada pasaje. Estos resultados sugieren que es posible incrementar la expresión de la adherencia en *A. butzleri* por pasajes intraperitoneales sucesivos en ratón.

© 2013 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

\*Corresponding author.

E-mail address: hfernand@uach.cl (H. Fernández).

The genus *Arcobacter* (Latin: *arcus*, arch, and Greek: *bacter*, bacteria) was proposed in 1991 to classify a group of bacteria previously considered to be members of the *Campylobacter* genus. Molecular studies show that these bacteria have different genotypic characteristics from those of *Campylobacter*, giving rise to the new genus *Arcobacter*<sup>1</sup>. This group is composed of spiral curved, or "S"-shaped, Gram-negative bacilli of 0.2 to 0.6  $\mu\text{m}$  diameters and 3  $\mu\text{m}$  length. They have one or two flagella and grow at temperatures ranging from 15 to 37 °C, although the optimal growth temperature is 30 °C. The primary isolation of these bacteria can be achieved under microaerobic conditions, but when using subsequent cultures, they can grow in aerobiosis. Today, the genus *Arcobacter* is composed of the following species: *Arcobacter cryaerophilus*, *Arcobacter butzleri*, *Arcobacter skirrowii*, *Arcobacter nitrofrigidus*, *Arcobacter cibarius*, *Arcobacter halophilus*, *Arcobacter mytili*, *Arcobacter thereius*, *Arcobacter marinus*, *Arcobacter trophiarum*, *Arcobacter defluvi*, *Arcobacter molluscorum* and *Arcobacter ellisii*; however, only the first three have been isolated from human and animal samples<sup>1-3</sup>.

*Arcobacter butzleri* is an emerging pathogen that has been associated with abortion and enteritis in animals, as well as with diarrhea and bacteremia in adults and children. Currently, *A. butzleri* is considered the most common species of the genus in environmental water, food and clinical samples, being ranked as the fourth most common campylobacter organism isolated from human feces<sup>1,2</sup>. Recently, *A. butzleri* has been considered a serious hazard to human health by the International Commission on Microbiological Specification for Foods<sup>1</sup>.

The presence of nine putative virulence genes has been reported in *A. butzleri* ATCC49616 strain and in several human and animal strains by Miller *et al.*<sup>4</sup> and Doudah *et al.*<sup>5</sup>, respectively. Some of them, such as *cadF* (fibronectin binding protein gene), *cj1349* (fibronectin binding protein gene), *flaA* (flagellin A protein gene) and *hecA* (hemagglutinin gene) share similarities with the virulence genes of *Campylobacter jejuni* and other bacteria to which they have been associated in adherence<sup>4,5</sup>.

Adhesive capacity to different epithelial cell lines is a bacterial characteristic associated to pathogenicity, which has been described in *A. butzleri* by several authors<sup>6,7</sup>. However, there is no information about changes in the pathogenicity of *A. butzleri* after animal passages as it occurs with *C. jejuni*. In fact, several other biological models, such as intraperitoneal animal passage<sup>8</sup>, intragastric passage in chicks<sup>9</sup> and chick embryo passage<sup>10,11</sup> have all been used to either enhance or restore virulence capacities and culturability in *Campylobacter* strains.

The aim of this study was to investigate the possibility of enhancing adhesive properties in weak-adherent *A. butzleri* strains through intraperitoneal passage (i.p.) in mice.

Four strains of *A. butzleri* from the collection owned by our Institute, originally isolated from river water (three strains) and from chicken liver for human consumption (one strain) were studied. These strains were previously characterized as weakly-adherent to HEp-2 cells by means of the protocol currently being used in our laboratory<sup>2</sup>. In brief, four Leighton tubes containing a coverslip with the


HEp-2 cell monolayers were used for each strain. They were inoculated with 1 ml of the bacterial suspensions ( $10^6$  CFU) prepared in the same medium that had been used for growing HEp-2 cells [RPMI medium (Gibco) supplemented with 10% fetal calf serum (Gibco)] and incubated at 37 °C during 3 h, under 5% CO<sub>2</sub>. Afterwards, coverslips were washed four times with phosphate-buffered saline pH 7.4, fixed with methanol, stained with 10% Giemsa stain and examined under light microscopy (1000 X).

In order to assess induction of adhesive capacity, each strain was subjected to five serial i.p. passages in Pockefeller (3-6 weeks age) mice, being the original strain considered as passage zero. Inocula were prepared in Brucella broth (BB) (Difco) and each animal received 1 ml of a bacterial suspension containing  $2 \times 10^9$  CFU/ml. After 24 h, mice were euthanized with CO<sub>2</sub> and their peritoneal contents were aseptically removed and seeded on blood agar plates that were incubated for 48 h at 30 °C in a microaerobic atmosphere. The adhesive capacity of each of the resulting pure cultures of the isogenic strain was determined by inoculating HEp-2 cell cultures following the above described method. The adhesion rate (number of cells showing adhered bacteria/total cells examined  $\times$  100) and the number of bacteria observed in each HEp-2 cell were determined by counting at least 200 cells.

The average number of adhering bacteria was estimated in a minimum of 35 HEp-2 cells with adhered bacteria. All the tests were carried out twice and in duplicate.

The expression of *flaA* gene in strains AC11 and HP12 was determined by means of the semi-quantitative RT-PCR (sqRT-PCR) technique described by Niehus *et al.*<sup>12</sup>, using the following *flaA* primer sequences: sense 5'-CAGTTGCACAGCTGACATT-3' and antisense 5'-TGCAAGAACTGCAAAAGGTG-3' (amplicon size 158 pb). The *rpoB* (housekeeping gen) primer sequences used were: sense 5'-CAACTCTTCAACACCATTAACAA-3' and antisense 5'-AGGTAGCGAAGTTGGTAAACCT-3' (amplicon size 230 bp). First, cDNA was synthesized from total RNA by TaqMan® (Roche) Reverse Transcription Reagents according to the manufacturer's instructions. Then, RT-PCR reaction was performed using KOD Hot Start DNA Polymerase Novagen®, according to the manufacturer's instructions. Products from sqRT-PCR were separated and visualized by electrophoresis in agarose gels stained with ethidium bromide. Images of RT-PCR ethidium bromide-stained agarose gels were taken and band quantification was performed with the ImageJ gel analysis free software to determine pixel intensity. Data were normalized with regard to *rpoB* pixel intensity and expressed as mean  $\pm$  standard error of the mean (SEM). Semi-quantitative measurements of RNAs were expressed as arbitrary units (AU). Statistical significance was verified by two-way ANOVA - Dunnett's Multiple Comparison Test. A value of  $p < 0.05$  was considered statistically significant. Data analysis was done with GraphPad Prism5 software.

In order to establish isogenicity, the original strains and the isolates obtained after each i.p. were genotyped using the enterobacterial repetitive intergenic consensus PCR (ERIC-PCR) technique according to Houf *et al.*<sup>13</sup>. All the isolates obtained after each i.p. passage showed the same electrophoretic pattern displayed by their respective original strains. As an example, Figure 1 shows the ERIC-PCR electrophoretic patterns of strains AC11 and HP 12.



**Figure 1** ERIC-PCR electrophoretic patterns of strains AC11 and HP12 and their isogenic strains derived after five successive i.p. passages. Lines 1, 2, 3, 4, 5, 6: strain AC11 P0, P1, P2, P3, P4, P5, respectively. Lines 7, 8, 9, 10, 11, 12: strain HP12 P0, P1, P2, P3, P4, P5, respectively. Line 13: negative control.

As shown in Table 1, all the strains enhanced their adhesive capacity even after the first i.p. passage. Expression of adhesive capacity appears to be enhanced


after each consecutive passage, with the adhesion rates and the number of adherent bacteria increasing from the first to the fifth passage. The adhesion rate increased between 5% (strain AC11) and 15.5% (strain AA10) after the first passage and between 15% (strain HP12) and 49.5% (strain AA10) after the fifth passage. The increase in the adhesion rates of all the passages of each strain showed statistical differences (Student's t-test) when compared with the adherence rate of passage 0, except for passage 2 of strain HP12.

Strain HP12, isolated from chicken liver, showed the lowest number of adhering bacteria per cell ( $1.49 \pm 0.93$  and  $2.57 \pm 1.10$ ) after the first and the fifth passages, whereas strains AC11 and AA10, isolated from river water, showed the highest numbers of adhered bacteria per cell after the fifth passage ( $3.98 \pm 2.79$  and  $4.09 \pm 3.42$ , respectively). However, numbers of adherent bacteria increased after each passage, with the exception of passage 5 of strain AA10. No differences (Student's t-test) were observed when comparing the increase of adherent bacteria in each passage with the number of adherent bacteria observed in passages 0 of each strain.

**Table 1** Enhancement of the adhesive capacity of *Arcobacter butzleri* through intraperitoneal passages in mice

Strain	Origin	Passage number	Adherence rate (%) <sup>a</sup>	Number of adherent bacteria/cell <sup>b</sup>
AC11	River water	0	46.0 $\pm$ 0.58	1.53 $\pm$ 1.03
		1	51.0 $\pm$ 1.53	1.91 $\pm$ 1.01
		2	65.0 $\pm$ 1.52	2.12 $\pm$ 1.34
		3	68.0 $\pm$ 1.53	2.41 $\pm$ 1.42
		4	67.0 $\pm$ 1.00	2.30 $\pm$ 1.45
		5	83.0 $\pm$ 2.87	3.98 $\pm$ 2.79
AP5	River water	0	44.5 $\pm$ 0.86	1.49 $\pm$ 0.93
		1	51.0 $\pm$ 4.58	2.66 $\pm$ 1.32
		2	62.5 $\pm$ 1.50	1.87 $\pm$ 1.33
		3	68.0 $\pm$ 1.00	1.65 $\pm$ 0.96
		4	66.5 $\pm$ 1.00	2.26 $\pm$ 1.51
		5	66.0 $\pm$ 0.76	2.77 $\pm$ 1.30
AA10	River water	0	31.0 $\pm$ 1.73	0.96 $\pm$ 0.61
		1	46.5 $\pm$ 1.32	1.72 $\pm$ 1.04
		2	59.5 $\pm$ 2.00	1.74 $\pm$ 1.01
		3	58.5 $\pm$ 3.46	2.14 $\pm$ 1.62
		4	73.5 $\pm$ 4.58	3.55 $\pm$ 1.18
		5	80.5 $\pm$ 1.80	4.09 $\pm$ 3.42
HP12	Chicken liver	0	45.0 $\pm$ 1.32	1.49 $\pm$ 0.93
		1	53.0 $\pm$ 0.57	2.66 $\pm$ 1.32
		2	47.5 $\pm$ 1.50	1.87 $\pm$ 1.33
		3	53.0 $\pm$ 1.52	1.65 $\pm$ 0.96
		4	59.0 $\pm$ 0.76	2.26 $\pm$ 1.51
		5	60.0 $\pm$ 2.64	2.57 $\pm$ 1.10

<sup>a</sup> $p < 0.05$  when comparing the results of each passage with that of passage 0, except passage 2 of strain AA10; <sup>b</sup> $p > 0.05$  when comparing the results of each passage with that of passage 0, except passage 5 of strain AA10.



**Figure 2** Relative semi-quantification of *flaA* gene expression in strains HP12 and AC11 and their isogenic strains derived after i.p. passages P1, P3 and P5. AU: arbitrary units; ns: non-significant; P0: Passage 0; P1: Passage 1; P3: Passage 3; P5: Passage 5.

\* $p < 0.05$ .

In general, despite the description of nine putative virulence genes<sup>4,5</sup>, little is known about the pathogenic mechanisms or potential virulence factors of *Arcobacter* spp. and how they could be expressed in both *in vitro* and *in vivo* models. In this study, we used the mouse i.p. passage model as a biological support, in an attempt to increase the adhesive capacity in *A. butzleri* strains that had previously shown low adherence rates. As shown in Table 1, all four strains enhanced their adhesive capacity after several consecutive passages, with the adhesion rates and the number of adherent bacteria increasing from the first to the fifth passage, with very few exceptions.

Several biological models have been used to enhance virulence in different bacteria. Fernández *et al.*<sup>8</sup> reported that *C. jejuni* and *Campylobacter coli* strains enhanced their enterotoxigenic and invasive capacities after i.p. passages in mice. Hänel *et al.*<sup>9</sup> demonstrated that upon passage through the chicken gut, a *C. jejuni* strain enhanced its adherence to eukaryotic cells.

Despite the existence of several studies on adhesion, invasion and toxigenic capacities, the pathogenicity and virulence mechanisms of *A. butzleri* are still poorly understood<sup>1</sup>. On the other hand, there are few experimental trials exploring the pathogenic properties of *Arcobacter* in animal models.

The first *in vivo* study inoculating *Arcobacter* strains intravenously or intraperitoneally in rodents was unsuccessful. No clinical symptoms were observed and no internal lesions were detected after postmortem examination<sup>14</sup>. However, the invasion and colonization capacity of *A. butzleri* has been demonstrated in White turkeys by Wesley and Baetz<sup>15</sup>, who reported that adapting *A. butzleri* to poultry via serial passage may increase the ability of the organism to colonize outbred birds.

The mouse i.p. passage model used in this study enhanced the adherence capacity of *A. butzleri* strains after the first passage, with an increase in adhesion rates after each passage giving a new insight into the pathogenicity mechanisms of this bacterium. These results show that this

model is suitable for inducing expression of adhesion capacity in *A. butzleri*, having the additional advantage that the isogenic variants can be isolated in pure cultures on non-selective media due to the absence of contaminant microbiota in the i.p. cavity.

In an attempt to have an approximation to the molecular basis that could explain the increase of the adhesion capacity in the strains studied, we analyzed the expression of the *flaA* gene in the original strains AC11 and HP12 and their isogenic strains derived from i.p. passages 1, 3 and 5. As shown in Figure 2, the expression of the *flaA* gene of strain AC11 was significantly increased.

Miller *et al.*<sup>4</sup> and Doudah *et al.*<sup>5</sup> reported nine putative virulence genes in *A. butzleri* that had been previously reported in *C. jejuni*. It is known that the *flaA* gene is associated with adherence capacity in *C. jejuni* and *A. butzleri*<sup>4,5</sup>. The increased expression of the *flaA* gene after the fifth i.p. passage in strain AC11, although still not conclusive, suggests that changes in the gene expression of *A. butzleri* can occur through successive passages. However, further studies are necessary to elucidate the molecular basis of this phenomenon and also to determine if *A. butzleri* can carry its own virulence genes, other than those reported by Miller *et al.*<sup>4</sup> and Doudah *et al.*<sup>5</sup>.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

The authors would like to thank Dr. Hans G. Richter (Departamento de Anatomía, Histología y Patología, Universidad Austral de Chile) for his advice and critical review. This work was financially supported by grant FONDECYT 1110202.

## References

1. Baffone W, Casaroli A, Citterio B, Pierfelici L, Campana R, Vittoria E, Guaglianone E, Donelli G. *Campylobacter jejuni* loss of culturability in aqueous microcosms and ability to resuscitate in a mouse model. *Int J Food Microbiol.* 2006;107:83-91.
2. Bückler R, Troeger H, Kleer J, Fromm M, Schulzke JD. *Arcobacter butzleri* induces barrier dysfunction in intestinal HT-29/B6 cells. *J Infect Dis.* 2009; 200:756-64.
3. Collado L, Figueras MJ. Taxonomy, epidemiology, and clinical relevance of the genus *Arcobacter*. *Clin Microbiol Rev.* 2011;24:1174-92.
4. Doudah L, de Zutter L, Baré J, DeVos P, Vandamme P, Vandenberg O, Van den Abeele AM, Houf K. Occurrence of putative virulence genes in *Arcobacter* species isolated from humans and animals. *J Clin Microbiol.* 2012;50:735-41.
5. Fernández H, Vivanco T, Eler G. Expression of invasiveness of *Campylobacter jejuni* subsp. *jejuni* after serial intraperitoneal passages in mice. *J Vet Med B.* 2000;47:635-9.
6. Fernández H, Flores S, Inzunza F. *Arcobacter butzleri* strains isolated from different sources display adhesive capacity to epithelial cells *in vitro*. *Acta Scient Vet.* 2010;38:287-91.
7. Field LH, Underwood JL, Payne SM, Berry LJ. Characteristics of an avirulent *Campylobacter jejuni* strain and its virulence-enhanced variants. *J Med Microbiol.* 1993;38:293-300.
8. Figueras MJ, Levican A, Collado L, Inza MI, Yustes C. *Arcobacter ellisii* sp. nov., isolated from mussels. *Syst Appl Microbiol.* 2011;34:414-8.
9. Hänel I, Borrmann E, Müller J, Müller W, Pauly B, Liebler-Tenorio EM, Schulze F. Genomic and phenotypic changes of *Campylobacter jejuni* strains after passage of the chicken gut. *Vet Microbiol.* 2009;136:121-9.
10. Higgins R, Degre R. Isolation of spirillum-like organisms from pig and bovine fetuses. *Vet Rec.* 1979;104:262-3.
11. Hoa TKH, Lipman LJA, Hendriks HGCJM, Tooten PCJ, Ultee T, Gastra W. Interaction of *Arcobacter* spp. with human and porcine intestinal epithelial cells. *FEMS Immunol Med Microbiol.* 2007;50:51-8.
12. Houf K, de Zutter L, Van Hoof J, Vandamme P. Assessment of the genetic diversity among arcobacters isolated from poultry products by using two PCR-based typing methods. *Appl Environ Microbiol.* 2002;68:2172-8.
13. Miller WG, Parker CT, Rubenfield M, Mendz GL, Wösten MM, Usery DW, Stolz JF, Binnewies TT, Hallin PF, Wang G, Malek JA, Rogosin A, Stanker LH, Mandrell RE. The complete genome sequence and analysis of the epsilonproteobacterium *Arcobacter butzleri*. *PLoS One.* 2007;2:e1358.
14. Niehus E, Gressmann H, Fang Y, Schlapbach R, Dehio M, Dehio C, Stack A, Meyer TF, Suerbaum S, Josenhans C. Genome-wide analysis of transcriptional hierarchy and feedback regulation in the flagellar system of *Helicobacter pylori*. *Mol Microbiol.* 2004;52:947-61.
15. Wesley IV, Baetz AL. Natural and experimental infections of *Arcobacter* in poultry. *Poult Sci.* 1999;78:536-45.