



ORIGINAL ARTICLE

Genetic diversity of thermotolerant *Campylobacter* spp. isolates from different stages of the poultry meat supply chain in Argentina



María V. Zbrun^{a,c}, Analía Romero-Scharpen^c, Carolina Olivero^a,
Jorge A. Zimmermann^c, Eugenia Rossler^c, Lorena P. Soto^{a,c}, Diego M. Astesana^c,
Jesica E. Blajman^c, Ayelén Berisvil^c, Laureano S. Frizzo^{a,c,*}, Marcelo L. Signorini^{a,b}

^a Laboratory of Food Analysis, Institute of Veterinary Science (ICiVet Litoral), National University of the Littoral, National Council of Scientific and Technical Research (UNL/CONICET), Esperanza, Province of Santa Fe, Argentina

^b National Council of Scientific and Technical Research (CONICET), National Institute of Agricultural Technology (INTA), EEA Rafaela, Rafaela, Province of Santa Fe, Argentina

^c Department of Public Health, Faculty of Veterinary Science-National University of the Littoral, Esperanza, Province of Santa Fe, Argentina

Received 21 October 2016; accepted 3 March 2017

Available online 13 July 2017

KEYWORDS

Campylobacter coli;
Campylobacter jejuni;
Pulsed-field gel electrophoresis;
Poultry meat

Abstract The objective of this study was to investigate a clonal relationship among thermotolerant *Campylobacter* spp. isolates from different stages of the poultry meat supply chain in Argentina. A total of 128 thermotolerant *Campylobacter* spp. (89 *C. jejuni* and 39 *C. coli*) isolates from six poultry meat chains were examined. These isolates were from: a) hens from breeder flocks, b) chickens on the farm (at ages 1 wk and 5 wk), c) chicken carcasses in the slaughterhouse, and d) chicken carcasses in the retail market. Chickens sampled along each food chain were from the same batch. *Campylobacter* spp. isolates were analyzed using pulsed-field gel electrophoresis to compare different profiles according to the source. Clustering of *C. jejuni* isolates resulted in 17 profiles, with four predominant genotypes and many small profiles with just a few isolates or unique patterns, showing a very high degree of heterogeneity among the *C. jejuni* isolates. Some clusters included isolates from different stages within the same chain, which would indicate a spread of strains along the same poultry meat chain. Moreover, twenty-two strains of *C. coli* clustered in seven groups and the remaining 17 isolates exhibited unique profiles. Evidence for transmission of thermotolerant *Campylobacter* spp. through the food chain and cross contamination in the slaughterhouses were obtained. This collective

* Corresponding author.

E-mail address: lfrizzo@fvc.unl.edu.ar (L.S. Frizzo).

PALABRAS CLAVE

Campylobacter coli;
Campylobacter jejuni;
 Electroforesis de campos pulsados;
 Carne aviar

evidence should be considered as the scientific basis to implement risk management measures to protect the public health.

© 2017 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Diversidad genética de *Campylobacter* spp. termotolerantes aislados de diferentes etapas de la cadena cárnica aviar en Argentina

Resumen El objetivo de este trabajo fue investigar la relación clonal entre aislamientos de *Campylobacter* spp. termotolerantes obtenidos de diferentes etapas de la cadena cárnica aviar en Argentina. En total se examinaron 128 aislamientos de *Campylobacter* spp. (89 de *Campylobacter jejuni* y 39 de *Campylobacter coli*) obtenidos de 6 cadenas cárnicas muestreadas en los siguientes puntos del circuito productivo: a) gallinas reproductoras; b) pollos en las granjas (de una y 5 semanas de edad); c) carcasas de pollo en frigorífico, y d) carcasas de pollo en puntos de venta final. Las muestras de pollos fueron obtenidas a lo largo de las cadenas cárnicas siguiendo el mismo lote. Los aislamientos de *Campylobacter* spp. fueron analizados mediante electroforesis de campos pulsados y se compararon los diferentes perfiles. Los aislamientos de *C. jejuni* se agruparon en 17 perfiles, 4 de ellos predominantes y el resto en perfiles que agruparon pocos aislamientos o patrones únicos, lo que ilustra una gran heterogeneidad. Algunos agrupamientos incluyeron aislamientos obtenidos de diferentes etapas de una misma cadena cárnica, lo cual indicaría una dispersión de cepas a lo largo de las cadenas cárnicas. Por otra parte, 22 aislamientos de *C. coli* se agruparon en 7 grupos y otros 17 aislamientos presentaron perfiles únicos. Se obtuvieron evidencias de transmisión de *Campylobacter* spp. termotolerante en la cadena cárnica aviar y contaminación cruzada en frigoríficos. La evidencia reunida debería servir como base científica para implementar estrategias de manejo del riesgo, destinadas a proteger la salud de los consumidores de carne aviar.

© 2017 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Thermotolerant *Campylobacter* spp., especially *C. jejuni* and *C. coli*, causes significant morbidity both in developing and developed countries^{10,13} and constitutes the most frequent cause of foodborne illness worldwide^{4,5}. Poultry meat is one of the most important sources of human campylobacteriosis. The reduction and elimination of thermotolerant *Campylobacter* spp. in the food chain, particularly from chicken products, are major strategies in the efforts to control human exposure and to improve the public health¹¹.

In developing countries, information on food-borne disease is scant due to the inadequate data provided by the passive/non-existent surveillance systems. Additionally, outbreak information is frequently unsubstantial because health authorities lack the capabilities or resources for detection of diarrheal diseases¹⁸. A study conducted in Argentina⁶ concluded that thermotolerant *Campylobacter* spp. was the most important gastrointestinal pathogen in humans whose incidence rate was higher than other common pathogens such as *Salmonella* spp., *Shigella* spp., and *Escherichia coli*.

Recent studies have shown high prevalence of thermotolerant *Campylobacter* spp. in poultry carcasses in slaughterhouses and the retail market in Argentina¹⁹, most of which were resistant to quinolones and erythromycin²⁰. However, there is a lack of research in Argentina conducted

with the aim to investigate the epidemiology of campylobacteriosis in the whole food chain from farm to fork. Molecular typing is a useful tool to enhance epidemiological studies¹³. Pulsed-field gel electrophoresis (PFGE) is considered a gold standard due to its high discrimination potential and this information is essential to develop effective plans to control the disease¹⁷.

The objective of this study was to investigate a clonal relationship among thermotolerant *Campylobacter* spp. isolates from different stages of the poultry meat supply chain in Argentina.

Materials and methods

Collection of *Campylobacter* isolates

A total of 128 thermotolerant *Campylobacter* spp. strains were isolated from six different companies (here referred to as poultry meat food chains) in the Santa Fe region of Argentina during the years 2011–2012. The stages sampled in each poultry meat chain were: a) hens from breeder flocks (n=75), b) broilers in flocks (aged <1 wk (n=180) and >5 wk (n=180)), c) chicken carcasses in the slaughterhouse (n=60), and d) chicken carcasses in the retail market (n=60). The chickens sampled along the meat supply chain were from the same batch (defined as a group of chickens

from the same flock, sent to the same slaughterhouse at the same time, and sold together in the same retail market). Additionally, samples of litter (n=24), feed (n=24), and drinking water (n=24) were taken from the flocks. At the slaughterhouses, cecal (n=60) and liver (n=60) samples were randomly collected from the evisceration line.

Fecal samples (hens from breeder flocks and broilers in flocks) were randomly collected from the cloaca using sterile cotton swabs, which were placed in capped plastic tubes containing 10 ml of Cary-Blair (Britania[®]) transport medium and transported to the laboratory under refrigeration conditions within 4 h. Together with the cloacal samples, samples of chicken feed (500 g), drinking water (1 l) and litter (500 g) were also taken from each flock. Feed, drinking water, and litter samples were taken directly from the feeders. Cecal and liver samples in the slaughterhouses were randomly collected from the evisceration line by one of the researchers and placed into sterile plastic bags. Broiler carcasses were taken from the processing line after chilling, using a clean pair of latex gloves and put into a sterile bag with 200 ml of Ringer's solution ¼ strength. Carcasses were rinsed by shaking for 60 seconds in each of two directions to ensure that the solution came into contact with all the surfaces; the solution was recovered and transported to the lab in sterile plastic tubes (under refrigeration conditions), within 4 h. Chickens were packaged at the processing plant in the slaughterhouse and transported to the retail market where whole chickens from the same flock were randomly sampled, following the same procedure described for the broiler carcasses in the slaughterhouse.

Campylobacter spp. were isolated using the selective media Bolton Broth (OXOID[®]) and Preston Agar (OXOID[®])². Isolation of *Campylobacter* spp. was performed by a previously described methodology^{2,8}. All incubations were performed under microaerophilic conditions (5% O₂, 10% CO₂ and 85% H₂) at 42 °C. Preliminary identification of thermotolerant *Campylobacter* spp. isolates was based on colony morphology, microscopic appearance (curved gram-negative bacilli with typical motility), and the following phenotypic characteristics: oxidase and catalase production⁸. All presumptive *Campylobacter* spp. isolates were identified to the species level (*C. jejuni* and *C. coli*) by multiplex PCR, as proposed by Vandamme et al.¹⁵. DNA was extracted using

a Wizard genomic DNA purification kit (Promega[®]) and PCR products were analyzed on 1.5% agarose gels and stained with GelRed (Biotium[®]).

Positive isolates were subcultured on Columbia blood agar and stored in glycerol broth (15% glycerol and 85% serum broth) at -80 °C¹⁴.

PFGE-typing

Analysis of *C. jejuni* (n=89) and *C. coli* (n=39) isolates by PFGE was performed according to the method described in the PulseNet protocol¹² using *Sma*I (Fermentas[®]) as restriction endonuclease. *Salmonella* spp. H9812 was used as reference marker (digested with *Xba*I Fermentas[®])⁷. PFGE banding patterns were analyzed using BioNumerics version 6.6 (Applied Maths, Belgium). Images of gels were normalized by alignment with the appropriate size standard lanes. Matching and dendrogram of fingerprints were determined by the unweighted pair group method with averages (UPGMA) and performed using the Dice coefficient (position tolerance, 1.0%). The PFGE cluster was based on a 95% similarity cut off.

Results

Among the 128 isolates available, 89 were *C. jejuni* and 39 were *C. coli*. The origin of the isolates is shown in Table 1.

Overall, clustering of *C. jejuni* isolates resulted in 17 profiles, with four predominant genotypes (clusters E, F, H, and K) shared by four or five isolates, except profile F which was shared by 12 isolates (Table 2). Many small profiles with just a few isolates (two or three) were observed. Additionally, it is important to mention that only 56 of 89 isolates were grouped, the rest presented unique patterns (Fig. 1), showing a very high degree of heterogeneity among the *C. jejuni* isolates. Seven out of 17 profiles were constituted by isolates from a single stage. Among them, only group A consisted of isolates from breeding hens isolated from different chains. The remaining profiles contained isolates from different stages of the poultry meat chain. Some clusters (E, F, H, I, and L) included isolates from different stages within the same chain, which would indicate a spread of

Table 1 Origin of thermotolerant *Campylobacter* isolates from different stages of poultry meat supply chain

Poultry chain stage	Sample type	Number of isolates (% of total)	
		<i>C. jejuni</i>	<i>C. coli</i>
Hen's farm	Cloacal feces	19 (14.8%)	16 (12.5%)
Broiler's farm 1 wk	Cloacal feces	7 (5.5%)	2 (1.6%)
Broiler's farm 5 wk	Cloacal feces	22 (17.2%)	5 (3.8%)
	Drinking water	1 (0.8%)	0
	Feed	1 (0.8%)	0
	Carcasses	10 (7.8%)	7 (5.5%)
Slaughterhouse	Cecum	7 (5.5%)	1 (0.8%)
	Liver	3 (2.3%)	2 (1.6%)
	Carcasses	19 (14.8%)	6 (4.7%)
Retail market	Carcasses	19 (14.8%)	6 (4.7%)
Total		89 (69.5%)	39 (30.5%)

Table 2 PFGE profiles of *C. jejuni* isolated at different stages of the poultry meat chain

Profile	Isolates (n)	Samples/total (%)	Stage of poultry meat chain ^a	Chain (1–6)
A	2	2.2	H	1–6
B	3	3.3	H	5
C	2	2.2	RM	1
D	2	2.2	P(5 wk) Sl	5
E	4	4.4	P(5 wk) RM	5
F	12	13.3	Sl RM P(w5) Sce	3
G	3	3.3	Sce P(5w)	1–5–6
H	5	5.6	P(1w) W(5w)	4
I	3	3.3	Sce RM	4
J	2	2.2	Sl H	6–3
K	4	4.4	Sl H	1–3–4
L	3	3.3	P(5w) RM	2
M	3	3.3	Sl	3
N	2	2.2	P(5w)	2
O	2	2.2	RM	2
P	2	2.2	Sl (P(5w))	3–6
Q	2	2.2	H	3
Total	56	62.9		

^a References: H: breeding hens, P(1w): poultry <1 wk, P(5w): poultry >5 wk, W(5w): drinking water in poultry flocks >5 wk, Sl: carcass at slaughterhouse, Sce: caeca at slaughterhouse, RM: carcass at retail market.

strains along the same poultry meat chain. The same strain of *C. jejuni* in broiler flock and carcasses in the slaughterhouse and the retail market could be observed. Another remarkable finding was that four out of 17 profiles (G, J, K, and P) were represented by isolates obtained from different stages in different poultry meat chains.

On the one hand, twenty-two strains of *C. coli* were clustered into seven groups (Table 3) and the remaining 17 isolates presented unique profiles (Fig. 2). Profiles D and F were predominant and each of them had five isolates. The others were small profiles with just a few isolates (two or three). Three (A, C, and G) out of these seven profiles were represented by isolates obtained from the same stage within the same poultry meat chain. Profile B was constituted by isolates from breeding hens and drinking water in the broiler flock from two different chains. On the other hand, profile D included all isolates from the same poultry meat chain but isolated from two different stages (carcasses in the slaughterhouse and the retail market), which supports

the hypothesis of thermotolerant *Campylobacter* spp. transmission through the poultry meat chain. Profiles E and F were constituted by isolates from different stages of the chain and at the same time from different poultry meat chains.

Discussion

Transmission along the food chain is generally recognized as a major source of human campylobacteriosis^{4,5,21}. This study was conducted with the aim to evaluate the genetic diversity among thermotolerant *Campylobacter* spp. isolates and then to follow up *Campylobacter* spp. isolates along the poultry meat chain from hens through the carcass in the retail market. This transmission of strains throughout the poultry meat supply chain was observed in certain isolates and especially in the later stages of the poultry meat chain (broilers in the flocks and carcasses in slaughterhouses and the retail market). This finding supports the

Table 3 PFGE profiles of *C. coli* isolated at different stages of the poultry meat chain

Profile	Isolates (n)	Samples/total (%)	Stage of poultry meat chain ^a	Chain (1–6)
A	2	5.1	Sce	6
B	3	7.7	H W(1w)	1–2
C	2	5.1	H	5
D	5	12.8	Sce RM	5
E	2	5.1	P(5w) H	1–2
F	5	12.8	Sl Sce H RM	1–2–3–5
G	3	7.7	H	2
Total	22	56.4		

^a References: H: breeding hens, P(5w): poultry >5 wk, W(1w): drinking water in poultry flocks >1 wk, Sl: carcass at slaughterhouse, Sce: caeca at slaughterhouse, RM: carcass at retail market.




Figure 1 Dendrogram of *C. jejuni* Smal PFGE profiles isolated at different stages of the poultry meat supply chain. References: Ch: chain; S: stage of the poultry meat supply chain; H: breeding hens, P(1w): poultry <1 wk, P(5w): poultry >5 wk, W(5w): drinking water in poultry flocks >5 wk, SI: carcass at slaughterhouse, Sc: ceca at slaughterhouse, RM: carcass at retail market.




Figure 2 Dendrogram of *C. coli* Smal PFGE profiles isolated at different stages of the poultry meat supply chain. References: Ch: chain; S: stage of the poultry meat supply chain; H: breeding hens, P(1w): poultry <1 wk, P(5w): poultry >5 wk, W(5w): drinking water in poultry flocks >5 wk, SI: carcass at slaughterhouse, Sc: ceca at slaughterhouse, RM: carcass at retail market.

hypothesis that certain genotypes may be transmitted along the poultry meat chain as was observed in other countries such as Hungary⁴, Slovenia and Bosnia and Herzegovina²¹, and Spain¹⁰. These matching *Campylobacter* spp. profiles were able to survive under environmentally adverse conditions along the poultry production chain. However, not all the isolates showed this behavior because some strains isolated at early stages were not recovered at later stages of the poultry meat chain.

Moreover, the observed matching of isolates from different chains would have various explanations. For example, it may be a reflection of a common source of contamination. Because all farms were part of the same integrated production system and basically applied the same system for sanitization, feeding, animal health procedures between the farms, *Campylobacter* spp. may have survived in

drinking water in poultry flocks >5 wk, SI: carcass at slaughterhouse, Sc: ceca at slaughterhouse, RM: carcass at retail market.

certain niches, fomites or vectors that were present on any of these and transported to other farms and slaughterhouses. In this sense, further studies should be conducted with the aim to establish the role of animate (birds, insects, small mammals, poultry farmer) and inanimate agents (poultry litter, supply air, food, water, trucks) in the maintenance of thermotolerant *Campylobacter* spp. among different production cycles within a chain and among different chains.

The most common profile seems to support the idea that each stage of the poultry meat chain has stable and particular genotypes. Additionally, the results of our study showed that the flocks exhibited different *C. jejuni* and *C. coli* profiles. Several works showed similar quantity of clusters for *C. jejuni* and many unique profiles for this species^{9,10}. The genetic diversity of both *Campylobacter* species was also reported in previous studies^{4,17,21,22} and supported by an acquisition of foreign DNA or random recombination of large DNA segments, may well cause alterations detectable using PFGE¹⁶. As a previous work¹⁰, genotype diversity of the isolates for flocks increased through the poultry production chain, with the highest diversity being detected at the slaughterhouse level. However, it was interesting that the isolates recovered in the carcasses in the slaughterhouses and the retail market were not identified in previous stages of the same chain. It is possible that *Campylobacter* spp. can survive in the slaughterhouse environment even after cleaning, as suggested by Melero et al.¹⁰. The ability of *C. jejuni* to form biofilms as protective mechanisms and to survive longer and increase resistance to disinfectants, antimicrobials, and antibiotics was recognized³. Biofilm formation on abiotic surfaces may help *Campylobacter* spp. survive in the environment and these biofilms are a significant reservoir of antibiotic-resistant *Campylobacter* spp. even without antibiotic selective pressure¹. Thermotolerant *Campylobacter* spp. may survive in the slaughterhouses and be a contamination source to poultry from different chains. Therefore, future studies of biofilm in slaughterhouses should provide tools for both the *Campylobacter* spp. epidemiology in order to implement appropriate intervention measures to eliminate thermotolerant *Campylobacter* spp. from slaughterhouse surfaces.

Previous studies conducted by our group allowed us to conclude that: a) thermotolerant *Campylobacter* spp. was present in a high proportion in the different stages of the poultry meat chain, especially in broilers in flock and carcasses in the slaughterhouse and the retail market¹⁹ and b) a significant proportion of the thermotolerant *Campylobacter* spp. isolated was resistant to the antibiotics commonly used in human medicine and most of them showed multiresistance patterns²⁰. These findings show that poultry meat is a significant source of *Campylobacter* spp. contamination and that the consumers in Argentina are exposed to different strains of thermotolerant *Campylobacter* spp. which constitute a significant public health problem. Further studies should be conducted with the aim to establish the epidemiological link between the strains isolated from the poultry meat chain and human isolates. Furthermore, MLST typing could be useful for comparing our isolates with those circulating in other countries.

Conclusions

The present work is the first in Argentina to reveal that thermotolerant *Campylobacter* spp. shows high genetic diversity. Evidences for transmission through the food chain and cross contamination in slaughterhouses were found. This collective evidence should be considered as the scientific basis to implement risk management measures to protect the public health. It is clear that poultry are colonized on the farm and further studies should be conducted to evaluate the importance of different fomites and vectors and therefore, to understand the epidemiology of this food-borne pathogen.

Ethical responsibilities

Protection of people and animals. The authors state that the procedures followed conformed to the ethical standards of the responsible human experimentation committee and in agreement with the World Medical Association and the Declaration of Helsinki.

Confidentiality of data. The authors state that no patient data appears in this article.

Right to privacy and informed consent. The authors state that no patient data appears in this article.

Conflict of interest

The authors declare that they have no ‘‘conflicts of interest’’.

Acknowledgments

LSF, LPS, MVZ and MLS are Research Career Member from the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina). JEB, DMA, ER, APB, ARS and JAZ are doctoral fellow from CONICET (Argentina). The authors would like to thank Dr. Gerardo Leotta (Instituto de Genética Veterinaria (IGEVET-CONICET/UNLP) Facultad de Ciencias Veterinarias (UNLP) for the use of the Bionumerics software. We also thank Dr. Horacio Terzolo and Alejandra Velilla (National Institute of Agricultural Technology-EEA Balcarce, Argentina) for their assistance and cooperation in the training of our working group in bacteriological techniques of thermotolerant *Campylobacter*. This study was funded by CONICET (PIP 11220120100481CO) and SENASA (Research, Transfer and Communication – Health, Safety and Quality Agrifood Research Group for training. Collective award from SENASA).

References

1. Bae J, Oh E, Jeon B. Enhanced transmission of antibiotic resistance in *Campylobacter jejuni* biofilms by natural transformation. *Antimicrob Agents Chemother*. 2014;58:7573–5.
2. Bolton FJ, Coates D. Development of a blood-free *Campylobacter* medium: screening tests on basal media and supplements,

- and the ability of selected supplements to facilitate aerotolerance. *J Appl Bacteriol.* 1983;54:115–25.
3. Brown HL, Reuter M, Salt LJ, Cross KL, Betts RP, van Vliet AH. Chicken juice enhances surface attachment and biofilm formation of *Campylobacter jejuni*. *Appl Environ Microbiol.* 2014;80:7053–60.
 4. Damjanova I, Jakab M, Farkas T, Mészáros J, Galántai ZS, Turcsányi I, Bistyák A, Juhász A, Pászti J, Kiss I, Kardos G. From farm to fork follow-up of thermotolerant campylobacters throughout the broiler production chain and in human cases in a Hungarian county during a ten-month period. *Int J Food Microbiol.* 2011;150:95–102.
 5. EFSA. Analysis of the baseline survey on the prevalence of *Campylobacter* in broiler batches and of *Campylobacter* and *Salmonella* on broiler carcasses in the EU, 2008, Part A: *Campylobacter* and *Salmonella* prevalence estimates. *EFSA J.* 2010;8:1503.
 6. Fuentes LS. Prevalencia y perfiles de sensibilidad en cepas de *Campylobacter* spp. aisladas de diarreas en Córdoba, Argentina. *Gastroenterology.* 2010;12:2–5.
 7. Hunter SB, Vauterin P, Lambert-Fair MA, Van Duynne MS, Kubota K, Graves L, Wrigley D, Barrett T, Ribot E. Establishment of a universal size standard strain for use with the PulseNet standardized pulsed-field gel electrophoresis protocols: converting the national databases to the new size standard. *J Clin Microbiol.* 2005;43:1045–50.
 8. Lior H. New, extended biotyping scheme for *Campylobacter jejuni*, *Campylobacter coli* and *Campylobacter lariidis*. *J Clin Microbiol.* 1984;20:636–40.
 9. Ma L, Wang Y, Shen J, Zhang Q, Congming W. Tracking *Campylobacter* contamination along a broiler chicken production chain from the farm level to retail in China. *Int J Food Microbiol.* 2014;181:77–84.
 10. Melero B, Juntunen P, Hanninen ML, Jaime I, Rovira J. Tracing *Campylobacter jejuni* strains along the poultry meat production chain from farm to retail by pulsed-field gel electrophoresis and the antimicrobial resistance of isolates. *Food Microbiol.* 2012;32:124–8.
 11. Newell DG, Fearnley C. Sources of *Campylobacter* colonization in broiler chickens. *Appl Environ Microbiol.* 2003;69:4343–51.
 12. Ribot EM, Fitzgerald C, Kubota K, Swaminathan B, Barrett TJ. Rapid pulsed-field gel electrophoresis protocol for subtyping of *Campylobacter jejuni*. *J Clin Microbiol.* 2001;39:1889–94.
 13. Scheppard SK, Dallas JF, Strachan NJC, MacRae M, McCarthy ND, Wilson DJ, Gormley FJ, Falush D, Ogden IE, Maden MCJ, Forbes KJ. *Campylobacter* genotyping to determine the source of human infection. *Clin Infect Dis.* 2009;48:1072–8.
 14. Terzolo HR, Lawson GHK, Angus KW, Snodgrass DR. Enteric *Campylobacter* infection in gnotobiotic calves and lambs. *Res Vet Sci.* 1987;43:72–7.
 15. Vandamme P, van Doorn LJ, Rashid STA, Quint WGV, Plas Jvd, Chan VL, On SLW. *Campylobacter hyoilei* Alderton et al. 1995 and *Campylobacter coli* Veron and Chatelain 1973 are subjective synonyms. *Int J Syst Bacteriol.* 1997;47:1055–60.
 16. Wassenaar TM, Newell DG. Genotyping of *Campylobacter* spp. *Appl Environ Microbiol.* 2000;66:1–9.
 17. Wieczorek K, Denis E, Osek J. Comparative analysis of antimicrobial resistance and genetic diversity of *Campylobacter* from broilers slaughtered in Poland. *Int J Food Microbiol.* 2015;210:24–32.
 18. Zaidi MB, Calva JJ, Estrada-García MT, León V, Vázquez G, Figueroa G, López E, Contreras J, Abbott J, Zhao S, McDermott P, Tollefson L. Integrated food chain surveillance system for *Salmonella* spp. in Mexico. *Emerg Infect Dis.* 2008;14:429–35.
 19. Zbrun MV, Romero-Scharpen A, Olivero C, Rossler E, Soto LP, Rosmini MR, Sequeira GJ, Signorini ML, Frizzo LS. Occurrence of thermotolerant *Campylobacter* spp. at different stages of the poultry meat supply chain in Argentina. *N Z Vet J.* 2013;61:337–43.
 20. Zbrun MV, Olivero C, Romero-Scharpen A, Rossler E, Soto LP, Astesana DM, Blajman JE, Berisvil A, Signorini ML, Frizzo F L.S. Antimicrobial resistance in thermotolerant *Campylobacter* isolated from different stages of the poultry meat supply chain in Argentina. *Food Control.* 2015;57:136–41.
 21. Zorman T, Heyndrickx M, Uzunović-Kamberović S, Smole Mozina S. Genotyping of *Campylobacter coli* and *C. jejuni* from retail chicken meat and humans with campylobacteriosis in Slovenia and Bosnia and Herzegovina. *Int J Food Microbiol.* 2006;110:24–33.
 22. Zweifel C, Scheu KD, Keel M, Renggli F, Stephan R. Occurrence and genotypes of *Campylobacter* in broiler flocks other farm animals, and the environment during several rearing periods on selected poultry farms. *Int J Food Microbiol.* 2008;125:182–7.