



BRIEF REPORT

Persistence of the same genetic type of *Mycoplasma hyopneumoniae* in a closed herd for at least two years

Florencia Rebaque^a, Pablo Camacho^{a,c}, Julián Parada^{a,c}, Paula Lucchesi^{b,c}, Arnaldo Ambrogi^a, Pablo Tamiozzo^{a,*}

^a Departamento de Patología Animal, Facultad de Agronomía y Veterinaria, Universidad Nacional de Río Cuarto, Ruta 36 Km 601, Río Cuarto, Córdoba CP 5800, Argentina

^b CIVETAN, UNCPBA, CICPBA, Departamento de Sanidad Animal y Medicina Preventiva, Facultad de Ciencias Veterinarias, Tandil, Buenos Aires, Argentina

^c CONICET, Argentina

Received 24 February 2017; accepted 21 May 2017

Available online 20 October 2017

KEYWORDS

Swine;
Mycoplasma hyopneumoniae;
Typing;
Multiple Locus
Variable-number
tandem repeat
Analysis;
Persistence;
Polymerase chain
reaction

PALABRAS CLAVE

Cerdo;
Mycoplasma hyopneumoniae;
Tipificación;

Abstract Two cross-sectional studies were carried out in 2013 and 2015 monitoring for *Mycoplasma hyopneumoniae* presence in a swine farm. In these studies, the genetic diversity of *M. hyopneumoniae* was assessed in clinical specimens using a Multiple Locus Variable-number tandem repeat Analysis (MLVA) targeting P97 R1, P146 R3 and H4 loci. The samples from August 2015 showed the MLVA profile prevalent in June 2013, therefore it can be concluded that a same genetic type of *M. hyopneumoniae* can persist for at least two years in a closed herd. In addition, the nested PCR reactions implemented in this study showed to be useful for MLVA typing in non-invasive clinical samples.

© 2017 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Persistencia del mismo tipo genético de *Mycoplasma hyopneumoniae* por al menos 2 años en una piara sin reposición externa de animales

Resumen Dos estudios transversales fueron realizados en los años 2013 y 2015 monitorizando la presencia de *Mycoplasma hyopneumoniae* en una piara. En esos estudios la diversidad genética de *M. hyopneumoniae* fue evaluada a partir de muestras clínicas utilizando un análisis multilocus de regiones repetidas en tándem (MLVA) de los loci P97 R1, P146 R3 y H4. Las muestras colectadas en agosto del 2015 mostraron el perfil de MLVA prevalente en junio del 2013, por lo tanto, se puede concluir que el mismo tipo genético de *M. hyopneumoniae* puede

* Corresponding author.

E-mail address: ptamiozzo@ayv.unrc.edu.ar (P. Tamiozzo).

Análisis Multilocus de
Regiones Repetidas
en Tándem;
Persistencia;
Reacción en cadena
de la polimerasa

persistir por al menos 2 años en una piara sin reposición externa de animales. Además, las reacciones de PCR anidadas implementadas en este estudio mostraron ser útiles para la tipificación por MLVA a partir de muestras clínicas no invasivas.

© 2017 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Mycoplasma hyopneumoniae is the etiological agent of enzootic pneumonia, an important and highly prevalent respiratory disease affecting pigs in almost all swine-producing areas around the world. Certain epidemiological aspects, such as persistence and genetic diversity of *M. hyopneumoniae* strains circulating within herds, have been addressed in some studies^{9,11}, however, most of the longitudinal studies were carried out over a period of less than one year.

Even though there is no molecular marker able to discriminate between high and low virulence *M. hyopneumoniae* strains, the knowledge of the persistence of certain genotypes of the agent might be an important indicator of the severity of the disease when comparing clinical signs and lung lesions.

The aim of this study was to determine genotypes of *M. hyopneumoniae* present in pigs in a commercial farm on two sampling occasions, two years apart, to assess persistence of the strains.

The study was performed according to the international guidelines of the Council for International Organizations of Medical Sciences (CIOMS).

Typing of *M. hyopneumoniae* was performed by Multiple-Locus Variable-Number Tandem-Repeat Analysis (MLVA) using DNA samples positive for *M. hyopneumoniae* by a nested PCR (nPCR) described by Calsamiglia et al.¹ P97 R1, P146 R3 and H4 loci were analyzed in 35 DNA samples obtained from nasal swab (NS) and bronchoalveolar lavage (BAL) specimens of sows and pigs. All samples were collected in two cross-sectional studies performed on June 2013 (27 specimens) and on August 2015 (8 specimens) in order to monitor enzootic pneumonia control measures in a two-site, 1300-sow, commercial wean-to-finish farm. The control measures applied since June 2013 included sows vaccination against *M. hyopneumoniae* (Porcilis M hyo, MSD Animal Health) and strategic in-feed antibiotic treatment (tiamulin at 112 ppm and chlortetracycline at 300 ppm) to pigs at 6 weeks of age during a 2-week period.

For MLVA, VNTR loci P97 R1 and H4 were amplified by nPCR format combining primers designed in previous studies. The primers reported by de Castro et al.², were used for the first round of amplification, and those reported by Vranckx et al.¹⁰, for the second round. The P146 R3 locus was amplified using a nPCR previously reported by Tamiozzo et al.^{8,9} After amplification, the amplicons were purified (Puriprep-GP Kit, Inbio Highway), quantified and sequenced (ABI 3130xl; Applied Biosystems) with the primers described by Vranckx et al.¹⁰, for P97 and H4, and those of Mayor

et al.⁶, for P146 R3. The number of tandem repeats was determined viewing the sequences in the BioEdit software.

All the samples from June 2013 and August 2015 characterized by MLVA showed the same 11 repeat units in P97 R1 (repeat motif encoding AAKP[EV]), and 8 repeat units for H4 (QTTQKD). Regarding P146 R3, tandem repeats with 26, 27 and 28 units (S) were identified in samples from June 2013, whereas all typed samples from August 2015 showed 26 repeat units (Table 1).

Considering that all the samples from 2015 analyzed in the present study showed the same 3-loci profile as most samples taken in 2013, persistence of a *M. hyopneumoniae* strain over this period is highly suggested. The high discriminatory power reported by Dos Santos et al.³, for two of these loci (P97 R1 and P146 R3) gives support to that conclusion. Furthermore, the MLVA approach was reinforced in the present study by adding another polymorphic locus (H4).

Persistence of same MLVA type of *M. hyopneumoniae* over time is not an unexpected result due to the fact that it was a closed herd and the presence of only one distinct strain within the herds and low genetic diversity at herd-level has already been reported¹¹. The access of another similar genetic strain through personnel, fomites or airborne transmission is highly improbable, because the presence of *M. hyopneumoniae* strain with 26 polyserine repeat motif of *p146* gene had never been found in our country.

Apparently, the strain circulated among the pig population without any pressure of selection in the genome, or at least in the loci that were analyzed, in spite of vaccination and antibiotic treatment of the pigs. This may result in the persistence of the same genetic type. In this way a recent study has demonstrated the stability of P97 R1 and P146 R3 as *in vivo* as *in vitro*³.

Kuhnert and Overesch⁴ have shown, using MLST typing, the persistence of one *M. hyopneumoniae* strain in some farms, causing recurrent outbreaks, relating this fact to unsuccessful sanitation measures. In addition, it has been determined that *M. hyopneumoniae* can remain in pig organs even after antibiotic treatment⁵. More in-depth studies are necessary to determine the role of these MLVA loci in the development of mechanisms of persistence and evasion of the host immune system.

The VNTR PCR tests implemented in a nested format proved to be useful to type *M. hyopneumoniae* from clinical specimens without killing animals or performing invasive samplings. Considering that the 3-loci MLVA profile detected in the present study cannot be easily compared with results from other studies because of not all these three loci were analyzed or the number of repeat units present were not

Table 1 *Mycoplasma hyopneumoniae* detection and typing using DNA extracted from nasal swabs and broncho-alveolar lavages (BAL) of sows and pigs at different ages and at slaughter^a

			nPCR for <i>M. hyopneumoniae</i> detection (positive samples/analyzed samples)			MLVA loci		
						P97 R1	P146 R3	H4
						Number of repeat units (number of samples)	Number of repeat units (number of samples)	Number of repeat units (number of samples)
June 2013	Sows (nasal swabs)	4/30		None positive	26 (1)		None positive	
	Pigs (nasal swabs)	17/105		11(4)	26 (8), 27 (1), 28 (1)		8 (1)	
	Slaughter (BAL)	6/30		11(4)	26 (6)		8 (4)	
August 2015	Pigs (nasal swabs)	3/45		11(1)	None positive		8 (1)	
	Slaughter (BAL)	5/30		11(5)	26 (5)		8 (5)	

^a A two-site, 1300-sow, commercial wean-to-finish farm was monitored looking for *M. hyopneumoniae* in cross-sectional studies carried out in June 2013 and in August 2015. In these studies, some nasal swabs and broncho-alveolar lavages of sows and pigs had rendered nPCR positive results. Typing was performed to assess the genetic diversity from some of these clinical specimens. The number repeat units of *p97*, *p146* and *H4* genes was determined by a nested-polymerase chain reactions as following: For P146 using as outer primers those described by Tamiozzo et al.⁸, and as inner primers those described by Mayor et al.⁶ Sequencing was performed using this last primer pair. For P97 and H4 using as outer primers those described by de Castro et al.², and as inner primers those described by Vranckx et al.¹⁰ Sequencing was performed using this last primer pair.

determined^{3,6,7}; we consider that the use of a standardized nomenclature for MLVA loci and alleles of *M. hyopneumoniae* is highly needed to aid comparison of results from different studies.

To the best of our knowledge, this is the first report showing the persistence of a same genetic type of *M. hyopneumoniae* for such a long time period. We are aware that knowledge of *M. hyopneumoniae* persistence and genetic variability are essential to understand some epidemiological aspects of enzootic pneumonia and to develop and evaluate strategies for disease control.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this investigation.

Confidentiality of data. The authors declare that they have followed the protocols of their work centre on the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appears in this article.

Funding

This study was financially supported in part by PICT 0442/2015 FONCyT-ANPCyT-MinCyT, República Argentina and PPI 2016–2018, 188/2016- UNRC. F. Rebaque was holder of an “Incentivo a las Vocaciones Científicas” scholarship

during the development of this study (CIN-SPU-Ministerio de Educación), República Argentina.

Conflict of interest

None reported.

References

1. Calsamiglia M, Pijoan C, Trigo A. Applications of a nested-polymerase chain reaction assay to detect *Mycoplasma hyopneumoniae* from nasal swabs. *J Vet Diag Inv.* 1999;1:246–51.
2. de Castro LA, Rodrigues Pedrosa T, Kuchiishi SS, Ramenzoni M, Kich JD, Zaha A, Henning Vainstein M, Bunselmeyer Ferreira H. Variable number of tandem aminoacid repeats in adhesion-related CDS products in *Mycoplasma hyopneumoniae* strains. *Vet Microbiol.* 2006;116:258–69.
3. Dos Santos LF, Sreevatsan S, Torremorell M, Moreira MA, Sibila M, Pieters M. Genotype distribution of *Mycoplasma hyopneumoniae* in swine herds from different geographical regions. *Vet Microbiol.* 2015;175:374–81.
4. Kuhnert P, Overesch G. Molecular epidemiology of *Mycoplasma hyopneumoniae* from outbreaks of enzootic pneumonia in domestic pig and the role of wild boar. *Vet Microbiol.* 2014;174:261–6.
5. Le Carrou J, Laurentie M, Kobisch M, Gautier-Bouchardon AV. Persistence of *Mycoplasma hyopneumoniae* in experimentally infected pigs after marbofloxacin treatment and detection of mutations in the *parC* gene. *Antimicrob Agents Chemother.* 2006;50:1959–66.
6. Mayor D, Zeeh F, Frey J, Kuhnert P. Diversity of *Mycoplasma hyopneumoniae* in pig farms revealed by direct molecular typing of clinical material. *Vet Res.* 2007;38:391–8.

7. Stakenborg T, Vicca J, Maes D, Peeters J, de Kruif A, Haesebrouck F, Butaye P. Comparison of molecular techniques for the typing of *Mycoplasma hyopneumoniae* isolates. *J Microbiol Methods*. 2006;66:263–75.
8. Tamiozzo P, Lucchesi PMA, Ambrogi A. Diversidad genética de *Mycoplasma hyopneumoniae* en granjas porcinas de Argentina [*Mycoplasma hyopneumoniae* genetic diversity in swine farms from Argentina]. *In Vet*. 2011;13:27–35.
9. Tamiozzo P, Lucchesi PMA, Ambrogi A. Monitoring for *Mycoplasma hyopneumoniae* before and after a partial depopulation program using a typing scheme based on the polyserine repeat motif of *p146*. *J Swine Health Prod*. 2013;21:309–12.
10. Vranckx K, Maes D, Calus D, Villarreal I, Pasmans F, Haesebrouck F. Multiple-locus variable-number tandem-repeat analysis is a suitable tool for differentiation of *Mycoplasma hyopneumoniae* strains without cultivation. *J Clin Microbiol*. 2011;49:2020–3.
11. Vranckx K, Maes D, Sacristán R, del P, Pasmans F, Haesebrouck F. A longitudinal study of the diversity and dynamics of *Mycoplasma hyopneumoniae* infections in pig herds. *Vet Microbiol*. 2012;156:315–21.