



MICROBIOLOGICAL IMAGE

Infectivity by nematode–bacteria association on the potato weevil *Phyrdenus muriceus*



Infectividad de una asociación nematodo-bacteria en el gorgojo de la papa *Phyrdenus muriceus*

Daiana P. Eliceche, José M. Rusconi*, Matías N. Rosales, Augusto Salas, María F. Achinelly

Laboratorio de Nematodos de Vida Libre y de Importancia Agro-económica, Centro de Estudios Parasitológicos y de Vectores (CEPAVE)-CCT-La Plata-CONICET-UNLP, CIC, Boulevard 120 S/N e/ 61 y 64, 1900 La Plata, Buenos Aires, Argentina

Received 2 October 2018; accepted 26 November 2018
Available online 9 January 2020

Bacteria of the genus *Photorhabdus* (*Enterobacteriaceae*) are motile, gram negative, symbiotically associated with infective juveniles of entomopathogenic nematodes of the genus *Heterorhabditis* (*Heterorhabditidae*). Nematodes retain the *Photorhabdus* bacterial cells in their intestine and transport them into an insect host⁶. Upon locating an insect host, the nematode of the genus *Heterorhabditis* penetrates through natural openings (mouth, anus, spiracles)¹, or directly into the hemocoel via the integument, subsequently releasing bacteria into the hemolymph⁴. Once in the hemolymph, bacterium of the genus *Photorhabdus* begins to multiply simultaneously releasing toxins virulent enough to kill the insect within 24 h². *Phyrdenus muriceus* (Coleoptera: Curculionidae) is a pest of cultivated Solanaceae affecting egg-plant, potato and tomato. Considerable damages on these crops are produced by this pest affecting both the aerial and underground parts of the plants^{3,5}. Infectivity of the entomopathogenic complex *Heterorhabditis*

bacteriophora and the symbiotic bacterium *Photorhabdus luminescens* isolated from soil samples collected from an orchard of La Plata was tested against pupae of *P. muriceus*. This stage was considered to be a suitable target to control in the field due to its permanence in the soil. Plastic containers (8.5 cm diameter × 5.5 cm height) were used with sterile soil forming a 2 cm high layer. Pupae were added to the soil surface and exposed to a concentration of 5000 infective juveniles (IJs)/insect. Seventy percent (70%) mortality (n = 10) occurred at 48 h after infection. In [Figure 1](#), we can observe nematodes inside the body of the pupae (B and C). The cadavers showed the typical red wine coloration due to the release of the endosymbiotic bacteria in the hemocoel ([Figs. 1B–D](#)) unlike non-parasitized pupae ([Fig. 1A](#)). The nematode–bacterium complex was effective to parasitize and kill *P. muriceus* pupae. Future studies should be carried out to determine its use as biological control for this horticultural pest.

* Corresponding author.

E-mail address: rusconi@cepave.edu.ar (J.M. Rusconi).



Figure 1 Image of pupae of *Phyrdenus muriceus* taken with an Olympus DP-71 camera. (A) Pupa with no infection. (B) Dorsal view of infected pupa with nematodes inside the body (arrows). (C) Ventral view of infected pupa with nematodes inside the body (arrows). (D) Emerging nematodes from the lesion of the tegument (arrow).

Conflict of interest

The authors declare that they have no conflicts of interest.

References

1. Boemare N. Interactions between the partners of the entomopathogenic bacterium nematode complexes, *Steinrenema-Xenorhabdus* and *Heterorhabditis-Photorhabdus*. Brill Academic Publishers; 2002. p. 601–3.
2. Chiche TA, Ensing JC. For the insect pathogen *Photorhabdus luminescens* which end of a nematode is out? *Appl Environ Microbiol*. 2003;60:1890–7.
3. Espul JC, Magistretti G. Biotecnología de “gorgojo del tomate” *Phyrdenus muriceus* (Germ.) y su control en la Provincia de Mendoza. *Rev Investig Agropecu (AR) Ser 5 Patol Veg*. 1969;6:95–117.
4. Forst S, Dowds B, Boemare N, Stackenbrandt E. *Xenorhabdus* and *Photorhabdus* spp: bugs that kill bugs. *Annu Rev Microbiol*. 1997;51:47–72.
5. Novo RJ, Viglianco A, Vaudagna E. Efectos de insecticidas sobre el gorgojo de la papa, *Phyrdenus muriceus* (Germ.) (Coleoptera: Curculionidae). *AgriScientia*. 2002;19:3–10.
6. Peat S, French-Constant R, Waterfield N, Marokházi J, Fodor A, Adams B. A robust phylogenetic framework for the bacterial genus *Photorhabdus* and its use in studying the evolution and maintenance of bioluminescence: a case for *16S*, *gyrB*, and *glnA*. *Mol Phylogenet Evol*. 2010;57:728–40.