



Original

Yeasts isolation from bovine mammary glands under different mastitis status in the Mexican High Plateau

Carolina Segundo Zaragoza^{a,*}, Roberto Arnulfo Cervantes Olivares^a, Andrés Ernesto Ducoing Watty^b, Alejandro de la Peña Moctezuma^a, Lourdes Villa Tanaca^c

^a Departamento de Microbiología e Inmunología, Laboratorio de Micología Veterinaria, Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, Mexico

^b Departamento de Producción Animal Rumiantes, Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, Mexico

^c Departamento de Microbiología, Laboratorio de Genética Microbiana, Escuela Nacional de Ciencias Biológicas, IPN, Mexico

ARTICLE INFO

Article history:

Received 18 October 2010

Accepted 17 January 2011

Available online 22 February 2011

Keywords:

Candida

Bovine

Milk

Mastitis

Yeast

ABSTRACT

Background: The mastitis subclinical and clinical in cows caused by fungi has been increased specially by yeast of the genus *Candida*.

Objective: To identify what yeasts were present in milk samples obtained from mammary glands of healthy cows, and others suffering subclinical or clinical mastitis.

Methods: From a total of 1,095 milk samples 342 were from mammary glands of healthy dairy cows, 383 with subclinical mastitis, and 370 with clinical mastitis, were taken, in the states of Querétaro, Hidalgo, Puebla and Mexico City (Distrito Federal) in the so called Mexican High Plateau. The clinical status of the mammary glands was determined by clinical examination and the California Mastitis Test. Yeasts identification was carried out by morphology and biochemical methods.

Results: Twenty different species of *Candida* were identified out of 282 (25.75%) milk samples. The most frequently identified species in the healthy cows and cows with clinical mastitis groups were *Candida glabrata* and *Candida krusei*. On the other hand, samples from the subclinical mastitis group showed a diversity of *Candida* species, including *Candida zeylanoides*, *Candida norvegica*, *Candida viswanathii*, *Candida guilliermondii*, and *Candida tropicalis*. *Candida albicans* was isolated only in 11 (3.9%) samples from the clinical and subclinical mastitis groups.

Conclusions: These results suggest the possible role that *Candida* species other than *C. albicans* may play in mycotic mastitis in cows.

© 2010 Revista Iberoamericana de Micología. Published by Elsevier España, S.L. All rights reserved.

Aislamiento de levaduras de glándulas mamarias con diferentes presentaciones de mastitis en el Altiplano Mexicano

RESUMEN

Antecedentes: Las mastitis subclínicas y clínicas en bovinos por hongos se han incrementado principalmente por levaduras del género *Candida*.

Objetivo: Conocer las levaduras presentes en la leche de glándulas mamarias de bovinos clínicamente sanos, y de aquellos con mastitis subclínica y clínica.

Método: Se evaluó la presencia de levaduras en 1.095 muestras de leche de 342 glándulas mamarias sanas, 383 con mastitis subclínica y 370 con mastitis clínica, de los estados de Querétaro, Hidalgo, Puebla y la ciudad de México, Distrito Federal, que forman parte del Altiplano Mexicano. El estado sanitario de las glándulas mamarias se determinó por examen clínico y la prueba de California. La identificación de levaduras fue realizada por métodos morfológicos y bioquímicos.

Resultados: Se identificaron 20 especies diferentes del género *Candida* a partir de 282 (25,75%) de las muestras de leche. Las especies encontradas con mayor frecuencia en los bovinos sanos y con

Palabras clave:

Candida

Bovinos

Leche

Mastitis

Levadura

* Corresponding author.

E-mail address: c.segund@yahoo.com.mx (C. Segundo Zaragoza).

mastitis clínica fueron *Candida glabrata* y *Candida krusei*. El grupo de las muestras con mastitis subclínica mostró una diversidad de especies de *Candida*, incluidas *Candida zeylanoides*, *Candida norvegica*, *Candida viswanathii*, *Candida guilliermondii* y *Candida tropicalis*. *Candida albicans* fue aislada solo en 11 (3,9%) de las muestras de mastitis clínica (6) y subclínica (5).

Conclusiones: Estos resultados sugieren el posible papel de otras especies de *Candida* diferentes a *C. albicans* como causantes de mastitis micótica.

© 2010 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Bovine mastitis is a disease caused by a wide variety of microorganisms that causes large economical losses and damages to the dairy industry by decreasing milk production and through increased costs of antibiotic treatments and culling⁶. In most cases, bacteria are recognized as the primary pathogens while fungi, particularly yeasts, have been regarded as secondary mastitis pathogens. Yeasts are considered opportunistic pathogens which colonize the cows' udder. The use and abuse of antibacterial drugs, treatment with contaminated antibiotic solutions, as well as syringes, or other materials brought in contact with the mammary gland may favor yeast colonization of cows udders^{4,6,12,13}.

Different fungi have been reported as a cause of mycotic mastitis, such as *Aspergillus fumigatus*, *Aspergillus terreus*, *Candida* spp., *Cephalosporium* spp., *Coccidioides* spp., *Cryptococcus neoformans*, *Geotrichum candidum*, *Histoplasma* spp., *Mucor* spp., *Rhizopus* spp., *Torulopsis* spp., and *Trichosporon* spp. *Candida* species have been regarded in subclinical and clinical mastitis, being the most reported species *Candida albicans*, *Candida krusei*, *Candida rugosa*, and *Candida guilliermondii*^{1,4,16,26}. Furthermore, other species of *Candida*, as well as other fungi such as *Cryptococcus* spp., *Rhodotorula* spp., *Trichosporum cutaneum*, *Aureobasidium pullulans* and *Pichia ohmeri*, have also been isolated from the milk of healthy glands^{5,18,21,27}.

The purpose of this study is to report the presence of *Candida* species isolated from cows with different mammary gland health status in the Mexican High Plateau.

Materials and methods

One thousand and ninety-five milk samples were collected from Holstein-Friesian cows kept under intensive production conditions in the states of Queretaro (n = 216), Hidalgo (n = 697), Puebla (n = 52) and Distrito Federal (n = 130) in central Mexico, each sample corresponds to a different animal and from single quarter, in the clinical cases the sample was taken from the most affected quarter. The samples were obtained at convenience, from two-year and older cows, with different mammary glands health status: 342 samples from cows with healthy mammary glands, 383 from cows with subclinical mastitis as determined by the California Mastitis Test (CMT), and 370 from cows diagnosed with clinical mastitis¹⁵. Clinical mastitis was defined by: swelling, reduced milk flow, and abnormal milk appearance (watery to viscous with clots varying from gray-white to yellowish). Additionally, other signs of infection such as fever, inappetence, ataxia, and depression were also considered⁸. CMT was used to identify subclinical mastitis on mammary gland of the cows. For this study, milk samples from glands affected with subclinical mastitis were included when the reaction to CMT was at least grade 1, corresponding with an appearance of viscous milk that does not adhere to the bottom of the CMT plate, and correlates to 400,000–1,500,000 somatic cells/ml²⁵.

Sampling was carried out prior to milking, after washing the udder with soap, drying and disinfecting it with the disinfectant on use in the farm at the moment of sampling, and drying again. All farms included in the study used either 2% iodine or 70% alcohol, both known to have a fungicidal effect^{2,3,28}. Fifteen ml of milk

from each sampled gland were collected in sterile screw cap containers. Samples were refrigerated (4 °C) during transportation to the laboratory of Mycology at the Faculty of Veterinary Medicine of the National Autonomous University of Mexico (UNAM), and kept at 4 °C until processing (no longer than 24 h after collection). Milk samples were incubated during 15 min at 25 °C and homogenized by vigorous shaking; 0.5 ml of the homogenized milk were inoculated in 4.5 ml of Sabouraud dextrose broth (Difco™), at pH of 3.5, and then incubated at 37 °C during 10 days¹⁷. Thereafter, 50 µl of each broth culture was plated on Sabouraud dextrose agar (Difco™), with chloramphenicol (Merck™) (0.05 mg/ml). Inoculated plates were then incubated at 37 °C and examined daily for colonies for five days.

Yeast identification was performed taking into consideration morphological characteristics, like formation of chlamydoconidium, pseudohyphae and germinal tube development. Additional characteristics were also evaluated, such as growth in the presence of 0.1% cyclohexamide (Sigma™), acidic pH tolerance, urea hydrolysis and carbohydrates assimilation and/or fermentation, accordingly to the methodology described by Barnnet and Payne, and Kurtzman and Fell^{9,10,23,29}. Data obtained in this study were evaluated using cross-tabulation analyses through Chi square distribution²⁴ with the statistical package JMP software, version 5.1.

Results

Out of the 1.095 milk samples analyzed, 282 (25.75%) were positive to yeast isolation (Table 1), obtaining in Queretaro 48/216, Hidalgo 119/697, Puebla 23/52 and Distrito Federal 92/130. Yeast isolates were obtained in 23.39% (80/342) out of milk samples from healthy mammary glands), 9.92% (38/383) from the subclinical mastitis group and 43.27% (164/370) from mammary glands with clinical mastitis. Significant differences (Chi square = 121.7, df = 2.1092, P < 0.01) were found among the three groups for the yeast isolation frequencies. Only the genus *Candida* was identified in the samples that were positive to yeast isolation.

Twenty *Candida* species were identified, being the most frequently found *Candida glabrata* 30.14% (85/282), and *C. krusei* 27.65% (78/282) (Table 1).

Out of the 342 milk samples analyzed from healthy mammary glands, *Candida* isolates were obtained in 23.39% (80/342) from which 12 different species were identified. The species most frequently found was *C. glabrata* with 58.75% (47/80) isolates, followed by *C. krusei* with 11.25% (9/80) and *Candida viswanathii* with 7.5% (6/80). From the total isolates obtained, 8.75% (7/80) isolates could not be identified at to the species level, being considered as «*Candida*-like» according their growth characteristics, morphology and microscopy characteristics.

From the 383 samples analyzed from mammary glands with subclinical mastitis, only 9.92% (38/383) *Candida* isolates were obtained, and identified 15 different species. In this group, the most frequent species found were *C. albicans* 13.15% (5/38), *C. krusei* 7.89% (3/38), and *Candida zeylanoides* 7.89% (3/38), being 34.21% (13/38) of the isolates considered as *Candida*-like species.

Table 1
Milk samples and *Candida* species isolated from mammary glands of healthy cows and cows with subclinical and clinical mastitis.

Candida species	Mastitis status of mammary glands			Frequency isolation	
	Healthy, n = 342	Subclinical, n = 383	Clinical, n = 370	n = 282	%
<i>Candida glabrata</i>	47	0	38	85	30.14
<i>Candida krusei</i>	9	3	66	78	27.65
<i>Candida albicans</i>	0	5	6	11	3.90
<i>Candida zeylanoides</i>	2	3	2	7	2.48
<i>Candida norvegica</i>	1	1	5	7	2.48
<i>Candida viswanathii</i>	6	1	0	7	2.48
<i>Candida kefir</i>	1	0	4	5	1.77
<i>Candida guilliermondii</i>	1	1	3	5	1.77
<i>Candida tropicalis</i>	1	1	3	5	1.77
<i>Candida parapsilosis</i>	0	0	4	4	1.41
<i>Candida intermedia</i>	2	0	2	4	1.41
Other <i>Candida</i> species*	3	10	4	17	6.02
<i>Candida</i> -like species	7	13	27	47	16.66
Total	80/342 (23.39%) ^a	38/383 (9.92%) ^b	164/370 (44.32%) ^c	282/1095 (25.75%)	

**Candida* species isolated in < 1.41%.

^{a,b,c} Different superscripts indicates statistical differences ($p < 0.01$) among groups for isolates frequencies.

In the 370 samples analyzed from mammary glands with clinical mastitis, 44.32% (164/370) *Candida* isolates were obtained, from which 13 different species were identified, being *C. krusei* the most frequently found with 40.24% (66/164) isolates, and *C. glabrata* with 23.17% (38/164). From these isolates, 16.46% (27/164) were classified as *Candida*-like species.

Other *Candida* species, such as *C. zeylanoides*, *Candida norvegica* and *C. viswanathii* were also found in 2.48% and *Candida kefir*, *C. guilliermondii*, *Candida tropicalis*, *Candida parapsilosis*, *Candida intermedia*, *Candida clausenii*, *Candida lusitanae*, *Candida macedoniensis*, *Candida brumptii*, *Candida slooffii*, *Candida lambica*, *Candida cantarelli*, *Candida lipolytica* and *Candida incommunis*, were also found at very low frequency ($\leq 1.7\%$).

Discussion

In this study, yeasts were isolated in the 25.75% of all samples analyzed, being *Candida* the only genus identified. *C. glabrata* was the predominant species, isolated in 58.75% of all positive samples (47/80) from the mammary glands from healthy cows, and in 23.17% of the *Candida* isolates (38/164) from those with clinical mastitis. The isolation of this *Candida* species from cattle affected with mastitis has also recently been reported in New Zealand by Williamson and di Menna²⁹. However, a previous report from Lagneau¹⁸, also showed the presence of *C. glabrata* in the mammary glands of healthy animals, in a rather small frequency (0.2%) when compared with our findings. In contrast, no *C. glabrata* was obtained from milk samples of our group of cows with subclinical mastitis. This could be partially explained by the fact that in subclinical mastitis it is common to find a greater number of bacteria competing for substrates; in addition, some bacterial metabolites are known as yeasts antagonists, restricting their development²⁰. It seems necessary to put more our attention to this species in order to establish its possible role in bovine mastitis.

The second most frequent species of *Candida* found in this study was *C. krusei*. This species was isolated in 11.25% of the healthy glands positive milk samples (9/80); in 7.89% of those with subclinical mastitis (3/38); and in 40.24% of those diagnosed with clinical mastitis (66/164). The presence of *C. krusei* in dairy cattle with mastitis has been recognized since the 1970s^{14,22}, although it has been more frequently reported since the late 1990s^{4,11,16,19}. There is, however, some discrepancy among the rates in which this yeast has been found. Whereas Casia⁴, reported in 2005 a 44.5% of *C. krusei* isolates from cows with mastitis, similarly to ours findings; Krukowski in 2000¹⁶, and Langoni in 1995¹⁹, found this species in 15.5% of mastitis samples and in less percentage than 2% in glands

with subclinical and clinical mastitis, respectively. Both findings are lower than our results. This might be a consequence effect of several factors, such as the abuse of intramammary antibiotic treatment or the use of home made antibiotic infusions for mastitis therapy¹⁶; the natural resistance of *C. krusei* to antimycotics, like fluconazol¹¹; yeast contaminated food or environment^{7,27}; and inadequate milking procedures²⁷, in addition to the presence of pathogenic strains of *C. krusei*²⁷. Recently, Williamson and di Menna reported a self-limiting clinical mastitis infection due to *C. krusei* in cows at calving and at up to 5 weeks post-partum²⁹. The isolation of *C. krusei* from healthy mammary glands might indicate the yeast as part of their normal microbiota and therefore, acting as an opportunistic pathogen in mastitis cases.

Regardless the health status of the mammary glands analyzed, *C. zeylanoides*, *C. norvegica*, *C. guilliermondii*, and *C. tropicalis* were always isolated. All these species have been shown to be involved in clinical mastitis. *C. zeylanoides* and *C. norvegica* are considered opportunistic pathogens, while *C. guilliermondii* and *C. tropicalis* are recognized as pathogens^{4,16}. However, our findings suggest that *C. norvegica* might act as a pathogen similar to *C. guilliermondii* and *C. tropicalis*, since the three species were found predominantly in milk from glands with clinical mastitis. In addition, our results on the isolation of *Candida parapsilosis*, only from animals with clinical mastitis, might support the probably pathogenicity of this *Candida* species^{16,18}.

The isolation of *C. albicans* only from glands from the groups of cows with clinical and subclinical mastitis (but not from healthy mammary glands), is consistent with the recognized pathogenicity of this species. The percentages in which this yeast was isolated in this study were similar to those reported in cows with mastitis by Langoni¹⁹, Krukowski¹⁶, and Casia⁴.

Our findings in relation to *C. viswanathii*, which was identified in 7.5% (6/80) of the healthy glands group, and in 2.6% (1/38) of the subclinical mastitis group, suggest the opportunistic pathogen role that this *Candida* species could be playing in this disease.

C. kefir, on the other hand, was identified in 2.43% (4/164) isolates in the clinical mastitis group, in partial agreement with previous reports in Europe that found *C. kefir* in milk samples at frequencies as high as 24.1%^{16,25}.

The isolation of numerous *Candida* non-*albicans* species from bovine milk is also in agreement with other reports^{4,16,25}, recognizing among others, antibacterial therapy without previous susceptibility tests as the principal predisposing factor for yeast infections. Furthermore, the wide distribution of yeasts in nature, e.g. soil, plants and water, to which cows are in contact with, might have implications regarding the presence of fungi in mammary

glands^{4,8,16}. Similar to our findings, there are some reports where *Candida* species other than *C. albicans*, have been isolated from milk of cows with mastitis^{2,3,8}.

Of the yeast isolates identified as *Candida*-like species, 16.66% (47/282) could not be further characterized by the biochemical methodology used in this study. At present, it is well known that the use of routine yeasts identification methods based on morphology and biochemical test are time-consuming, and they often fail to identify yeast species other than *C. albicans*. Alternative identification methods, such as PCR or RAPD-PCR might be necessary to support identification and further characterization of these mycotic agents¹⁷.

The presence of yeasts in udders of healthy cows, as well as in those of cows with mastitis, may be an important factor in the dissemination of disease. These agents might increase their presence due to handling of dairy cattle, bacterial mastitis treatment and prevention as well as inadequate milking practices. Furthermore, the presence of *Candida* non-*albicans* species that have gained ground as pathogens of the bovine mammary gland must be highlighted. For instance, the association of *C. glabrata* and *C. krusei* with clinical mastitis is clear, probably acting as opportunistic pathogens, considering that they can also be found in milk samples coming from healthy cows.

Although *C. albicans* has been reported as the most common species of yeast pathogen found, this study shows that *C. krusei* seems to be of interest as a probable cause of mastitis problems.

Financing

Programa de Apoyo a Proyectos de Investigación e Innovación Tecnológica (PAPIIT) grant IN 209908, Universidad Nacional Autónoma de México (UNAM). Consejo Nacional de Ciencia y Tecnología (CONACYT) 151676.

Conflict of interest

Authors have no conflict of interests.

Acknowledgements

The authors like to thank Sarahi Luna-Castro who collaborated with the biochemical characterization of the yeasts. We are particularly grateful to Dr. Cristina Escalante-Ochoa for her valuable review of this manuscript.

References

1. Aalbaek B, Stenderup J, Jensen HE. Mycotic and algal bovine mastitis in Denmark. *APMIS*. 1994;102:451–6.
2. Arenas R. *Micología Médica Ilustrada*. 2nd ed. Mexico: Interamericana-McGraw-Hill; 2003.
3. Barnett JA, Payne RW, Yarrow D, editors. *Yeast: Characteristics and Identification*. London: Cambridge University Press; 1990.
4. Casia dos Santos R, Marín JM. Isolation of *Candida* spp. from mastitic bovine milk in Brazil. *Mycopathologia*. 2005;159:251–3.
5. Costa EO, Gandra CR, Pires MF, Coutinho SD, Castillo W, Teixeira CM. Survey of bovine mycotic mastitis in dairy herds in the State of Sao Paulo, Brazil. *Mycopathologia*. 1993;124:7–13.
6. Costa EO, Ribeiro AR, Watanabe ET, Melvilla PA. Infectious bovine mastitis caused by environmental organisms. *J Vet Med B*. 1998;45:65–71.
7. Elad D, Shpigiel NY, Winkler M, Klinger I, Fuchs V, Saran A, et al. Feed contamination with *Candida krusei* as a probable source of mycotic mastitis in dairy cows. *JAVMA*. 1995;207:620–2.
8. Erskine RJ, Sears PM, Bartlett PC, Gage CR. Efficacy of postmilking disinfection with benzyl alcohol versus Iodophor in the prevention of new intramammary infections in lactating cows. *J Dairy Sci*. 1998;81:116–20.
9. Farnsworth RJ, Sorensen DK. Prevalence and species distribution of yeast in the mammary gland of dairy cows in Minnesota. *Can J Comp Med*. 1972;36:329–32.
10. Farnsworth RJ, Sorensen DK. The effect of penicillin, dihydrostreptomycin and prednisolone treatment of experimental *Candida krusei* infections of the mammary glands of dairy cattle. *Can J Comp Med*. 1974;39:340–8.
11. Freydiere AM, Guinet R, Boiron P. Yeast identification in the clinical microbiology laboratory: phenotypical methods. *Med Mycol*. 2001;39:9–33.
12. García ME, Blanco JL. Principales enfermedades fúngicas que afectan a los animales domésticos. *Rev Iberoam Micol*. 2000;17:S2–7.
13. Gibbony WJ, Catcott EJ, Smithcory JF. *Bovine Medicine and surgery*. USA American Veterinary Publications; 1970.
14. Holm C, Jepsen L, Larsen M, Jespersen L. Predominant microflora of downgraded Danish bula tank milk. *J Dairy Sci*. 2004;87:1151–7.
15. Kansas State University College of Veterinary Medicine [homepage on the Internet]. College of Veterinary Medicine, Kansas State University ©1996–2007. Bovine Mastitis [updated 31 Jan 2005]. Available from: www.vet.k-state.edu/depts/fahm/fam/mammary/mam.htm.
16. Krukowski H, Tietze M, Majewski T, Rozanski P. Survey of yeast mastitis in dairy herds of small-type farms in the Lublin region, Poland. *Mycopathologia*. 2000;150:5–7.
17. Kurtzman CP, Fell JW, editors. *The Yeast: a Taxonomic Study*. Amsterdam: Elsevier Science; 1998.
18. Lagneau PE, Lebtani K, Swinne D. Isolation of yeast from bovine milk in Belgium. *Mycopathologia*. 1996;135:99–102.
19. Langoni H, Domingues PF, Chi KD, Pardo RB, Silva AV, Cabral KG, et al. Participación de levaduras, algas y hongos en mastitis bovina. *Vet e Zoot*. 1998;10:89–98.
20. Moretti A, Pasquali P, Mencaroni G, Boncio L, Piergili F. Relationship between cell counts in bovine milk and the presence of mastitis pathogens (yeasts and bacteria). *Zentralbl Veterinarmed B*. 1998;45:129–32.
21. Ognean L, Pusta D, Oana L. Signals regarding the isolation of chlorophyll-free algae in the milk of some healthy cows and some with mastitis. *JCEA*. 2001;2:1–2.
22. Panizo MM, Reviákina V, Flores Y, Montes W, González G. Actividad de fosfolipasas y proteasas en aislados clínicos de *Candida* spp. *Rev Soc Ven Microbiol*. 2005;5:64–71.
23. Roostita R, Fleet GH. Growth of yeasts in milk and associated changes to milk composition. *Int J Food Microbiol*. 1996;31:205–19.
24. SAS Institute, Inc. JMP 7.0. Cary, North Carolina. 2007.
25. Scott EM, Gorman SP, McGrath SJ. An assessment of the fungicidal activity of antimicrobial agents used for hard-surface and skin disinfection. *J Clin Hosp Pharm*. 1986;11:199–205.
26. Silva V, Díaz C, Febré N. Red de diagnóstico en Micología Médica. Vigilancia de la resistencia de levaduras a antifúngicos. *Rev Chil Infect*. 2002;19Supl2:S149–156.
27. Spanemberg A, Wüender A, Brayer PDI, Argenta J, Cavallini SEM, Valente P, et al. Diversity of yeasts from bovine mastitis in Southern Brazil. *Rev Iberoam Micol*. 2008;25:154–6.
28. Waltimo TM, Orstavik D, Sirén EK, Haapasalo MP. In vitro susceptibility of *Candida albicans* to four disinfectants and their combinations. *Int Endod J*. 1999;32:421–9.
29. Williamson JH, Di Menna ME. Fungi isolated from bovine udders, and their possible sources. *N Z Vet J*. 2007;55:188–90.