



Revista Iberoamericana de Micología

www.elsevier.es/reviberoammicol



Special Article

Recommendations for the diagnosis of candidemia in Latin America

Arnaldo Lopes Colombo^{a,m,*}, Jorge Alberto Cortes^{b,m}, Jeannete Zurita^{c,m}, Manuel Guzman-Blanco^{d,m}, Tito Alvarado Matute^{e,m}, Flavio de Queiroz Telles^{f,m}, María E. Santolaya^{g,m}, Iris Nora Tiraboschi^{h,m}, Juan Echevarría^{i,m}, Jose Sifuentes^{j,m}, Luis Thompson-Moya^{k,m}, Marcio Nucci^{l,m}

^a Federal University of São Paulo, São Paulo, Brazil

^b Universidad Nacional de Colombia, Bogotá, Colombia

^c Hospital Vozandes, Facultad de Medicina, Pontificia Universidad Católica del Ecuador, Quito, Ecuador

^d Hospital Privado Centro Médico de Caracas, Caracas, Venezuela

^e Hospital Escuela, Tegucigalpa, Honduras

^f Hospital de Clínicas, Universidade Federal do Paraná, Paraná, Brazil

^g Hospital Luis Calvo Mackenna, Universidad de Chile, Santiago, Chile

^h Hospital de Clínicas José de San Martín, University of Buenos Aires, Buenos Aires, Argentina

ⁱ Universidad Peruana Cayetano Heredia, Lima, Perú

^j National Institute of Medical Sciences and Nutrition, Tlalpan, Mexico

^k Clínica Alemana, Universidad del Desarrollo, Santiago, Chile

^l Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

^m Latin America Invasive Mycosis Network

ARTICLE INFO

Article history:

Received 26 March 2013

Accepted 16 May 2013

Available online 10 June 2013

Keywords:

Recommendations

Candidemia

Diagnosis

Latin America

ABSTRACT

Candidemia is one of the most frequent opportunistic mycoses worldwide. Limited epidemiological studies in Latin America indicate that incidence rates are higher in this region than in the Northern Hemisphere. Diagnosis is often made late in the infection, affecting the initiation of antifungal therapy. A more scientific approach, based on specific parameters, for diagnosis and management of candidemia in Latin America is warranted.

'Recommendations for the diagnosis and management of candidemia' are a series of manuscripts that have been developed by members of the Latin America Invasive Mycosis Network. They aim to provide a set of best-evidence recommendations for the diagnosis and management of candidemia.

This publication, 'Recommendations for the diagnosis of candidemia in Latin America', was written to provide guidance to healthcare professionals on the diagnosis of candidemia, as well as on the usefulness and application of susceptibility testing in patients who have a confirmed diagnosis of candidemia.

Computerized searches of existing literature were performed by PubMed. The data were extensively reviewed and analyzed by members of the group. The group also met on two occasions to pose questions, discuss conflicting views, and deliberate on a series of management recommendations.

'Recommendations for the diagnosis of candidemia in Latin America' includes diagnostic methods used to detect candidemia, *Candida* species identification, and susceptibility testing. The availability of methods, their costs and treatment settings are considered.

This manuscript is the first of this series that deals with diagnosis and treatment of invasive candidiasis. Other publications in this series include: 'Recommendations for the management of candidemia in adults in Latin America', 'Recommendations for the management of candidemia in children in Latin America', and 'Recommendations for the management of candidemia in neonates in Latin America'.

This article is also published in Spanish in this issue. It can be found in <http://dx.doi.org/10.1016/j.riam.2013.05.009>

© 2013 Revista Iberoamericana de Micología. Published by Elsevier España, S.L. All rights reserved.

* Corresponding author.

E-mail address: colomboal@terra.com.br (A.L. Colombo).

Recomendaciones para el diagnóstico de la candidemia en América Latina

R E S U M E N

Palabras clave:
Recomendaciones
Candidemia
Diagnóstico
América Latina

La candidemia es una de las micosis oportunistas más frecuentes en todo el mundo. El escaso número de estudios epidemiológicos llevados a cabo en América Latina indica que las tasas de incidencia en esta región son mayores que las descritas en el hemisferio norte. A menudo el diagnóstico de la infección se establece tardíamente, lo que afecta al inicio del tratamiento antimicótico. Por esta razón, para el diagnóstico y el manejo de la candidemia está justificada una estrategia más científica, basada en parámetros específicos.

Recomendaciones para el diagnóstico y manejo de la candidemia constituye una serie de artículos preparados por miembros del grupo Latin America Invasive Mycosis Network. Su objetivo es proporcionar las mejores evidencias disponibles para el diagnóstico y el manejo de la candidemia.

El presente artículo, *Recomendaciones para el diagnóstico de la candidemia en América Latina*, ha sido redactado con el objetivo de brindar asesoramiento a los profesionales de la salud en lo referente al diagnóstico de la candidemia en pacientes que la padecen o están en riesgo de padecerla.

Mediante la base de datos PubMed se emprendió una búsqueda informatizada de los estudios publicados. Los miembros del grupo revisaron y analizaron exhaustivamente los datos. El grupo también se reunió en 2 ocasiones para proponer preguntas, abordar los puntos de vista conflictivos y deliberar sobre las recomendaciones terapéuticas.

Recomendaciones para el diagnóstico de la candidemia en América Latina incluye diversas recomendaciones sobre aspectos relacionados con los métodos diagnósticos para la detección de la candidemia, la identificación de las especies de *Candida* y las pruebas de sensibilidad antifúngica. Se expone también la disponibilidad de los métodos, sus costes y el marco en el que se aplican los tratamientos.

Este manuscrito es el primero de los artículos de esta serie dedicada al diagnóstico y tratamiento de las candidiasis invasoras. Otras publicaciones de esta serie son *Recomendaciones para el diagnóstico de la candidemia en adultos en América Latina*, *Recomendaciones para el manejo de la candidemia en niños en América Latina*, y *Recomendaciones para el manejo de la candidemia en neonatos en América Latina*.

Este artículo está publicado en español en este mismo número. Puede encontrarlo en <http://dx.doi.org/10.1016/j.riam.2013.05.009>

© 2013 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Challenges of invasive candidiasis diagnosis

The diagnosis of invasive candidiasis is often made late in the course of infection, causing a delay in the initiation of antifungal therapy. Late diagnosis can be the result of non-specific clinical signs and symptoms,¹⁸ the variable accuracy of the available diagnostic tests,²⁷ delayed time to grow *Candida*,¹⁹ blood cultures that are not positive until late-stage infection,^{18,25} inadequate sample size for blood culture,²⁵ and false-negative results due to the use of antifungal agents in prophylaxis.²⁵

Overcoming the challenges of diagnosis can improve time-to-treatment of *Candida* infection. Here, we provide recommendations for the diagnosis of invasive candidiasis, including methods for detecting infection, *Candida* species identification, and susceptibility testing.

Methods for detecting hematogenous *Candida* infection

Diagnosis can be achieved through various procedures, including conventional methods such as blood culture and serological assays for surrogate markers (1-3- β -D-glucan [BDG] and mannan antigen/antibody). Newer diagnostic methods include polymerase chain reaction (PCR) from blood or tissue biopsy, and enzyme-linked immunosorbent assay (ELISA) to detect an antigen produced by *Candida*.

Blood culture

Blood culture is by far the best method available for the diagnosis of candidemia; however, it is a lengthy procedure. This can affect timing of the initiation of antifungal therapy and contribute to increased morbidity and mortality.⁴⁹ Time to detection can be influenced by *Candida* spp. In one retrospective study, mean times

to positive culture for *Candida albicans* and *Candida glabrata* were 35.3 ± 18.1 h and 80.0 ± 22.4 h, respectively.¹⁹ The type of medium used for culture can also influence time to detection. A comparison of a selective fungal medium and a standard aerobic medium for the diagnosis of candidemia demonstrated a significant time saving when the fungal medium was used, with a mean time saving of 8.8 h for *C. albicans* and 43.7 h for *C. glabrata*.⁴⁶

As mentioned previously, blood culture is the gold standard for the diagnosis of candidemia; however, its sensitivity is variable.⁵¹ In order to optimize this method, factors which are known to influence sensitivity (e.g. blood volume, number of cultures, and time to detection) should be carefully considered, as should the size of the inoculum, as well as type of culture bottles and media used. In one study with 15 *Candida* spp., higher inoculum sizes (10, 100, 1000 yeast cells per bottle) resulted in 70%, 73% and 79% detection of growth, respectively.²⁷ A comparable rate of detection of *Candida* in aerobic and mycology bottles has been observed,^{8,27,45} which is higher than that in anaerobic bottles.^{8,27} The Working Group recommends the use of aerobic bottles for the diagnosis of candidemia by blood culture, as this is the standard procedure in every hospital.

Blood culture systems

A number of blood culture systems are available, which vary in sensitivity. These include conventional (manual), automated, and lysis-centrifugation methods. The use of biphasic media appears to be slightly better than or equivalent to conventional broth in yeast recovery.³³ The sensitivity of automated blood culture systems, such as BacT/ALERT, BACTEC, and Isolator, is influenced by the type of culture bottle and media used.^{28,46} The lysis-centrifugation blood culture technique is the most sensitive method available. It has better diagnostic yields compared with conventional blood culture systems and reduces the time to result.³⁸ However, lysis centrifugation is labor intensive, expensive, and has a high rate of

contamination,¹⁸ and for this reason the Working Group considers automated blood culture systems the best culture option for the diagnosis of fungemia due to *Candida* spp.

Serology

A number of serological assays for the detection of candidemia are currently under development. Due to limited information on their diagnostic value, the Working Group cannot make a recommendation on their usage. These methods include the BDG assay,⁶⁵ and mannan antigen and antibody assays.^{1,47} A recent systematic review of serological methods for the diagnosis of candidemia in critically ill patients showed a higher diagnostic sensitivity for the combined use of antigen and antibody.¹¹

BDG is a structural component of *Candida* cell walls that shows promise as a biomarker for candidemia. Studies using the serum BDG assay for the diagnosis of candidemia reported assay sensitivities between 57% and 100% and specificities between 44% and 92%, much greater than blood culture methods.⁶⁵ However, the BDG test is prone to false-positive results from β -glucan contamination by certain antibiotics and materials.²⁵ Sources of β -glucan include dialysis membranes and filters made from cellulose, specific immunoglobulin products, cotton gauze, and sponges used in surgeries, as well as some drugs (including lentinan, crestin, scleroglucan, and schizophyllan).^{35,37,50,63} BDG has been extensively used in research and has been approved by regulatory agencies in both the USA and Europe for the diagnosis of *Candida* infection. However, owing to lack of availability and high costs, very few centers in Latin America are likely to be able to use this assay. Given its high negative predictive value, repeated negative BDG tests may help to rule out candidemia. Nevertheless, at present it is not clear how this test should be utilized at bedside, in terms of helping the clinician to decide whether or not an antifungal drug should be prescribed to a patient at risk of developing candidemia.

Screening of *Candida* deep-seated infections

Imaging

Echocardiography has demonstrated to be an effective tool for the diagnosis of *Candida*-induced endocarditis,¹⁵ and magnetic resonance imaging (MRI) and computed tomography (CT) have shown to be effective non-invasive tools for the identification of hepatosplenic fungal disease.^{26,43} Although MRI and CT are sensitive in detecting the fungal microabscesses associated with chronic disseminated candidiasis (CDC), the requirement for repeat images during the course of an infection limits their usage. Computer-assisted ultrasonography has also been shown to successfully detect microabscesses in the liver and/or spleen and can be repeated as often as required, making it a useful method for the detection of CDC and for the follow-up of patients with CDC.³⁶

Biopsy

The diagnosis of infections due to *Candida* spp. from particular specimens can be complicated. These organisms commonly colonize human skin and mucous membranes and, therefore, culture from specimens (sputum, tracheal secretions, and urine) from these sites is not necessarily a proof of invasion.³³ Deep-organ *Candida* infection may require tissue biopsy to establish diagnosis, as in the case of *Candida*-induced pneumonia; however, tissue samples may contain few organisms, resulting in negative culture.⁶⁴ CDC is diagnosed through identification of fungal structures under the microscope, or fungal growth from biopsy materials, as blood cultures are positive in only <20% of CDC patients.⁴³ However, *Candida* may not always be detected upon biopsy, and a biopsy may not be feasible in some patients owing to the risk of complications.⁴³ In at-risk patients who develop skin lesions, biopsy is utilized for the

confirmation of diagnosis of *Candida* pneumonia via the identification of fungal elements.

New perspectives

Recently an ELISA was developed for the detection of a 65 kDa antigen produced by *C. albicans*, *Candida tropicalis* and *Candida parapsilosis*. In one study, this novel diagnostic test detected the 65 kDa antigen in 80% of patients with candidemia.⁴ The advent of PCR for the detection of fungal nucleic acids has provided a faster and more accurate alternative to blood culture for the detection of *Candida* in blood samples.⁴¹ In addition, PCR has been used in the detection of *Candida* in tissue-biopsy specimens from patients with CDC.^{22,39}

A recent meta-analysis covering 54 studies found PCR-positivity rates among patients with proven or possible candidemia to be 85% (78–91%), compared with just 38% (29–46%) for blood cultures.³ An improved sensitivity for PCR to detect candidemia has been achieved with the use of serum and plasma samples as compared with whole blood. In one study, *Candida* DNA was detected in 71% of serum samples and 75% of plasma samples, compared with 54% of whole blood samples.⁴¹ A limitation of this assay was the inability to detect fungal DNA in 25% of whole blood specimens drawn at the same time as positive blood culture samples.⁴¹

Reports of false-positive, as well as false-negative, results when PCR has been used for the detection of *Candida* in seeded clinical specimens such as blood, urine, and peritoneal fluid, have been described.⁷ In another study, a false-positive result was reported in two patients who had been treated with azole antifungal agents, which may indicate that PCR was able to detect non-viable organisms that could not be recovered by culture.²¹ Compared with antigen and antibody methods, PCR has a higher sensitivity and specificity for diagnosis.¹¹ Despite this, no international standardization exists for this method, and there are no commercially available PCR kits for diagnosing *Candida*. Consequently, PCR is not considered a reliable tool for use in clinical laboratories to help clinicians to identify patients with candidemia early.

Recommendations summary for detecting hematogenous *Candida* infection

1. Blood culture is the gold-standard method for the diagnosis of invasive candidiasis.
2. Aerobic bottles and automated blood culture systems are recommended to achieve optimum sensitivity.
3. Following diagnosis, *Candida* isolates should be identified at species level.

Additional diagnostic method recommendations

1. Imaging (e.g. echocardiography, MRI and CT) may play a part in the diagnosis of endocarditis and CDC.
2. Histopathology and/or direct microscopy can be considered for the diagnosis of deep-seated *Candida* infections, as with CDC or pneumonia. In case of CDC, a negative microscopic examination does not indicate the absence of disease.

Methods for *Candida* identification

Following the detection of a pathogen, early species identification is important to initiate appropriate antifungal therapy or to modify existing antifungal therapy. Species identification can be achieved by one or more of the following methods: conventional

direct examination and staining, screening (germ tubes or chromogenic media), the MicroScan Yeast Identification Panel, manual commercial systems (e.g. API 20C, and API 32C), automated commercial systems (e.g. VITEK 2) and molecular identification (PCR, matrix-assisted laser desorption ionization time-of-flight [MALDI-TOF] or peptide nucleic acid fluorescent *in situ* hybridization [PNA FISH]).

Conventional direct examination and staining

In general, direct microscopy, using a number of different stains, is a quick and relatively cheap method to initially identify a pathogen as yeast or bacterium.¹⁷ In patients with deep-seated *Candida* infections, microscopic identification of fungal elements in biopsy material or fungal growth from such biopsies has been considered the gold standard for the diagnosis of CDC³² or *Candida*-induced pneumonia. It may also be useful for the diagnosis of hematogenous candidiasis in patients who develop skin lesions secondary to *Candida* dissemination.

The microbiological diagnosis of fungal infections via direct microscopy of tissue sections is generally performed using wet preparations (potassium hydroxide, calcofluor white) and Gram, Wright or Giemsa stains.⁴⁸ Specimens should be examined for the presence of small, round-to-oval, thin-walled budding yeast cells in clusters and branching pseudohyphae⁶²; however, other genera of fungi may also exhibit this morphology. For this reason, further examination may be necessary to exclude other genera. Following Gram staining, the capsule of *Cryptococcus* can be visualized as a faint red halo, and the presence of arthroconidia may indicate *Trichosporon*.⁵³ It is important to note that a negative microscopic examination does not necessarily exclude infection.^{23,48}

Peptide nucleic acid-fluorescent *in situ* hybridization

The Yeast Traffic Light PNA FISH is a novel molecular method which may help the fast identification of the most clinically relevant species of *Candida* as soon as the culture is positive⁶⁰. Following Gram stain to check for the presence of *Candida*, slides are prepared to generate fluorescence signals characteristic of individual *Candida* species. Despite being a useful test, the need of fluorescent microscopy and the cost of the reagents limit its use in clinical laboratories in Latin America.

Phenotypic culture examination

Fast screening

A germ tube test (GTT) is a prompt and easy method for the screening of *C. albicans*.⁵⁷

Chromogenic media

In Latin America, the Working Group recommends the use of CHROMagar media for species identification as it is more accurate, although less expensive chromogenic media are also available (e.g. Oxoid media). In one study, CHROMagar *Candida* medium allowed the isolates from 93% of clinical specimens to be identified at the species level, with up to 92% accuracy, within 48 h. The cost of this technique was comparable with conventional methods, making it an affordable testing method for laboratories from countries with limited resources.³⁰ A restriction of the CHROMagar method is the lack of a wide range of colors, which confines identification to a limited number of species. These include *C. albicans* (indistinguishable by this method from *Candida dubliniensis*), *C. tropicalis* and *Candida krusei*. The isolates of *C. glabrata*, *C. parapsilosis*, *Candida guilliermondii*, *Candida kefyr* and *Candida lusitanae* have various tones on CHROMagar from off-white to pink, which are difficult to define because they are similar pastel tones and not typical for any

Table 1

Colony colors of yeast isolates incubated for 2 days on CHROMagar *Candida* medium at 37 °C.

Species	Color and morphology ⁵²
<i>Candida albicans</i>	Green
<i>Candida tropicalis</i>	Dark blue to blue-gray (with dark halo in agar)
<i>Candida krusei</i>	Pale pink to purple (rough with spreading pale edges)

species.⁶⁶ This limitation highlights the requirement for alternative methods for extended *Candida* spp. identification.⁵ When culturing biological materials or yeast colonies in chromogenic media, it is important to check for mixed cultures and infection by more than one fungal pathogen.

CHROMagar is prepared according to the manufacturer's instructions and poured into sterile Petri dishes or tubes and allowed to gel and dry. Prepared media plates can be kept for 24 h at ambient temperature or for up to one month if refrigerated and protected from light and dehydration.⁹ If refrigerated, plates should be warmed to room temperature prior to inoculation via streaking the sample onto the plate and incubation at 30–37 °C for 48 h.⁹ Identification of the species is based on colony color (Table 1).^{29,52}

Phenotypic tests for *Candida* identification

It is essential that all yeasts causing invasive fungal infections in humans are identified to species level. Species identification can be determined by the following methods.

"In-house" classical method

It is possible to verify the biochemical profile of yeast colonies by using "in-house" sugar assimilation and fermentation tests. Several laboratories in Latin America prepare their own biochemical test panels, which include 7–12 different sugars for assimilation and fermentation. Results from such assays can then be compared against existing databases (in textbooks) to verify species.

Manual commercial systems

The biochemical profile of yeasts under identification may be checked by commercially available tests. Following inoculation with yeast, a number of commercially available systems provide different parameters associated with the growth of colonies and a particular substrate. Some detect change in turbidity compared with a control well (e.g. API 20C AUX, and API 32C), whereas others use color generation in a series of wells (e.g. API *Candida*, Auxa-color, and Uni-Yeast-Tek).¹⁸ These tests provide accurate results for the commonest *Candida* species; however, they are not as reliable for the identification of uncommon species of *Candida*. Accuracy of results is highly dependent on the number of tests available in each commercial system as well as their database (i.e. limited or broad).

Automated commercial systems

VITEK 2 is a fully automated instrument which uses a fluorescent-based technology for identification of yeasts and yeast-like organisms within 15 h.²⁴ The fluorometric card has been replaced by a colorimetric card to broaden the database.² The MicroScan Yeast Identification Panel is a 4-hour microdilution system for the identification of yeast isolates. It has a 94% identification success rate for 22 species of *Candida*.^{40,59}

Molecular identification

As mentioned previously, PCR can be used for the detection of fungal nucleic acids. It can also serve to identify *Candida* in blood

samples, with species identification time reduced from a mean of 3.5 days by routine phenotypic methods to 7 h.⁵⁸ Furthermore, PCR is able to detect the presence of more than one species of *Candida* in the same patient, whereas blood cultures may yield only one species. The ability to detect and identify more than one species of *Candida* may affect therapy and outcome, particularly when one of the species is known to be resistant to azole compounds.¹⁴

New perspectives (proteomics)

Mass spectrometry (MALDI-TOF system) has been used for the identification of *Candida* species in less than 30 min with a 91% success rate.²⁰ MALDI-TOF has been shown to successfully identify 100% of *Candida* isolates, compared with 92% identification using existing biochemical methods (VITEK 2 and API C AUX).⁴²

The availability of specific diagnostic methods for the detection and identification of candidemia is dependent on the clinical setting. For small hospitals that do not provide care for transplant patients or treat many hematological and immunocompromised patients, the minimal requirements suggested by the Working Group for yeast identification are the capability to evaluate the micromorphology of the colonies complemented by a screening of the main relevant species of *Candida* by means of CHROMagar *Candida* medium and/or any in-house or a commercial kit to conduct biochemical tests. For tertiary hospitals, in addition to micromorphology and CHROMagar, species identification should be determined by manual commercial systems (API 20C, API 32C), automated commercial systems (VITEK 2) and/or molecular methods. Molecular methods should be considered in the identification of emerging pathogens where conventional tools usually generate inconsistent identification and when facing outbreaks of fungal infections.

Recommendations summary for *Candida* species identification

1. The Working Group recommends, as the minimum requirement, colony micromorphology observation complemented by macromorphology using CHROMagar *Candida* medium.
2. For secondary hospitals, species identification may be determined using one or more of the following methods:
 - (a) Colony micromorphology.
 - (b) Colony macromorphology (CHROMagar *Candida* medium).
 - (c) Biochemical tests.
 - I. In-house conventional methods.
 - II. Manual commercial systems with a limited database (e.g. Auxacolor and Uni-Yeast-Tek).
3. For tertiary care hospitals that provide care for transplant patients or treat many hematological and immunocompromised patients, the Working Group recommends, as minimum requirement:
 - (a) Micromorphology observation complemented by biochemical tests (API 20C, API 32C, VITEK 2 or MicroScan Yeast Identification Panel).
 - (b) Molecular methods in specific situations.
4. Molecular methods (PCR and MALDI-TOF) should be considered in the identification of emerging pathogens and when investigating outbreaks.

Methods for susceptibility testing

It is important to note that yeast identification should always take priority over susceptibility testing. Knowledge of the

susceptibilities of local clinical isolates to individual antifungal agents can influence treatment choice. The Working Group encourages laboratories to collaborate with each other when performing susceptibility testing as this will ultimately reduce costs and increase accuracy.

If susceptibility testing is performed, it should be noted that there are still some controversies in the definition of clinical breakpoints for echinocandins and amphotericin B. Screening tests for fluconazole resistance provide the most reliable results compared with other antifungal agents. A number of methods exist for *Candida* susceptibility testing. These include reference broth-based methods (e.g. standard M27-A of the Clinical and Laboratory Standards Institute [CLSI] and standard EDef 7.1 of the European Committee on Antimicrobial Susceptibility Testing [EUCAST]), disk-based reference methods (e.g. CLSI standard M44-A), commercially available systems (including Fungitest, Sensititre YeastOne, Etest and Neo-sensitabs) and other methods (including flow cytometry and VITEK 2).

Candida susceptibility testing reference methods

The reference broth-based method (M27-A) became a National Committee for Clinical Laboratory Standards Institute approved method for antifungal-susceptibility testing in 1997.⁵⁴ Since the Institute's change of name in 2005, it is now known as the CLSI method.¹⁰ The EUCAST⁶¹ reference broth-based method is known as EDef 7.1 Both CLSI M27-A and EUCAST EDef 7.1 specify the size of the inoculum, test media, incubation temperature and duration, and end-point readings for the test drugs (Table 2).

Both methods produce similar minimum inhibitory concentrations (MICs [although EDef 7.1 results usually produce slightly lower MICs]),^{16,55} which indicates that methodology does not pose any obstacle to obtaining uniform standards in antifungal-susceptibility testing of yeasts. However, these methods are expensive and can be very laborious. Therefore, their actual usage in hospital laboratories is limited; many hospital laboratories prefer to use commercially available products, which claim to be faster and more accurate than the M27-A method.

The disk-based susceptibility testing method (CLSI M44-A) provides a zone of inhibition, a measurement that can be correlated with the MIC value. A good correlation between disk-based susceptibility testing and the M27-A reference method has been shown for fluconazole.^{44,54} This method is not widely used in Latin America, as fluconazole and voriconazole disks have a limited distribution.

Commercially available susceptibility tests

It is important to note that some of the commercially available products presented in this section may not currently be available in Latin America. Commercial broth-based MIC systems on the market include Candifast, Integral Systems Yeasts, and Fungitest. However, such methods show a limited correlation with the M27-A reference method.⁵⁴

The Sensititre YeastOne generates a colorimetric response. This method has $\geq 85\%$ rate of concordance with the M27-A reference method,⁵⁴ and a 92% rate of concordance with the EDef 7.1 reference method.¹³ A more recent paper presented high concordance between Sensititre Yeast One and EUCAST, with an essential agreement rate of $\geq 95.5\%$ (dependent on antifungal agent tested).¹²

E-test is an antimicrobial-susceptibility test which has been adapted for using with fungal agents. This simple method involves surface inoculation of an agar plate followed by the application of a plastic strip, which contains a concentration gradient of the antifungal agent. Following incubation of a susceptible yeast, a zone of

Table 2

CLSI and EUCAST conditions for antifungal susceptibility testing.

Characteristic	CLSI M27-A3 ¹⁰	EUCAST Def. 7. ⁶¹
Suitability	Yeasts	Fermentative yeasts
Inoculum	0.5×10^3 to 2.5×10^3 CFU/mL	0.5×10^5 to 2.5×10^5 CFU/mL
Inoculum standardization	Spectrophotometric, with reference to a 0.5 McFarland BaSO ₄ turbidity standard	Spectrophotometric, with reference to a 0.5 McFarland BaSO ₄ turbidity standard
Test medium	RPMI 1640, buffered to pH 7.0 with 0.165 M MOPS	RPMI 1640, buffered to pH 7.0 with 0.165 M MOPS and glucose 2% w/v
Format	Macro- or micro-dilution	Micro-dilution
Temperature	35 °C	35 °C
Duration of incubation	24–48 h	24 h (<i>Candida</i> spp.)
End point	Optically clear well for amphotericin B, ~50% reduction in visible growth (microdilution testing with flucytosine, echinocandins, and azoles ^a)	Optically clear well for amphotericin B, ~90% reduction in visible growth; for anidulafungin, caspofungin, flucytosine, and azoles ^a , the MIC end point is the lowest drug concentration at which the growth of the isolates is reduced by 50% or more compared with that of the control (MIC-2 or ≥50%)
Quality control isolates and drugs	Two <i>Candida</i> isolates for checking their susceptibility to amphotericin B, flucytosine, azoles, ^a caspofungin, and anidulafungin	<i>Candida parapsilosis</i> ATCC 22019 and <i>Candida krusei</i> ATCC 6258

Adapted from Ref.⁵⁵

CFU = colony-forming unit; CLSI = Clinical and Laboratory Standards Institute; EUCAST = European Committee for Antimicrobial Susceptibility Testing; RPMI = Roswell Park Memorial Institute; MIC = minimum inhibitory concentration; MOPS = 3-(N-morpholino)propanesulfonic acid; w/v = weight/volume.

^a Fluconazole, voriconazole, ketoconazole, itraconazole, ravuconazole and posaconazole.

inhibition can be seen and the MIC is read at the point at which the zone intersects the strip.³¹ The correlation between the E-test method and the reference methods M27-A and EDef 7.1 has been described as acceptable for most species of *Candida* and azole-based antifungal agents.^{13,54}

Neo-Sensitabs is a commercial disk-based system. It has been reported that the results from this system do not correlate well with the M27-A and EDef 7.1 reference methods.^{13,54}

Other susceptibility testing methods

The use of flow cytometry as a potential tool for antifungal-susceptibility testing has long been recognized and it is reported to provide rapid results which compare well with standard reference methods.^{34,56} The VITEK 2 system, which is used for the identification of yeasts, can also be used for susceptibility testing. A comparative evaluation study found an excellent categorical agreement between VITEK 2 and the M27-A reference methods for determining the MIC values of antifungal agents (92–98.2%); however, discrepancies were observed within species.⁶

Recommendations summary for *Candida* susceptibility testing

1. Yeast identification should always take priority over susceptibility testing.
2. Laboratories should collaborate to reduce costs and increase accuracy of tests.
3. Fluconazole-susceptibility testing is recommended for tertiary care hospitals.

Conflict of interests

A.L. Colombo has received research grants from Pfizer, MSD, United Medical and Luminex, medical education grants from Pfizer, MSD, United Medical and Astellas. Moreover, he has also been a consultant for MSD, Pfizer and Gilead. J.A. Cortes has received research grants and support to attend educational meetings from

Pfizer and MSD. M. Nucci has received research grants from Pfizer and MSD, and has acted as a consultant and speaker for Pfizer, MSD, Astellas and Gilead. F. de Queiroz Telles has participated in Continuing Education activities in laboratories for Astellas, MSD, Pfizer and United Medical, and in research activities in laboratories for Astellas, MSD and Pfizer. I.N. Tiraboschi has been a speaker for Pfizer and Gilead. J. Zurita has been advisory board member and consultant for Pfizer, and has received research grants from Wyeth and MSD for participating in the SMART study.

Acknowledgements

Editorial support in the form of assistance with the first draft, collating author comments and editorial suggestions to draft versions of this manuscript was provided by Jacqueline Adam, PhD, of Choice Healthcare Solutions and funded by Pfizer. Responsibility for opinions, conclusions and recommendations lies with the authors.

References

1. Alam FF, Mustafa AS, Khan ZU. Comparative evaluation of (1, 3)-beta-D-glucan, mannan and anti-mannan antibodies, and *Candida* species-specific snPCR in patients with candidemia. BMC Infect Dis. 2007;7:103.
2. Aubertine CL, Rivera M, Rohan SM, Larone DH. Comparative study of the new colorimetric VITEK 2 yeast identification card versus the older fluorometric card and of CHROMagar Candida as a source medium with the new card. J Clin Microbiol. 2006;44:227–8.
3. Avni T, Leibovici L, Paul M. PCR diagnosis of invasive candidiasis: systematic review and meta-analysis. J Clin Microbiol. 2011;49:665–70.
4. Berzaghi R, Colombo AL, Machado AM, de Camargo ZP. New approach for diagnosis of candidemia based on detection of a 65-kilodalton antigen. Clin Vac Immunol. 2009;16:1538–45.
5. Bishop JA, Chase N, Lee R, Kurtzman CP, Merz WG. Production of white colonies on CHROMagar Candida medium by members of the *Candida glabrata* clade and other species with overlapping phenotypic traits. J Clin Microbiol. 2008;46:3498–500.
6. Borghi E, Iatta R, Sciota R, Biassoni C, Cuna T, Montagna MT, et al. Comparative evaluation of the Vitek 2 yeast susceptibility test and CLSI broth microdilution reference method for testing antifungal susceptibility of invasive fungal isolates in Italy: the GISIA3 study. J Clin Microbiol. 2010;48:3153–7.
7. Burgener-Kairuz P, Zuber JP, Jaunin P, Buchman TG, Bille J, Rossier M. Rapid detection and identification of *Candida albicans* and *Torulopsis (Candida) glabrata* in clinical specimens by species-specific nested PCR amplification of

- a cytochrome P-450 lanosterol- α -demethylase (L1A1) gene fragment. *J Clin Microbiol.* 1994;32:1902–7.
8. Chiarini A, Palmeri A, Amato T, Immordino R, Distefano S, Giammanco A. Detection of bacterial and yeast species with the Bactec 9120 automated system with routine use of aerobic, anaerobic, and fungal media. *J Clin Microbiol.* 2008;46:4029–33.
9. CHROMagar. CHROMagar Candida chromogenic medium for the isolation and differentiation of major *Candida* species; 2009.
10. CLSI. Reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard – third edition, 2008.
11. Cortes JA, Concha MA, Cediell TL, Castillo JS. Diagnostic methods in candidemia: a systematic review of literature with meta-analysis. *Rev Chilena Infectol.* 2011;28:423–8.
12. Cuenca-Estrella M, Gomez-Lopez A, Alastruey-Izquierdo A, Bernal-Martinez L, Cuesta I, Buitrago MJ, et al. Comparison of the Vitek 2 antifungal susceptibility system with the clinical and laboratory standards institute (CLSI) and European Committee on Antimicrobial Susceptibility Testing (EUCAST) Broth Microdilution Reference Methods and with the Sensititre YeastOne and Etest techniques for in vitro detection of antifungal resistance in yeast isolates. *J Clin Microbiol.* 2010;48:1782–6.
13. Cuenca-Estrella M, Gomez-Lopez A, Mellado E, Rodriguez-Tudela JL. Correlation between the procedure for antifungal susceptibility testing for *Candida* spp. of the European Committee on Antibiotic Susceptibility Testing (EUCAST) and four commercial techniques. *Clin Microbiol Infect.* 2005;11:486–92.
14. Del Negro GM, Delgado AF, Manuli ER, Yamamoto L, Okay TS. Dual candidemia detected by nested polymerase chain reaction in two critically ill children. *Med Mycol.* 2010;48:1116–20.
15. Donal E, Abgueuen P, Coisne D, Gouello JP, McFadden EP, Allal J, et al. Echocardiographic features of *Candida* species endocarditis: 12 cases and a review of published reports. *Heart.* 2001;86:179–82.
16. Duarte C, Pulido N, Rivas P, Sánchez R, Cortés JA, Cuervo S, et al. CLSI microdilution M27-A2 and EUCAST method comparison for *Candida* spp. in patients with cancer. *Rev Infect.* 2010;14(S2):S107–15.
17. Dunn DL. Diagnosis and treatment of infection. In: Norton JA, Barie PS, Bollinger R, Chang AE, Lowry SF, Mulvihill SJ, et al., editors. *Surgery: basic science and clinical evidence.* Springer; 2008. p. 209–35.
18. Ellepola AN, Morrison CJ. Laboratory diagnosis of invasive candidiasis. *J Microbiol.* 2005;43(Spec. No.):65–84.
19. Fernandez J, Erstad BL, Petty W, Nix DE. Time to positive culture and identification for *Candida* blood stream infections. *Diagn Microbiol Infect Dis.* 2009;64:402–7.
20. Ferroni A, Suarez S, Beretti JL, Dauphin B, Bille E, Meyer J, et al. Real-time identification of bacteria and *Candida* species in positive blood culture broths by matrix-assisted laser desorption ionization-time of flight mass spectrometry. *J Clin Microbiol.* 2010;48:1542–8.
21. Flahaut M, Sanglard D, Monod M, Bille J, Rossier M. Rapid detection of *Candida albicans* in clinical samples by DNA amplification of common regions from *C. albicans*-secreted aspartic proteinase genes. *J Clin Microbiol.* 1998;36:395–401.
22. Fleischacker M, Schulz S, Johrens K, von Lilienfeld-Toal M, Held T, Fietze E, et al. Diagnosis of chronic disseminated candidosis from liver biopsies by a novel PCR in patients with haematological malignancies. *Clin Microbiol Infect.* 2012;18:1010–6.
23. Gilad J, Giladi M, Schwartz D. *Candida albicans* masquerading as Gram-negative bacilli in the clinical laboratory. *Scand J Infect Dis.* 2007;39:907–10.
24. Graf B, Adam T, Zill E, Gobel UB. Evaluation of the VITEK 2 system for rapid identification of yeasts and yeast-like organisms. *J Clin Microbiol.* 2000;38:1782–5.
25. Guery BP, Arendrup MC, Auzinger G, Azoulay E, Borges Sa M, Johnson EM, et al. Management of invasive candidiasis and candidemia in adult non-neutropenic intensive care unit patients: Part I. Epidemiology and diagnosis. *Intensive Care Med.* 2009;35:55–62.
26. Halkic N, Ksontini R. Images in clinical medicine. Hepatosplenic candidiasis. *N Engl J Med.* 2007;356:e4.
27. Horvath LL, George BJ, Hospenthal DR. Detection of fifteen species of *Candida* in an automated blood culture system. *J Clin Microbiol.* 2007;45:3062–4.
28. Horvath LL, George BJ, Murray CK, Harrison LS, Hospenthal DR. Direct comparison of the BACTEC 9240 and BacT/ALERT 3D automated blood culture systems for *Candida* growth detection. *J Clin Microbiol.* 2004;42:115–8.
29. Horvath LL, Hospenthal DR, Murray CK, Dooley DP. Direct isolation of *Candida* spp. from blood cultures on the chromogenic medium CHROMagar Candida. *J Clin Microbiol.* 2003;41:2629–32.
30. Iyampillai T, Michael JS, Mathai E, Mathews MS. Use of CHROMagar medium in the differentiation of *Candida* species: is it cost-effective in developing countries? *Ann Trop Med Parasitol.* 2004;98:279–82.
31. Johnson EM. Issues in antifungal susceptibility testing. *J Antimicrob Chemother.* 2008;61 Suppl. 1:i13–8.
32. Johnson TL, Barnett JL, Appelman HD, Nostrant T. *Candida* hepatitis. Histopathologic diagnosis. *Am J Surg Pathol.* 1988;12:716–20.
33. Jones JM. Laboratory diagnosis of invasive candidiasis. *Clin Microbiol Rev.* 1990;3:32–45.
34. Joong YH, Kim HR, Lee MK, Park AJ. Fluconazole susceptibility testing of *Candida* species by flow cytometry. *J Infect.* 2007;54:504–8.
35. Kanda H, Kubo K, Hamasaki K, Kanda Y, Nakao A, Kitamura T, et al. Influence of various hemodialysis membranes on the plasma (1 $\rightarrow$ 3)- β -D-glucan level. *Kidney Int.* 2001;60:319–23.
36. Karthaus M, Huebner G, Elser C, Geissler RG, Heil G, Ganster A. Early detection of chronic disseminated *Candida* infection in leukemia patients with febrile neutropenia: value of computer-assisted serial ultrasound documentation. *Ann Hematol.* 1998;77(1/2):41–5.
37. Kato A, Takita T, Furuhashi M, Takahashi T, Maruyama Y, Hishida A. Elevation of blood (1 $\rightarrow$ 3)- β -D-glucan concentrations in hemodialysis patients. *Nephron.* 2001;89:15–9.
38. Kiehn TE, Wong B, Edwards FF, Armstrong D. Comparative recovery of bacteria and yeasts from lysis-centrifugation and a conventional blood culture system. *J Clin Microbiol.* 1983;18:300–4.
39. Kirby A, Chapman C, Hassan C, Burnie J. The diagnosis of hepatosplenic candidiasis by DNA analysis of tissue biopsy and serum. *J Clin Pathol.* 2004;57:764–5.
40. Land GA, Salkin IF, el-Zaatari M, McGinnis MR, Hashem G. Evaluation of the Baxter-MicroScan 4-hour enzyme-based yeast identification system. *J Clin Microbiol.* 1991;29:718–22.
41. Lau A, Halliday C, Chen SC, Playford EG, Stanley K, Sorrell TC. Comparison of whole blood, serum, and plasma for early detection of candidemia by multiplex-tandem PCR. *J Clin Microbiol.* 2010;48:811–6.
42. Martinez-Lamas L, Perez Del Molino ML, Pardo F, Varela E, Regueiro BJ. Matrix-assisted laser desorption ionization time-of-flight (MALDI-TOF) mass spectrometry vs conventional methods in the identification of *Candida* non-albicans. *Enferm Infect Microbiol Clin.* 2011;29:568–72.
43. Masood A, Sallah S. Chronic disseminated candidiasis in patients with acute leukemia: emphasis on diagnostic definition and treatment. *Leuk Res.* 2005;29:493–501.
44. Matar MJ, Ostrosky-Zeichner L, Paetznick VL, Rodriguez JR, Chen E, Rex JH. Correlation between E-test, disk diffusion, and microdilution methods for antifungal susceptibility testing of fluconazole and voriconazole. *Antimicrob Agents Chemother.* 2003;47:1647–51.
45. McDonald LC, Weinstein MP, Fune J, Mirrett S, Reimer LG, Reller LB. Controlled comparison of BacT/ALERT FAN aerobic medium and BATEC fungal blood culture medium for detection of fungemia. *J Clin Microbiol.* 2001;39:622–4.
46. Meyer MH, Letscher-Bru V, Jaulhac B, Waller J, Candolfi E. Comparison of Mycosis IC/F and plus Aerobic/F media for diagnosis of fungemia by the Bactec 9240 system. *J Clin Microbiol.* 2004;42:773–7.
47. Mikulska M, Calandra T, Sanguinetti M, Poulain D, Viscoli C. The use of mannan antigen and anti-mannan antibodies in the diagnosis of invasive candidiasis: recommendations from the Third European Conference on Infections in Leukemia. *Crit Care.* 2010;14:R222.
48. Morace G, Borghi E. Fungal infections in ICU patients: epidemiology and the role of diagnostics. *Miner Anestesiol.* 2010;76:950–6.
49. Morrell M, Fraser VJ, Kollef MH. Delaying the empiric treatment of *Candida* bloodstream infection until positive blood culture results are obtained: a potential risk factor for hospital mortality. *Antimicrob Agents Chemother.* 2005;49:3640–5.
50. Nakao A, Yasui M, Kawagoe T, Tamura H, Tanaka S, Takagi H. False-positive endotoxemia derives from gauze glucan after hepatectomy for hepatocellular carcinoma with cirrhosis. *Hepatogastroenterology.* 1997;44:1413–8.
51. Neofytos D, Marr K. Diagnosis of invasive fungal disease. In: Safdar A, editor. *Management of infections in cancer patients.* New York: Springer; 2011. p. 261–72.
52. Odds FC, Bernaerts R. CHROMagar Candida, a new differential isolation medium for presumptive identification of clinically important *Candida* species. *J Clin Microbiol.* 1994;32:1923–9.
53. Reiss E, Shadomy HJ, Lyon GM. Part III: systemic mycoses caused by opportunistic yeasts and *Pneumocystis*. *Fundamental medical mycology.* Hoboken (New Jersey): John Wiley & Sons; 2011.
54. Rex JH, Pfaller MA, Walsh TJ, Chaturvedi V, Espinel-Ingroff A, Ghanoun MA, et al. Antifungal susceptibility testing: practical aspects and current challenges. *Clin Microbiol Rev.* 2001;14:643–58.
55. Rodriguez-Tudela JL, Donnelly JP, Pfaller MA, Chrysantou E, Warn P, Denning DW, et al. Statistical analyses of correlation between fluconazole MICs for *Candida* spp. assessed by standard methods set forth by the European Committee on Antimicrobial Susceptibility Testing (E.Dis. 7.1) and CLSI (M27-A2). *J Clin Microbiol.* 2007;45:109–11.
56. Rudensky B, Broidie E, Yinnon AM, Weitzman T, Paz E, Keller N, et al. Rapid flow-cytometric susceptibility testing of *Candida* species. *J Antimicrob Chemother.* 2005;55:106–9.
57. Sheppard DC, Locas MC, Restieri C, Laverdiere M. Utility of the germ tube test for direct identification of *Candida albicans* from positive blood culture bottles. *J Clin Microbiol.* 2008;46:3508–9.
58. Shin JH, Nolte FS, Morrison CJ. Rapid identification of *Candida* species in blood cultures by a clinically useful PCR method. *J Clin Microbiol.* 1997;35:1454–9.
59. St Germain G, Beauchesne D. Evaluation of the MicroScan Rapid Yeast Identification panel. *J Clin Microbiol.* 1991;29:2296–9.
60. Stone NR, Gorton RL, Barker K, Ramnarain P, Kibbler CC. Evaluation of PNA-FISH yeast traffic light for rapid identification of yeast directly from positive blood cultures and assessment of clinical impact. *J Clin Microbiol.* 2013;51:1301–2.
61. Subcommittee on Antifungal Susceptibility Testing (AFST) of the ESCMID European Committee for Antimicrobial Susceptibility Testing (EUCAST). EUCAST definitive document EDef 7.1: method for the determination of broth dilution

- MICs of antifungal agents for fermentative yeasts. *Clin Microbiol Infect.* 2008;14:398–405.
62. The University of Adelaide. Mycology online; 2011. Available from <http://www.mycology.adelaide.edu.au> [cited 15 August 2011].
63. Usami M, Ohata A, Horiuchi T, Nagasawa K, Wakabayashi T, Tanaka S. Positive (1→3)-beta-D-glucan in blood components and release of (1→3)-beta-D-glucan from depth-type membrane filters for blood processing. *Transfusion.* 2002;42:1189–95.
64. Vazquez JA, Sobel JD. Candidiasis. In: Kauffman CA, Pappas PG, Sobel JD, Dis-
mukes WE, editors. *Essentials of Clinical Mycology*. 2nd ed. New York: Springer; 2011. p. 167–206.
65. Wheat LJ. Approach to the diagnosis of invasive aspergillosis and candidiasis. *Clin Chest Med.* 2009;30:367–77, viii.
66. Yucesoy M, Marol S. Performance of CHROMAGAR Candida and BIGGY agar for identification of yeast species. *Ann Clin Microbiol Antimicrob.* 2003; 2:8.