



Original article

New lipid-dependent *Malassezia* species from parrots



F. Javier Cabañes^{a,*}, S. Dall'Acqua Coutinho^b, Laura Puig^a, M. Rosa Bragulat^a, Gemma Castellá^a

^a Veterinary Mycology Group, Department of Animal Health and Anatomy, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain

^b Laboratory of Molecular and Cellular Biology, Veterinary Medicine, Paulista University, São Paulo, Brazil

ARTICLE INFO

Article history:

Received 29 July 2015

Accepted 14 March 2016

Available online 13 May 2016

Keywords:

Malassezia brasiliensis

Malassezia psittaci

Birds

Parrots

Speciation

ABSTRACT

Background: All the currently recognized *Malassezia* species have been isolated from mammals. However, only a few of them have been isolated from birds. In fact, birds have been less frequently studied as carriers of *Malassezia* yeasts than mammals.

Aim: In this study we describe two new taxa, *Malassezia brasiliensis* sp. nov. and *Malassezia psittaci* sp. nov.

Methods: The isolates studied in this publication were isolated from pet parrots from Brazil. They were characterized using the current morphological and physiological identification scheme. DNA sequencing and analysis of the D1/D2 regions of the 26S rRNA gene, the ITS-5.8S rRNA gene sequences and the β -tubulin gene were also performed.

Results: The strains proposed as new species did not completely fit the phenotypic profiles of any the described species. The validation of these new species was supported by analysis of the genes studied. The multilocus sequence analysis of the three loci provides robust support to delineate these species.

Conclusions: These studies confirm the separation of these two new species from the other species of the genus *Malassezia*, as well as the presence of lipid-dependent *Malassezia* yeasts on parrots.

© 2016 Asociación Española de Micología. Published by Elsevier España, S.L.U. All rights reserved.

Nuevas especies lipodependientes del género *Malassezia* procedentes de loros

RESUMEN

Antecedentes: Todas las especies del género *Malassezia* actualmente identificadas se han aislado de mamíferos. Sin embargo, tan solo unas pocas de ellas se han aislado de aves. De hecho, las aves han sido estudiadas con menos frecuencia como portadoras de estas levaduras que los mamíferos.

Objetivos: En este estudio describimos dos nuevas especies del género *Malassezia*: *Malassezia brasiliensis* sp. nov. y *Malassezia psittaci* sp. nov.

Métodos: Las cepas estudiadas en esta publicación se aislaron de loros utilizados como animales de compañía en Brasil. Las cepas se caracterizaron mediante los criterios morfológicos y fisiológicos actualmente utilizados para la identificación de estas levaduras. También se llevó a cabo la secuenciación y el análisis de los fragmentos génicos D1/D2 26S e ITS-5.8S del ADN ribosómico y del gen de la β -tubulina.

Resultados: Los perfiles fenotípicos de las cepas propuestas como nuevas especies no encajaron completamente con los de las especies descritas en este género. Además, el análisis de los genes estudiados respaldó la validez de las nuevas especies. El análisis *multilocus* de secuencias de los tres *loci* estudiados reforzó con mayor firmeza la definición de las nuevas especies.

Conclusiones: Todos estos estudios confirman la separación de estas dos nuevas especies del resto de las especies descritas del género *Malassezia*, así como la existencia de especies dependientes de lípidos del género *Malassezia* en loros.

© 2016 Asociación Española de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Malassezia brasiliensis

Malassezia psittaci

Aves

Loros

Especiación

* Corresponding author.

E-mail address: javier.cabanes@uab.es (F.J. Cabañes).

Malassezia species are lipophilic basidiomycetous yeasts that inhabit the skin and mucosal sites of a variety of homoothermic animals, such as mammals or birds. In some conditions, these yeasts can be opportunistic pathogens of humans and animals.²⁹ This monophyletic genus is the only genus included in the order Malasseziales, which has an uncertain taxonomic position in the subphylum *Ustilagomycotina* (e.g., smut fungi). Recently, the class Malasseziomycetes has been proposed to accommodate these fungi.³¹ On the other hand, some culture-independent studies of fungi from environmental samples (e.g. Antarctic soils, corals, deep-sea sediments, nematodes) have shown that *Malassezia* are exceedingly widespread and ecologically diverse.¹ However, these yeasts have been only recovered from mammals and birds at the moment.

Nowadays *Malassezia* yeasts have been isolated mainly from domestic animals, different wild animals held in captivity, and also from wildlife.²⁹ However, the occurrence of *Malassezia* yeasts on the skin of most animals remains unknown. At present, the genus *Malassezia* includes 14 species. Most of them are lipid-dependent yeasts which require long-chain fatty acids to grow, while the lipophilic but non-lipid-dependent species *Malassezia pachydermatis* is the only species in the genus that does not require lipid supplementation for development in culture medium.⁵

These species have been isolated mainly from mammals, and only a few of them have been isolated from birds. In fact, birds have been less frequently studied as carriers of *Malassezia* yeasts than mammals. Therefore, few papers have been published about these yeasts from birds. A search in PubMed (www.ncbi.nlm.nih.gov/pubmed; US National Library of Medicine) early in 2015 yielded more than 1500 journal citations containing the keywords “*Malassezia* AND mammals”, but only 7 using the keywords “*Malassezia* AND birds”.

Besides, little is known about *Malassezia* yeasts isolated from birds.²⁹ At the moment, only *M. pachydermatis*, *Malassezia furfur* and *Malassezia sympodialis* have been reported from these animals. As far as *M. pachydermatis* is concerned, this non lipid-dependent species has been reported to be isolated from feathers of a variety of birds,¹¹ from a diseased throat of a scarlet macaw² and from several

wild bird droppings.²³ Some strains of the lipid-dependent species *M. furfur* were isolated from the wing and mouth of ostriches and from the wing of a pelican,¹⁵ and also from pigeon droppings.⁸ The lipid-dependent species *M. sympodialis* was the most frequently yeast isolated from clinically altered combs of adult chickens.¹⁸ In this study, the isolates were identified only in the basis of phenotypic characteristics without DNA sequencing confirmation. However, *Malassezia* yeasts have been isolated from birds, including parrots, in a previous study.²⁴ Nonetheless, in this study, the isolates were not identified at species level. On the other hand, some studies have failed to demonstrate the presence of lipophilic yeasts in these animals. *Malassezia* yeasts were not detected on the skin of healthy psittacine birds and neither on psittacine birds with feather-destructive behavior.²⁵

In this paper, we describe two new lipid-dependent species in the genus *Malassezia* isolated from parrots. For these isolates we propose the names *Malassezia brasiliensis* sp. nov. and *Malassezia psittaci* sp. nov.

Materials and methods

Strains

The strains proposed as new species in this publication were isolated from pet parrots from Brazil. Four isolates were recovered in the Laboratory of Molecular and Cellular Biology of the Paulista University (São Paulo, Brazil) from four individual parrots (Table 1). They were isolated from lesions on beak (MA 1453, *Amazona aestiva*; MA 1454, *Pionus menstruus*; MA 1455, *A. aestiva*) and oropharynx (MA 1456, *Nymphicus hollandicus*) in these animals. Unfortunately the etiological significance of these isolates in these lesions was not determined. The strains were stored at -80°C .⁹

Morphological and physiological characterization

The characterization of lipid-dependent yeasts was based on their inability to grow on Sabouraud's glucose agar (SGA; Oxoid, Basingstoke, UK), on their ability to use certain polyoxyethylene

Table 1
Strains studied, their hosts and GenBank accession numbers.

Strains	Host	GenBank accession no.		
		D1/D2	ITS	β-Tubulin
<i>M. caprae</i> CBS 10434 ^T	Goat	AY743616	AY743656	KC573795
<i>M. cuniculi</i> CBS 11721 ^T	Rabbit	GU733708	GU733709	KC573808
<i>M. dermatis</i> CBS 9169 ^T	Human	AB070361	AY390284	KC573796
<i>M. equina</i> CBS 9969 ^T	Horse	AY743621	AY743641	KC573798
<i>M. furfur</i> CBS 1878 ^{NT}	Human	AY743602	AY743634	KC573799
<i>M. furfur</i> CBS 7019 ^{NT}	Human	AY743603	AY743635	KC573800
<i>M. furfur</i> CBS 7984	Elephant	AY387226	AY387130	–
<i>M. furfur</i> CBS 7985	Ostrich	AY387225	AY387129	–
<i>M. furfur</i> CBS 9366	Human	AY387203	AY387107	–
<i>M. furfur</i> CBS 9368	Human	AY387205	AY387109	–
<i>M. furfur</i> MA 1453	Parrot	KC57390	KC573787	KR872309
<i>M. furfur</i> MA 1456	Parrot	KC57390	KC573787	KC573811
<i>M. globosa</i> CBS 7966 ^T	Human	AY743604	AY387132	KC573806
<i>M. japonica</i> CBS 9431 ^T	Human	EF140672	EF140669	KC573801
<i>M. nana</i> CBS 9557 ^T	Cat	EF140671	EF140666	HM594270
<i>M. obtusa</i> CBS 7876 ^T	Human	AY743629	AY387137	KC573802
<i>M. pachydermatis</i> CBS 1879 ^{NT}	Dog	AY743605	AY387139	KC573803
<i>M. restricta</i> CBS 7877 ^T	Human	AY743607	AY387143	KC573807
<i>M. slooffiae</i> CBS 7956 ^T	Pig	AY743606	AY387146	KC573805
<i>M. sympodialis</i> CBS 7222 ^T	Human	AY743626	AY387157	KC573797
<i>M. yamatoensis</i> CBS 9725 ^T	Human	AB125263	AB125261	KC573804
<i>M. yamatoensis</i> CBS 9726	Human	AB125264	AB125262	–
<i>M. brasiliensis</i> sp. nov. MA 1455	Parrot	KR872314	KR872315	KR872311
<i>M. psittaci</i> sp. nov. MA 1454	Parrot	KR872312	KR872313	KR872310

CBS, Centraalbureau voor Schimmelcultures; MA, culture collection of the Veterinary Mycology group. Prefixes KR of accession numbers correspond to the sequences generated in this study.

Table 2
Main phenotypical characteristics of *Malassezia* species.^a

Species	Cell morphology	Growth on mDA	Growth on SGA	T 20 ^e	T 40 ^e	T 60 ^e	T 80 ^e	Cremophor EL	Catalase	β-Glucosidase	Growth at 37 °C	Growth at 40 °C
<i>M. caprae</i>	Globose, ellipsoidal	+	–	– ^c (+)	+ ^b	+ ^b	+ ^b , (–)	– ^d	+	+, (–)	–, (w)	–
<i>M. cuniculi</i>	Globose	–, (w)	–	–	–	–	–	–	+	+	+	+
<i>M. dermatitis</i>	Ellipsoidal, globose	+	–	+	+	+	+, (w)	–	–	–	+	–
<i>M. equina</i>	Ellipsoidal	+	–	w ^c	+ ^d	+ ^{b,d}	+ ^b	–	+	–	w	–
<i>M. furfur</i>	Globose, ellipsoidal, cylindrical	+	–	+	+	+	+	+	+	+	+	+
<i>M. globosa</i>	Globose	+	–	– ^c	–	–	– ^c	–	+	–	–, (w)	–
<i>M. japonica</i>	Globose, ellipsoidal	+	–	– ^c	– ^c	+	+	w	+	+	+	–
<i>M. nana</i>	Ellipsoidal	+	–	– ^c	+	+	+	–	+	+	+	–
<i>M. obtusa</i>	Ellipsoidal, cylindrical	+	–	–	– ^d	– ^d	–	–	+	+	+	–
<i>M. pachydermatis</i>	Ellipsoidal	+	+	+ ^b	+ ^b	+	+	+ ^b	+, w, (–)	+, (–)	+	+
<i>M. restricta</i>	Globose, ellipsoidal	+	–	–	– ^d	– ^d	–	–	–	–	v	–
<i>M. slooffiae</i>	Ellipsoidal, cylindrical	+	–	+	+	+	w	–	+	–	+	+
<i>M. sympodialis</i>	Ellipsoidal	+	–	–, w ^c	+	+	+	–, (w)	+	+	+	+
<i>M. yamatoensis</i>	Ellipsoidal	+	–	+	+	+	+	+, (w)	+	–	+	–
<i>M. brasiliensis</i> sp. nov.	Ovoidal, ellipsoidal	+	–	+	+	+	+	+	+	–	+	+
<i>M. psittaci</i> sp. nov.	Globose, ovoidal	+	–	+	+	+	+	+	+	–	–	–

^a With the exception of *M. brasiliensis* and *M. psittaci* data are from Guého et al.¹⁴ and Cabañes et al.⁴; +, positive; –, negative; v, variable; w, weak; () indicate rare deviations from main pattern.

^b Growth may be inhibited near the well where the substrate is placed.

^c Growth may occur at some distance from the well where the substrate is placed.

^d Opaque zone may occur.

^e Tween diffusion test proposed by Guillot et al.¹⁶

sorbitanesters (Tweens 20, 40, 60 and 80), and the use of additional tests such as the Cremophor EL assimilation test, or the splitting of esculin (β-glucosidase activity), following the identification schemes of Guého et al.¹⁴ Other tests, such as the catalase reaction, growth at different temperatures (32 °C, 37 °C, 40 °C and 45 °C) on modified Dixon agar (mDA; 36 g malt extract, 10 g peptone, 20 g desiccated ox-bile, 10 ml Tween 40, 2 ml glycerol, 2 ml oleic acid and 12 g agar per liter, pH 6.0), and the morphological characteristics after incubation at 32 °C for 7 days in the same culture medium were also performed.¹⁴

DNA extraction, gene amplification, sequencing and phylogenetic analysis

DNA of the isolates MA 1453, MA 1454 and MA 1455 was extracted and purified directly from seven day-old cultures on modified Dixon agar according to the FastDNA Spin kit protocol with the FastPrep FP-24 instrument (MP Biomedicals, Biolink, Barcelona, Spain). The DNA was kept at –20 °C until used as a template for PCR amplification. Sequences of D1/D2 26S rRNA (D1/D2), ITS-5.8S rRNA (ITS) and β-tubulin (β-tub) genes of currently recognized species and the isolate MA 1456 were obtained from previous studies.^{3,4,6} D1/D2, ITS, and β-tub genes of the strains isolated from parrots were amplified and sequenced as described previously.⁶

Sequence alignments were performed using the software program Clustal X v2.0.12.²² Regions of ambiguous alignment were removed with Gblocks.⁷ Parsimony analyses of the individual and combined data matrices were conducted using PAUP* version 4.0b10 software.³⁰ One hundred heuristic searches were conducted with random sequence addition and tree bisection reconnection branch-swapping algorithms, collapsing zero-length branches and

saving all minimal-length trees (MulTrees) on different data sets. The gaps were treated as missing data, and support of internal branches was assessed using heuristic parsimony search of 1000 bootstrapped data sets. Tree length, consistency index (CI), retention index (RI) and the homoplasy index (HI) values were also calculated. The combined data set was tested for incongruence with the partition homogeneity test (PHT) as implemented in PAUP*.

Results

Morphology and physiology

The phenotypic characteristics of the new species, *M. brasiliensis* (MA 1455) and *M. psittaci* (MA 1454) and the other currently accepted *Malassezia* species, are summarized in Table 2. None of these isolates grew on SGA without any lipid supplementation. *M. brasiliensis* cells were ovoidal to ellipsoidal (Table 2), and *M. psittaci* cells were globose to ovoidal (Table 2).

All the isolates studied have good growth on mDA at 32 °C. The isolates MA 1453 (2.4–5.1 mm in diameter; average diameter = 4.2 mm) and MA 1456 (3.6–6.4 mm in diameter; average diameter = 5.4 mm) grew faster and formed larger colonies than MA 1455 (1.6–3.2 mm in diameter; average diameter = 2.5 mm) and MA 1454 (1.9–3 mm in diameter; average diameter = 2.6 mm) on mDA at 32 °C after 7 days of incubation. With the exception of MA 1454 (*M. psittaci*) that did not grow at 37 °C, the rest of the isolates showed good growth at both 37 °C and 40 °C after 7 days of incubation, but did not grow at 45 °C. All the isolates had similar good growth around Tweens 20, 40, 60 and 80, and Cremophor EL. All, except one (MA 1453), were β-glucosidase negative.




Fig. 1. Molecular phylogenetic tree inferred from a parsimony analysis of D1/D2 26S rRNA sequences of members of the genus *Malassezia*. MP bootstrap values >50% in 1000 replications are shown at nodes. The tree is rooted with *Cryptococcus neoformans*. CI, consistency index; RI, retention index; HI, homoplasy index.

Molecular and phylogenetic analysis

With the primers used we were able to amplify and sequence 619 bp, 717–829 bp and 950–1115 bp of the D1/D2, the ITS and the β -tub genes, respectively. The nucleotide sequences determined in this study have been deposited at the GenBank database under accession numbers KR872309–KR872315 (Table 1). Figs. 1, 2 and 3 show the molecular phylogenetic trees based on the maximum parsimony analysis of the sequences of D1/D2, ITS and β -tub, respectively. When the three loci were combined (Fig. 4), the data set included 1943 characters. Each sequence contributed to that length as follows: D1/D2, 501 characters; ITS, 599 characters; and β -tub, 843 characters. From these characters, 921 were constant, and 643 were parsimony informative (D1/D2 84, ITS 263 and β -tub 296). The result of the partition homogeneity test ($p = 0.01$) showed that the data set could be combined without reducing phylogenetic accuracy.¹⁰

The isolates MA 1453 and MA 1456 clustered with *M. furfur* strains. These isolates had identical D1/D2 and ITS sequences. The similarity of these sequences was 100% between these isolates and *M. furfur* CBS 7985, which was isolated from an ostrich. A different topology was observed using β -tub sequencing. The isolates

MA 1453 and MA 1456 formed a different clade from the *M. furfur* neotype species.

The isolate MA 1455, belonging to the novel proposed species *M. brasiliensis*, clustered close to the *M. furfur* reference strains, but in a different clade with high support in most of the individual gene trees (D1/D2: 67%; ITS: 99%; β -tub: 98%). Dissimilarities between this isolate and *M. furfur* CBS 1878NT and *M. furfur* CBS 7019NT in D1/D2 sequences were 1.4% and 1%, respectively. Besides, dissimilarities between MA 1455 and *M. furfur* CBS 1878NT and *M. furfur* CBS 7019NT in ITS and β -tub sequences were 4.8% and 5.3%, and 4.3% and 4.5%, respectively.

The isolate MA 1454, belonging to the novel proposed species *M. psittaci*, clustered close to the type strain of *Malassezia yamatoensis*, but in a different clade with high support (100%) in all the individual gene trees. The dissimilarity between this isolate and *M. yamatoensis* CBS 9725T in D1/D2 sequences was 1.7%. Besides, dissimilarities between MA 1454 and *M. yamatoensis* CBS 9725T in ITS and β -tub sequences were 3.6% and 5.9%, respectively.

Both *M. brasiliensis* and *M. psittaci* received strong support (100% bootstrap value) as distinct clades in the combined D1/D2-ITS- β -tub MP tree. They were clearly separated from *M. furfur* and *M. yamatoensis*, respectively.




Fig. 2. Molecular phylogenetic tree inferred from a parsimony analysis of ITS-5.8S rDNA sequences of members of the genus *Malassezia*. MP bootstrap values >50% in 1000 replications are shown at nodes. The tree is rooted with *Cryptococcus neoformans*. CI, consistency index; RI, retention index; HI, homoplasy index.

Description of the proposed new species

M. brasiliensis Cabañes, Coutinho, Bragulat et Castellá, sp. nov. Mycobank MB 812624

M. brasiliensis (*brasiliensis*: bra.si.li.en'sis M.L. adj, brasiliensis pertaining to Brazil, South America; this Latin-derived species epithet refers to the country from which the yeast was first isolated).

On mDA, after 7 days at 32 °C, colonies are large (average diameter 2.5 mm, 1.6–3.2 mm), whitish to cream-colored, smooth, dull, butyrous, moderately convex and slightly elevated in the center with entire margins. Cells are ovoidal to ellipsoidal, 2.8–3.9 μm × 1.9–3.3 μm, with buds formed monopolarly on a broad base. No growth is obtained on SGA. Catalase reaction is positive, and β-glucosidase activity is negative. Growth occurs on glucose-peptone agar with Tween-20, Tween-40, Tween-60, Tween-80 and Cremophor EL as sole source of lipid. Good growth appears at 37 °C and 40 °C and no growth occurs at 45 °C. The teleomorph is unknown.

The type strain CBS 14135 (= CECT 13138; originally strain MA 1455) was isolated from lesions on the beak of a parrot (*A. aestiva*)

in São Paulo, Brazil. The strains were deposited in the CBS Fungal Biodiversity Centre, Utrecht, The Netherlands, and in the Spanish Type Culture Collection, Valencia, Spain, as CBS 14135 and CECT 13138, respectively.

M. psittaci Cabañes, Coutinho, Bragulat et Castellá, sp. nov. Mycobank MB 812625

M. psittaci (*psittaci*: psit'ta.ci. L. n. N. *psittacus* a parrot; M.L. gen. n. *psittaci* of a parrot; this Latin-derived species epithet refers to the host animal from which the yeast was first isolated).

On mDA, after 7 days at 32 °C, colonies are large (average diameter 2.6 mm, 1.9–3 mm), whitish to cream-colored, smooth, shiny, butyrous, moderately convex and slightly elevated in the center with entire margins. Cells are globose to ovoidal, 2.5–4.4 μm × 2.2–3.4 μm, with buds formed monopolarly on a broad base. No growth is obtained on SGA. Catalase reaction is positive, and β-glucosidase activity is negative. Growth occurs on glucose-peptone agar with Tween-20, Tween-40, Tween-60, Tween-80 and Cremophor EL as sole source of lipid. No growth appears at 37 °C. The teleomorph is unknown.




Fig. 3. Molecular phylogenetic tree inferred from a parsimony analysis of β -tubulin sequences of members of the genus *Malassezia*. MP bootstrap values >50% in 1000 replications are shown at nodes. The tree is rooted with *Cryptococcus neoformans*. CI, consistency index; RI, retention index; HI, homoplasy index.

The type strain CBS 14136 (=CECT 13137; originally strain MA 1454) was isolated from lesions on the beak of a parrot (*P. menstruus*) in São Paulo, Brazil. The strains were deposited in the CBS Fungal Biodiversity Centre, Utrecht, The Netherlands, and in the Spanish Type Culture Collection, Valencia, Spain, as CBS 14136 and CECT 13137, respectively.

Discussion

The number of recognized *Malassezia* species described in the present century ($n=7$) is the same that the number of recognized *Malassezia* species described in the two last centuries ($n=7$). In fact, with the exception of *M. furfur* (Robin) Baillon (1889) and *M. pachydermatis* (Weidman) Dodge (1925), most of the species were described in the last decade of the 20th century on the basis of morphological, physiological and molecular studies which were already available for these purposes.^{13,28}

Lipid-dependent *Malassezia* yeasts require complex media enriched with lipids. These particular requirements (e.g. ability to use Tweens 20, 40, 60 and 80) were found to be useful to separate *Malassezia* species.^{13,16} However, the addition of the last new species into this genus have resulted in similar physiological patterns among several species, and thus in a doubtful identification. In these cases, the identification should be confirmed by DNA sequencing analysis (e.g. D1/D2 and ITS).¹⁴

Malassezia yeasts have been isolated from almost all domestic animals, different wild animals held in captivity, and also from wildlife.²⁹ Despite this, the occurrence of *Malassezia* yeasts on the skin of most animals remains unknown. Besides, in a recent study of the occurrence of fungal species in human skin,¹² DNA

sequences that may represent unidentified *Malassezia* species were also detected. These facts made it possible to anticipate an increase in the number of new species in this genus, particularly if wild animals are studied further.⁵

This increase may difficult the description of future new species in this genus if we only use the classical techniques of phenotypical characterization (e.g. ability to use Tweens 20, 40, 60 and 80) and the present molecular “gold standard” D1/D2 gene to determine *Malassezia* species. Various molecular markers had been used to achieve discrimination among the recognized species of *Malassezia*. The most studied loci had been the D1/D2 and the ITS. In addition to the sequencing of the rRNA genes, other genes such as the chitin synthase-2 gene (*CHS2*)²⁰ and the β -tub gene⁶ have been proposed for taxonomic purposes in this genus.

In other fungal genera, such as *Penicillium* and *Aspergillus*, the molecular “gold standard” ITS gene does not always distinguish different species. Protein coding genes are widely used in mycology for the identification and have generally a higher interspecies variability than the ITS region. There is no standard choice of a protein-coding gene in the fungal kingdom, but β -tub and calmodulin sequences are frequently used for the identification of *Aspergillus* and *Penicillium* species and are better species markers than ITS.¹⁹ On the other hand, new tools such as MALDI-TOF mass spectrometry,²¹ may be useful for a proper characterization of future new species in the genus *Malassezia*.

The isolates MA 1453 and MA 1456 had both very similar physiological patterns and D1/D2 sequences to *M. furfur*. However, they form a different clade from the *M. furfur* neotype strains using both the β -tub sequencing and the multilocus analysis. Only a few strains of *M. furfur* from birds have been studied in depth. On the




Fig. 4. Molecular phylogenetic tree inferred from a parsimony analysis of D1/D2 26S rRNA, ITS-5.8S rRNA and β -tubulin gene sequences of members of the genus *Malassezia*. MP bootstrap values >50% in 1000 replications are shown at nodes. The tree is rooted with *Cryptococcus neoformans*. CI, consistency index; RI, retention index; HI, homoplasy index.

other hand, a considerable genetic diversity within this species has been observed. Four distinct LSU rRNA sequences have been reported in this species.¹⁵ Three isolates from ostriches (e.g. *M. furfur* JG 590 = CBS 7985) showed the most common LSU rRNA sequence (sequence “a”) together with other human and animal isolates. Two isolates from an elephant (JG 570 = CBS 7984) and a pelican (JG 593) clustered together and showed a unique LSU rRNA sequence (sequence “b”). Eight well-distinguished subtypes in this species were revealed using AFLP.¹⁷ Isolates from animals clustered in two subtypes. Subtype 2 contained two human isolates and three isolates from animals (e.g. elephant, elk) and subtype 8 only two isolates from an elephant (JG 570 = CBS 7984) and an ostrich (JG 590 = CBS 7985).

In our study, the topology of some species was not always concordant between the three markers investigated. This has been also described in other studies in *Malassezia* species.^{3,6,14,20} Some species tend to jump between clades depending on the gene analyzed and the set of other fungi included. This implies that further phylogenetic research is needed to establish the position of these species in the tree of life.¹⁴ A further explanation of these incongruent results may be that hybridization may have occurred during speciation, and cell fusion, karyogamy and meiosis may be possible within the genus.³ In the case of speciation through clonal divergence and genetic drift, probably followed by some host adaptation, one would expect concordance between the phylogenetic patterns of each individual gene. So, the lack of concordance may indicate that recombination has played a role in the divergence of these species. This is particularly interesting, as sexual reproduction is

unknown in *Malassezia*. Recently, a region corresponding to the mating type locus (MAT) has been identified for these yeasts.²⁶

On the other hand, in the present investigation we propose two new *Malassezia* species. Among other differences, the new species *M. brasiliensis* can be distinguished from *M. pachydermatis* by its inability to grow in SGA, from *Malassezia caprae*, *Malassezia dermatitis*, *Malassezia equina*, *Malassezia japonica*, *Malassezia nana*, *M. psittaci* and *M. yamatoensis* by its ability to grow at 40 °C, from *Malassezia cuniculi*, *Malassezia globosa* and *Malassezia obtusa* by its ability to assimilate Tween-20, Tween-40, Tween-60 and Tween-80, from *Malassezia slooffiae* and *M. sympodialis* by its ability to assimilate Cremophor EL, from *Malassezia restricta* by its catalase activity, and from *M. furfur* by its β -glucosidase activity. The new species *M. psittaci* can be distinguished from *M. pachydermatis* by its inability to grow in SGA, from *M. brasiliensis*, *M. dermatitis*, *M. furfur*, *M. japonica*, *M. nana*, *M. slooffiae*, *M. sympodialis* and *M. yamatoensis* by its inability to grow at 37 °C, from *M. cuniculi*, *M. globosa* and *M. obtusa* by its ability to assimilate Tween-20, Tween-40, Tween-60 and Tween-80, from *M. caprae* and *M. equina* by its ability to assimilate Cremophor EL, and from *M. restricta* by its catalase activity.

In addition, the validation of these new species was supported by the analysis of the D1/D2, the ITS and the β -tub genes which confirmed the separation of these new species from the other described species of the genus. Although a boundary between species (a prerequisite number of nucleotide differences in either regions to separate species) has been not clearly stated, the within-species D1/D2 sequence similarity is above 99% for *Malassezia* species (<1% of dissimilarity).^{17,29} In our study, the multilocus sequence analysis

of the three loci provides also robust support to delineate these new species. So, phylogenetic analysis of sequences from *M. brasiliensis* and *M. psittaci* showed that they were clearly distinct from the other 14 described *Malassezia* species, exceeding the variation generally observed to occur between basidiomycetous yeast species.²⁷ On the other hand, our study confirms also the presence of lipid-dependent *Malassezia* yeasts on parrots.

Conflict of interest

The authors have no conflict of interest.

Acknowledgements

The authors thank Lab&Vet Diagnóstico e Consultoria Veterinària for kindly providing the parrot samples and Suzana Maria Bezerra from the Laboratory of Molecular and Cellular Biology of the Paulista University (PU) and Carolina Gómez from the Veterinary Mycology Group of the Universitat Autònoma de Barcelona (UAB) for their valuable technical assistance. Financial support came from Servei Veterinari de Bacteriologia i Micologia of the UAB and from the Research Section of the PU.

References

- Amend A. From dandruff to deep-sea vents: *Malassezia*-like fungi are ecologically hyper-diverse. *PLoS Pathog.* 2014;10:e1004277.
- Breuer-Strosberg R, Hochleithner M, Kuttin E. *Malassezia pachydermatis* isolation from a scarlet macaw. *Mycoses.* 1990;33:247–50.
- Cabañes FJ, Theelen B, Castellá G, Boekhout T. Two new lipid-dependent *Malassezia* species from domestic animals. *FEMS Yeast Res.* 2007;7:1064–76.
- Cabañes FJ, Vega S, Castellá G. *Malassezia cuniculi* sp. nov., a novel yeast species isolated from rabbit skin. *Med Mycol.* 2011;49:40–8.
- Cabañes FJ. *Malassezia* yeasts: how many species infect humans and animals? *PLoS Pathog.* 2014;10:e1003892.
- Castellá G, Coutinho SDA, Cabañes FJ. Phylogenetic relationships of *Malassezia* species based on multilocus sequence analysis. *Med Mycol.* 2014;52:99–105.
- Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol.* 2000;17:540–52.
- Chae HS, Jang GE, Kim NH, Son HR, Lee JH, Kim SH, et al. Classification of *Cryptococcus neoformans* and yeast-like fungus isolates from pigeon droppings by colony phenotyping and ITS genotyping and their seasonal variations in Korea. *Avian Dis.* 2012;56:58–64.
- Crespo MJ, Abarca ML, Cabañes FJ. Evaluation of different preservation and storage methods for *Malassezia* spp. *J Clin Microbiol.* 2000;38:3872–5.
- Cunningham CW. Can three incongruence tests predict when data should be combined? *Mol Biol Evol.* 1997;14:733–40.
- Dufait R. Presence de *Malassezia pachydermatis* (syn. *Pityrosporum canis*) sur le poils et les plumes d'animaux domestiques. *Bull Soc Fr Mycol Méd.* 1985;14:19–22.
- Findley K, Oh J, Yang J, Conlan S, Deming C, Meyer JA, et al. Topographic diversity of fungal and bacterial communities in human skin. *Nature.* 2013;498:367–70.
- Guého E, Midgley G, Guillot J. The genus *Malassezia* with description of four new species. *Antonie Van Leeuwenhoek.* 1996;69:337–55.
- Guého E, Boekhout T, Begerow D. Biodiversity, phylogeny and ultrastructure. In: Boekhout T, Guého E, Mayser P, Velegriaki A, editors. *Malassezia and the skin.* Berlin: Springer; 2010. p. 17–63.
- Guillot J, Guého E. The diversity of *Malassezia* yeasts confirmed by rRNA sequence and nuclear DNA comparisons. *Antonie Van Leeuwenhoek.* 1995;67:297–314.
- Guillot J, Guého E, Lesourd M, Midgley G, Chevrier G, Dupont B. Identification of *Malassezia* species, a practical approach. *J Mycol Méd.* 1996;6:103–10.
- Gupta AK, Boekhout T, Theelen B, Summerbell RC, Batra R. Identification and typing of *Malassezia* species by amplified fragment length polymorphism (AFLP) and sequence analyses of the internal transcribed spacer (ITS) and large subunit (LSU) regions of ribosomal DNA. *J Clin Microbiol.* 2004;42:4253–60.
- Gründer S, Mayser, Redmann T, Kaleta E. Mycological examinations on the fungal flora of the chicken comb. *Mycoses.* 2005;48:114–9.
- Houbraken J, de Vries RP, Samson RA. Modern taxonomy of biotechnologically important *Aspergillus* and *Penicillium* species. *Adv Appl Microbiol.* 2014;86:199–249.
- Kano R, Aizawa T, Nakamura Y, Watanabe S, Hasegawa A. Chitin synthase 2 gene sequence of *Malassezia* species. *Microbiol Immunol.* 1999;43:813–5.
- Kolecka A, Khayhan K, Arabatzis M, Velegriaki A, Kostrzewa M, Andersson A, et al. Efficient identification of *Malassezia* yeasts by matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS). *Br J Dermatol.* 2014;170:332–41.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al. Clustal W and Clustal X version 2.0. *Bioinformatics.* 2007;23:2947–8.
- Mendes JF, Albano AP, Coimbra MA, Ferreira GF, Gonçalves CL, Nascente Pda S, et al. Fungi isolated from the excreta of wild birds in screening centers in Pelotas, RS, Brazil. *Rev Inst Med Trop Sao Paulo.* 2014;56:525–8.
- Midgley G, Clayton Y. The yeast flora of birds and mammals in captivity. *Antonie Van Leeuwenhoek.* 1969;35 Suppl.:E23–4.
- Preziosi D, Morris D, Johnston M, Rosenthal K, O'Shea K, Rankin S. Distribution of *Malassezia* organisms on the skin of unaffected psittacine birds and psittacine birds with feather-destructive behavior. *J Am Vet Med Assoc.* 2006;228:216–20.
- Saunders CW, Scheynius A, Heitman J. *Malassezia* fungi are specialized to live on skin and associated with dandruff, eczema, and other skin diseases. *PLoS Pathog.* 2012;8:e1002701.
- Scorzetti G, Fell JW, Fonseca A, Statzell-Tallman A. Systematics of basidiomycetous yeasts: a comparison of large subunit D1/D2 and internal transcribed spacer rDNA regions. *FEMS Yeast Res.* 2002;2:495–517.
- Simmons RB, Guého E. A new species of *Malassezia*. *Mycol Res.* 1990;94:1146–9.
- Sugita T, Boekhout T, Velegriaki A, Guillot J, Hadina S, Cabañes FJ. Epidemiology of *Malassezia*-related skin diseases. In: Boekhout T, Guého E, Mayser P, Velegriaki A, editors. *Malassezia and the skin.* Berlin: Springer; 2010. p. 63–119.
- Swofford DL. PAUP*. Phylogenetic analysis using parsimony (*and other methods). Sunderland: Sinauer Associates; 2002.
- Wang QM, Theelen B, Groenewald M, Bai FY, Boekhout T. *Moniliellomyces* and *Malasseziomyces*, two new classes in *Ustilaginomycotina*. *Persoonia.* 2014;33:41–7.