



Original article

Prevalence of *Candida albicans*, *Candida dubliniensis* and *Candida africana* in pregnant women suffering from vulvovaginal candidiasis in Argentina



María Josefina Mucci^a, María Luján Cuestas^b, María Fernanda Landanburu^b, María Teresa Mujica^{b,*}

^a Hospital Materno Infantil Ramón Sardá, Esteban de Luca 2151, CP 1246 Buenos Aires, Argentina

^b Instituto de Microbiología y Parasitología Médica, Universidad de Buenos Aires-Consejo Nacional de Investigaciones Científicas y Técnicas (IMPaM, UBA-CONICET), Paraguay 2155, piso 11, CP 1121ABG Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 14 August 2015

Accepted 13 September 2016

Available online 3 April 2017

Keywords:

Candida albicans

Candida dubliniensis

Vulvovaginal candidiasis

Candida africana

ABSTRACT

Background: Vulvovaginal candidiasis (VVC) is a vulvovaginitis commonly diagnosed in gynecology care. In recent years, the taxonomy of the most important pathogenic *Candida* species, such as *Candida albicans* have undergone significant changes.

Aims: This study examined the prevalence of *C. albicans*, *Candida africana*, and *Candida dubliniensis* in vaginal specimens from 210 pregnant women suffering from vulvovaginitis or having asymptomatic colonization.

Methods: Phenotypic and molecular methods were used for the identification of the species.

Results: During the studied period, 55 isolates of *Candida* or other yeasts were obtained from specimens collected from 52 patients suffering from vulvovaginitis (24.8%). *C. albicans* was the predominant *Candida* species in 42 isolates (80.7%), either alone or in combination with other species of the genus (5.7%, n = 3). Additionally, nine isolates of *C. albicans* (50%) were obtained from asymptomatic patients (n = 18). *C. dubliniensis* was the causative agent in 2 (3.8%) cases of VVC, and was also isolated in one asymptomatic patient. Molecular assays were carried out using specific PCR to amplify the *ACT1*-associated intron sequence of *C. dubliniensis*. The amplification of the *HWP1* gene also correctly identified isolates of the species *C. albicans* and *C. dubliniensis*. No *C. africana* was isolated in this work. Some *C. albicans* isolates were either homozygous or heterozygous at the *HWP1* locus. The distribution of heterozygous and homozygous *C. albicans* isolates at the *HWP1* locus was very similar among patients suffering from VVC and asymptomatic patients ($p = 0.897$).

Conclusions: The presence of *C. albicans* and *C. dubliniensis*, and the absence of *C. africana* in pregnant is noteworthy.

© 2016 Asociación Española de Micología. Published by Elsevier España, S.L.U. All rights reserved.

Prevalencia de *Candida albicans*, *Candida dubliniensis* y *Candida africana* en mujeres gestantes con candidiasis vulvovaginal, en Argentina

RESUMEN

Palabras clave:

Candida albicans

Candida dubliniensis

Candidiasis vulvovaginal

Candida africana

Antecedentes: La candidiasis vulvovaginal (CVV) es una vulvovaginitis comúnmente diagnosticada en la práctica ginecológica. En los años recientes la taxonomía de *Candida albicans* ha sufrido cambios significativos.

Objetivos: En este estudio se examinó la prevalencia de *C. albicans*, *Candida africana* y *Candida dubliniensis* en las secreciones vaginales de 210 mujeres embarazadas con vulvovaginitis o colonización asintomática.

Métodos: Se usaron métodos fenotípicos y moleculares para la identificación de las levaduras.

* Corresponding author.

E-mail address: mtmujica@gmail.com (M.T. Mujica).

Resultados: Un total de 55 aislamientos de *Candida* u otras levaduras se recuperaron a partir de muestras de 52 pacientes con vulvovaginitis (24,8%). La especie *C. albicans* fue predominante como especie única (42 aislamientos; 80,7%) o asociada con otras especies del género (5,7%, n=3). Nueve aislamientos de *C. albicans* (50%) se recuperaron de las pacientes asintomáticas (n=18). Se aisló *C. dubliniensis* en dos casos de CVV (3,8%) y en una paciente asintomática. Los estudios moleculares por PCR que amplifican la secuencia del intrón asociado al gen *ACT1* confirmaron la presencia de esta especie. La amplificación del gen *HWP1* confirmó la presencia de *C. dubliniensis* y *C. albicans*, y la ausencia de *C. africana*. Los aislamientos de *C. albicans* fueron homocigóticos o heterocigóticos en el locus *HWP1*; las pacientes con CVV y aquellas asintomáticas mostraron para esta especie una distribución similar de aislamientos homocigóticos o heterocigóticos en el locus *HWP1* ($p = 0,897$).

Conclusiones: Destacamos la presencia de *C. albicans* y *C. dubliniensis* y la ausencia de *C. africana* en las CVV o en la colonización asintomática de las secreciones vaginales de las gestantes.

© 2016 Asociación Española de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Vulvovaginal candidiasis (VVC) is caused by abnormal yeast growth in the mucosa of the female genital tract and is one of the most common conditions diagnosed in women seeking gynecological care.⁹ The distribution of the different *Candida* species identified in women suffering from VVC varies widely depending on the locations as well as the populations studied.² It was reported that 80–90% of VVC cases are caused by *Candida albicans* whereas 10–20% are due to other species of the genus.^{2,25} However, in recent years, the taxonomy of the most important *Candida* species such as *C. albicans*, *Candida parapsilosis* and *Candida glabrata*, has undergone significant changes due to the description of new closely related species and therefore they are, nowadays, recognized as ‘cryptic species complexes’.^{4,6,7,20,21,28,31–34} *Candida dubliniensis* is a yeast species that is closely related to *C. albicans*,²⁸ and has been found in vaginal samples.^{1,28,35} The very close phenotypic and genotypic relationship between *C. dubliniensis* and *C. albicans* has led to the misidentification of isolates of *C. dubliniensis* as *C. albicans*.^{5,7} Recently, several phenotypic methods for the identification of *C. dubliniensis* and its differentiation from *C. albicans* have been reported.^{12,27–29} However, phenotypic variation in *C. dubliniensis* can result in difficulties in accurate identification.^{12,28,30} *Candida africana* was first isolated from patients in Africa and Germany.³³ In 2001, it was proposed as a new *Candida* species that produces a germ tube but no chlamydospores and is able to assimilate xylose and grow in hypertonic broths.^{7,23,34} *C. africana* has been reported as a cause of VVC in African, German, Spanish, and Italian patients.^{16,21,22,33} *C. africana* and *C. dubliniensis* strains are often misidentified as *C. albicans*. Therefore, rapid nucleic acid-based methods to identify *C. dubliniensis*, *C. albicans* and *C. africana* have been performed using polymerase chain reaction (PCR) assays.^{8,20}

In the present study we evaluated the presence of *C. albicans* and the closely related species *C. africana* and *C. dubliniensis* in clinical samples collected from patients suffering from VVC and having asymptomatic colonization by *Candida* species.

Materials and methods

Clinical *C. albicans* and non-*C. albicans* isolates

This study was carried out in the Laboratory of Microbiology of “Ramón Sardá” Maternity Hospital in Buenos Aires, Argentina, during the period of August 2012 to January 2013. Two hundred and ten immunocompetent pregnant women attending the “Ramón Sardá” maternity hospital to be examined for vaginal secretion, regardless of the presence or absence of vulvovaginitis symptoms, were enrolled. The characterization of the vaginal discharge was performed at the time of the visit when the sample was taken according to the method proposed by Odds et al.¹⁷ The specimens

were collected using sterile cotton-tipped swabs with the aid of a disposable vaginal speculum. The swabs were placed into a tube filled with saline solution prior to direct microscopic examination on a wet slide. Specific yeasts were initially phenotypically identified using conventional methods, such as the color of colonies grown on chromogenic medium CHROMagar Candida (CHROMagar Company, Paris, France) at 37 °C for 48 h, and the micromorphological studies on milk agar plates supplemented with 1% Tween 80.¹³ All the isolates that grew as light-green colonies were examined for chlamydospore formation on Staib agar as a species-specific characteristic of *C. dubliniensis*.^{12,27} Finally, all the yeasts were identified using a standard system, API ID32C (bioMérieux, Marcy l'Etoile, France).

Molecular identification of *C. albicans*, *C. dubliniensis* and *C. africana*

Definitive species identification and discrimination of all members of the *C. albicans* species complex was performed by the amplification of the *hyphal wall protein 1 (HWP1)* gene²⁰ using a single yeast colony presumptively identified as *C. albicans* or *C. dubliniensis*. The primers CR-f (5' GCT ACC ACT TCA GAA TCA TCA TC 3') and CR-r (5' GCA CCT TCA GTC GTA GAG GAC G 3') (Genbiotech, Argentina) and the PCR mixture were previously described.²⁰ The PCR reaction conditions were as follows: denaturation at 95 °C for 5 min, 30 cycles of denaturation at 94 °C for 45 s, primer annealing at 53 °C for 60 s and extension at 72 °C for 90 s; followed by a final extension at 72 °C for 10 min in a Minicycler DNA thermal cycler (TMJ Research Inc., USA). The PCR products were separated and visualized on a 1.3% (wt/vol) agarose gel. The CR-f/CR-r primers amplified a DNA segment of approximately 941 bp for *C. albicans*, 569 bp for *C. dubliniensis* and 700 bp for *C. africana*.²⁰ Positive control isolates *C. albicans* ATCC 90028 and *C. dubliniensis* from the Center of Mycology of the Faculty of Medicine, University of Buenos Aires (GenBank accession number FJ8116791.1) were tested in the same manner.

Molecular identification of *C. dubliniensis*

We also decided to identify *C. dubliniensis* using the *ACT1* gene-PCR based method. The PCR reaction was performed using specific primers from the *ACT1* gene intron of *C. dubliniensis* employing primers DUBF (5' GTA TTT GTC GTT CCC CTT TC 3') and DUBFR (5' CTG TTG TGT GCA CTA ACG TC 3') (Genbiotech, Argentina),⁸ using a single yeast colony presumptively identified as *C. albicans* or *C. dubliniensis*. The PCR reaction conditions were as follows: denaturation at 94 °C for 3 min, 30 cycles of denaturation at 94 °C for 45 s, primer annealing at 55 °C for 45 s and extension at 72 °C for 45 s, followed by a final extension at 72 °C for 10 min in a Mini-cycler DNA thermal cycler (TMJ Research Inc.). The PCR products

were separated and visualized on a 1.3% (wt/vol) agarose gel. The primers DUBF/DUBR amplified a DNA segment of approximately 288 bp. The positive control isolate of *C. dubliniensis* from the Center of Mycology of the Faculty of Medicine, University of Buenos Aires (GenBank accession number FJ8116791.1) was tested in the same manner.

Statistical methods

The statistical analysis of the data was performed using GraphPad Prism version 6.00 for Windows (GraphPad Software Inc., USA). Statistical significance was set at $p < 0.05$.

Results

Two hundred and ten immunocompetent pregnant women aged 10–42 years were evaluated. The median gestation time was about 24.8 ± 6.8 weeks. Altogether, 61.9% (130 patients) were asymptomatic while 38.1% (80 patients) were symptomatic. Vaginal discharge (100%, 80/80), vulvovaginal itching (46.2%, 37/80), vulvovaginal burning sensation (22.5%, 18/80) and dysuria (6.2%, 5/80) were among the most common described symptoms. None of them reported dyspareunia. There were statistically significant differences in vaginal discharge characteristics between symptomatic and asymptomatic patients (p -value < 0.0001). While white vaginal discharges were predominant among symptomatic patients, clear vaginal secretions were most common in the asymptomatic ones.

During the study period, 55 isolates of *Candida* or other yeasts were obtained from specimens collected from 52 patients suffering from VVC (24.8%).²

Table 1 shows the species distribution of yeasts in symptomatic (patients suffering from VVC) and asymptomatic patients (colonization). *C. albicans* was the predominant *Candida* species (42 strains; 80.7%) and was associated with three non-*C. albicans* (5.7%) isolates in three specimens. Additionally, nine isolates of *C. albicans* (50%) from asymptomatic patients ($n = 18$) were also obtained by culture. Non-*C. albicans* *Candida* species and other yeasts were more frequent in the asymptomatic group ($p = 0.0038$).

Forty-four isolates were presumptively identified as *C. albicans* on the basis of the color of the colonies on CHROMagar Candida, the micromorphological studies on milk agar plates supplemented with 1% Tween 80,¹³ and the identification codes in the API ID32C system (bioMérieux, France). Furthermore, three isolates produced typical clustered chlamydospores at the tip of the hyphae on agar milk 1% Tween-80,¹³ as well as abundant pseudohyphae and chlamydospores on Staib agar.²⁷ These results together with the carbohydrate assimilation tests using a commercially available API kit suggested that these three isolates were *C. dubliniensis*. Two of these isolates (3.8%) were implicated as the causative agent of VVC, whereas the remaining one was isolated from an asymptomatic patient. None of the isolates were identified as *C. africana*.

Table 1
Yeast isolates from symptomatic ($n = 52$) and asymptomatic ($n = 18$) patients.

Yeast species	Symptomatic patients (VVC) n (%)	Asymptomatic patients n (%)
<i>C. albicans</i>	42 (80.7)	9 (50)
<i>C. dubliniensis</i>	2 (3.8)	1 (5.5)
<i>C. glabrata</i>	2 (3.8)	1 (5.5)
<i>C. parapsilosis</i>	1 (1.9)	0 (0)
<i>C. incoplicua</i>	0 (0)	1 (5.5)
<i>C. albicans/C. glabrata</i>	2 (3.8)	0 (0)
<i>C. albicans/C. tropicalis</i>	1 (1.9)	0 (0)
<i>Rhodotorula</i> spp.	2 (3.8)	5 (27.8)
<i>Saccharomyces cerevisiae</i>	0 (0)	1 (5.5)

Molecular identification of *C. dubliniensis*

A PCR reaction was carried out among the 57 isolates showing the characteristic green colony color on CHROMagar Candida, for the molecular identification of *C. dubliniensis* using the primers DUBF and DUBFR. Three of these isolates were identified as *C. dubliniensis*.

Molecular identification of *C. albicans*, *C. africana* and *C. dubliniensis*

Among the 57 isolates presumptively identified as *C. albicans* on the basis of colony appearance on CHROMagar Candida (green colonies), the use of *HWP1* gene amplification verified that 54 of them were *C. albicans* and 3 were *C. dubliniensis*, confirming the above results. No *C. africana* isolates were found in this study. **Fig. 1** shows a 941 bp-DNA fragment as expected for *C. albicans* and a 569 bp-fragment for *C. dubliniensis*. Some *C. albicans* isolates were heterozygous and produced two DNA fragments: one of 839 bp and the other one of 941 bp. Nevertheless, we found no statistically significant differences in the rate of heterozygous and homozygous *C. albicans* isolates at the *HWP1* locus when the symptomatic (VVC) and asymptomatic groups of patients were compared ($p = 0.897$).

Discussion

Colonization refers to the presence of *Candida* in the vagina that is not associated with any sign of disease. In patients who are asymptomatic carriers of yeasts, the results of microscopic examinations of vaginal secretions tend to be negative; colonization is usually identified on the basis of a positive culture for yeasts.^{2,26} In contrast to asymptomatic colonization, VVC is defined as an infection of the vulva and the vagina, which is characterized by signs and symptoms of inflammation in the presence of *Candida* spp. and in the absence of other infectious etiology.^{2,26} In the current study, the prevalence of VVC in pregnant women was practically 25% (52/210). Garcia-Heredia et al. obtained similar results to those described,¹⁰ although higher percentages were reported.^{11,19} Recent studies have demonstrated that the prevalence of *C. albicans* accounts for 40.5–85% of cases^{14,36} and these differences may reflect the population analyzed in such studies. This study also shows that *C. albicans* was the most frequent yeast isolated from vaginal secretions in both symptomatic and asymptomatic patients. However, there is a statistical trend reflecting a high frequency of non-*C. albicans* species in asymptomatic women.

These findings suggest that vaginal colonization by yeasts is associated with non-*C. albicans* species and that the evolution of the symptoms of VVC would depend, among other factors, on the replacement of such species by *C. albicans*.¹⁵ In addition, *C. albicans* was a commensal agent in asymptomatic women (50%). In this case, the development of VVC could depend on changes in the vaginal environment.²⁶

In VVC, *C. albicans* was present in all the associations between two *Candida* species even when their frequency was low. Typically, although a single species is identified, two or more species have been found in the same vaginal culture in a very small number of women (2–5%).² As it is shown in **Fig. 1**, *C. albicans* isolates produced either one or two DNA fragments of the *HWP1* gene (941 bp or 839 bp and 941 bp, respectively). The 839 bp-DNA fragment is considered a novel allele of the *HWP1* gene. The only 941 bp-DNA fragment demonstrates that *C. albicans* isolates were homozygous at the *HWP1* locus, while the two remaining DNA fragments indicate heterozygosity.²⁴ Although no correlation was found between




Fig. 1. Molecular identification of *C. albicans* and *C. dubliniensis* by using CR-f/CR-r primers. Lanes 1–8: PCR products of *C. albicans*; lane 9: negative control; lane 10: *C. albicans* ATCC 90028; lane 11: 100 bp molecular size marker; lane 12: positive control isolate *C. dubliniensis*, Center of Mycology of the Faculty of Medicine, University of Buenos Aires (GenBank accession number FJ8116791.1).

asymptomatic and symptomatic pregnant women with respect to the presence of homozygous or heterozygous *C. albicans* isolates at the *HWP1* locus, further studies are needed to elucidate an eventual pathogenic role of these homozygous or heterozygous strains in the adherence processes.

C. dubliniensis shares so many phenotypic similarities with *C. albicans* that it may easily be misidentified as such.^{5,7} This *Candida* species appears to be most commonly associated with recurrent episodes of oral candidiasis in HIV⁵ and non-HIV infected individuals.¹² In addition, there have also been reports of recovery of *C. dubliniensis* isolates from vaginal and fecal samples. Hence, the prevalence values of VVC caused by *C. dubliniensis* are scarce, ranging from 0.17 to 2.44%.^{13,5} As previously reported, the micromorphology on Staib agar¹² was concordant with the molecular identification by PCR using specific primers from the *ACT1*-associated intron sequence of *C. dubliniensis*⁸ and the *HWP1* gene,²⁰ and was useful for selecting presumptive *C. dubliniensis* isolates. However, DNA-based methods continue to be the most effective to allow the best discrimination between the two species.³

C. africana is a yeast particularly able to cause vaginitis. Although most *C. africana* isolates recovered from clinical samples worldwide were from vaginal samples,^{16,21,22,24,32,33} Odds et al.¹⁸ described its recovery from blood in Chile (South America). Interestingly, this Chilean strain could represent the first isolation of *C. africana* from a different clinical sample, suggesting that this fungus can also be associated with a broader clinical spectrum.²² However, the results of the *HWP1* gene amplification assay revealed that no *C. africana* isolate was recovered from our vaginal specimens, which may be partly due to the fact that *C. africana* generally has a low incidence in vaginal samples^{18,24} and probably a larger number of VVC cases are necessary in order to isolate this *Candida* species in our country. Molecular biological tools are essential for the reliable identification of *C. africana* and *C. dubliniensis*, species closely related to *C. albicans*.

Conflict of interest

The authors declare no conflict of interest.

References

- Acikgoz ZC, Sancak B, Gamberzade S, Misirlioglu M. Prevalence of *Candida dubliniensis* among the stored vaginal *Candida* isolates in a Turkish hospital. *Mycoses*. 2004;47:393–6.
- Achkar JM, Fries BC. Candida infections of the genitourinary tract. *Clin Microbiol Rev*. 2010;23:253–73.
- Albaina O, Sahand IH, Brusca MI, Sullivan DJ, Fernandez de Larrinoa I, Moragues MD. Identification and characterization of nine atypical *Candida dubliniensis* clinical isolates. *J Med Microbiol*. 2015;64:147–56.
- Alcoba-Flórez J, Méndez-Alvarez S, Cano J, Guarro J, Pérez-Roth E, Arévalo MP. Phenotypic and molecular characterization of *Candida nivariensis* sp. nov., a possible new opportunistic fungus. *J Clin Microbiol*. 2005;43:4107–11.
- Coleman DC, Sullivan DJ, Bennett DE, Moran GP, Barry HJ, Shanley DB. Candidiasis: the emergence of a novel species, *Candida dubliniensis*. *AIDS*. 1997;11:557–67.
- Correia A, Sampaio P, James S, Pais C. *Candida bracarensis* sp. nov., a novel anamorphic yeast species phenotypically similar to *Candida glabrata*. *Int J Syst Evol Microbiol*. 2006;56:313–7.
- Criseo G, Scordino F, Romeo O. Current methods for identifying clinically important cryptic *Candida* species. *J Microbiol Methods*. 2015;111:50–6.
- Donnelly SA, Sullivan DJ, Shanley DB, Coleman DC. Phylogenetic analysis and rapid identification of *Candida dubliniensis* based on analysis of *ACT1* intron and exon sequences. *Microbiology*. 1999;145:1871–82.
- Farage MA, Miller KW, Sobel JD. Dynamics of the vaginal ecosystem hormonal influences. *Infect Dis Res Treat*. 2010;3:1–15.
- García Heredia M, García SD, Copolillo EF, Cora EM, Barata AD, Vay CA, de Torres RA, Tiraboschi N, Famiglietti AMR. Prevalencia de candidiasis vaginal en embarazadas. Identificación de levaduras y sensibilidad a los antifúngicos. *Rev Arg Microbiol*. 2006;38:9–12.
- Guzel AB, Ilkit M, Burgut R, Urunsak IF, Ozgunen FT. An evaluation of risk factors in pregnant women with *Candida* vaginitis and the diagnostic value of simultaneous vaginal and rectal sampling. *Mycopathologia*. 2011;172:25–36.
- Jewtuchowicz VM, Mujica MT, Brusca MI, Sordelli N, Malzone MC, Pola SJ, Iovanitti CA, Rosa A. Phenotypic and genotypic identification of *Candida dubliniensis* from subgingival sites in immunocompetent subjects in Argentina. *Oral Microbiol Immunol*. 2008;23:505–9.
- Jitaurong S, Klamsiri S, Pattarakragn N. Milk medium for germ tube and chlamydoconidia production by *Candida*. *Mycopathologia*. 1993;123:95–8.

14. Liu XP, Fan SR, Peng YT, Zhang HP. Species distribution and susceptibility of *Candida* isolates from patient with vulvovaginal candidiasis in Southern China from 2003 to 2012. *J Mycol Med.* 2014;24:106–11.
15. Lopes Consolaro ME, Albertoni TA, Yoshida CS, Mazucheli J, Peralta RM, Estivalet Svidzinski TI. Correlation of *Candida* species and symptoms among patients with vulvovaginal candidiasis in Maringá, Paraná, Brazil. *Rev Iberoam Micol.* 2004;21:202–5.
16. Nnadi NE, Ayanbimpe GM, Scordino F, Okolo MO, Enweani IB, Criseo G, Romeo O. Isolation and molecular characterization of *Candida africana* from Jos, Nigeria. *Med Mycol.* 2012;50:765–7.
17. Odds FC, Webster C, Mayurathan P, Simmons PD. *Candida* concentrations in the vagina and their association with signs and symptoms of vaginal candidosis. *J Med Vet Mycol.* 1988;26:277–83.
18. Odds FC, Bougnoux ME, Shaw DJ, Bain JM, Davidson A, Diogo D, et al. Molecular phylogenetics of *Candida albicans*. *Euk Cell.* 2007;6:1041–52.
19. Okonkwo NJ, Umeanaeto PU. Prevalence of vaginal candidiasis among pregnant women in Nnewi Town of Anambra State, Nigeria. *Afr Res Rev.* 2010;4:539–48.
20. Romeo O, Criseo G. First molecular method for discriminating between *Candida africana*, *Candida albicans*, and *Candida dubliniensis* by using *HWP1* gene. *Diagn Microbiol Infect Dis.* 2008;62:230–3.
21. Romeo O, Criseo G. Morphological, biochemical and molecular characterization of the first Italian *Candida africana* isolate. *Mycoses.* 2009;52:454–7.
22. Romeo O, Criseo G. Molecular epidemiology of *Candida albicans* and its closely related yeasts *Candida dubliniensis* and *Candida africana*. *J Clin Microbiol.* 2009;47:212–4.
23. Romeo O, Tietz HJ, Criseo G. *Candida africana*: is it a fungal pathogen. *Curr Fungal Infect Rep.* 2013;7:192–7.
24. Shan Y, Fan S, Liu X, Li J. Prevalence of *Candida albicans*-closely related yeasts, *Candida africana* and *Candida dubliniensis*, in vulvovaginal candidiasis. *Medical Mycol.* 2014;52:636–40.
25. Sobel JD. Candidal vulvovaginitis. *Clin Obst Gynecol.* 1997;36:153–212.
26. Sobel JD. Vulvovaginal candidosis. *Lancet.* 2007;369:1961–71.
27. Staib P, Morschhäuser J. Chlamydospore formation on Staib agar as a species-specific characteristic of *Candida dubliniensis*. *Mycoses.* 1999;42:521–4.
28. Sullivan DJ, Westerneng TJ, Haynes KA, Bennett DE, Coleman DC. *Candida dubliniensis* sp. nov.: phenotypic and molecular characterization of a novel species associated with oral candidosis in HIV-infected individuals. *Microbiology.* 1995;141:1507–21.
29. Sullivan DJ, Moran G, Donnelly S, Gee S, Pinjon E, McCartan B, Shanley DB, Coleman DC. *Candida dubliniensis*: an update. *Rev Iberoam Micol.* 1999;16:72–6.
30. Sullivan DJ, Moran GP, Pinjon E, Al-Mosaid A, Stokes C, Vaughan C, et al. Comparison of the epidemiology, drug resistance mechanisms, and virulence of *Candida dubliniensis* and *Candida albicans*. *FEMS Yeast Res.* 2004;4:369–76.
31. Tavanti A, Davidson AD, Gow NA, Maiden MC, Odds FC. *Candida orthopsisilosis* and *Candida metapsilosis* spp. nov. to replace *Candida parapsilosis* groups II and III. *J Clin Microbiol.* 2005;43:284–92.
32. Theill L, Dudiuk C, Morano S, Gamarra S, Nardin ME, Méndez E, et al. Prevalence and antifungal susceptibility of *Candida albicans* and its related species *Candida dubliniensis* and *Candida africana* isolated from vulvovaginal samples in a hospital of Argentina. *Rev Argent Microbiol.* 2016, <http://dx.doi.org/10.1016/j.ram.2015.10.003>
33. Tietz HJ, Küssner A, Thanos M, Pinto de Andrade M, Presber W, Schönian G. Phenotypic and genotypic characterization of unusual vaginal isolates of *Candida albicans* from Africa. *J Clin Microbiol.* 1995;33:2462–5.
34. Tietz HJ, Hopp M, Schmalreck A, Sterry W, Czaika V. *Candida africana* sp. nov., a new human pathogen or a variant of *Candida albicans*? *Mycoses.* 2001;44:437–45.
35. Üs E, Cengiz SA. Prevalence and phenotypic evaluation of *Candida dubliniensis* in pregnant women with vulvovaginal candidosis in a university hospital in Ankara. *Mycoses.* 2007;50:13–20.
36. Verghese S, Padmaja P, Asha M, Elizabeth SJ, Anitha A, Kundavi KM, et al. Prevalence, species distribution and antifungal sensitivity of vaginal yeasts in infertile women. *Ind J Pathol Microbiol.* 2001;44:313–4.