



Note

Molecular detection of *Histoplasma capsulatum* indoors: A public health approach



Juliana Amorim-Conselheiro^a, Lilian Dias Orico^a, Maria Adelaide Galvão-Dias^a, Iraci Martins Grigorio^a, Hildebrando Montenegro Neto^a, Adriana Ruckert Da Rosa^b, Débora Cardoso Oliveira^b, Carlos Pelleschi Taborda^c, Adriana Araujo Reis-Menezes^{a,*}

^a Laboratory of Zoonoses and Vector-borne Diseases, Zoonoses Surveillance Division – Health Surveillance Coordination, São Paulo, SP, Brazil

^b Chiroptera Section, Zoonoses Surveillance Division – Health Surveillance Coordination, São Paulo, SP, Brazil

^c Laboratory of Pathogenic Dimorphic Fungi, University of São Paulo – Institute of Biomedical Sciences, Department of Microbiology, São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 25 June 2018

Accepted 7 January 2019

Available online 9 March 2019

Keywords:

Histoplasmosis

Bats

Indoors

Surveillance

qPCR

ABSTRACT

Background: Histoplasmosis, caused by the dimorphic fungus *Histoplasma capsulatum*, represents an important public health problem, especially in urban environments where bats and humans cohabit indoors.

Aims: To detect the presence of *H. capsulatum* indoors, using samples of bat droppings collected in roost sites inside houses.

Methods: A Real-Time TaqMan PCR assay targeting the ITS1 region of the ribosomal DNA of *H. capsulatum* was carried out.

Results: Fifty-nine sampling points in the municipality of São Paulo were inspected, all of them located at inhabited places. *H. capsulatum* was isolated from nine samples.

Conclusions: The rapid identification and monitoring of sites where the fungus is present may contribute to make a more reliable database of *H. capsulatum* distribution.

© 2019 Asociación Española de Micología. Published by Elsevier España, S.L.U. All rights reserved.

DetECCIÓN MOLECULAR DE *HISTOPLASMA CAPSULATUM* EN INTERIORES: UN ENFOQUE DE SALUD PÚBLICA

RESUMEN

Antecedentes: La histoplasmosis, causada por el hongo dimorfo *Histoplasma capsulatum*, representa un importante problema de salud pública, especialmente en los entornos urbanos donde los murciélagos y los humanos conviven en los espacios interiores.

Objetivos: Detectar la presencia de *H. capsulatum* en interiores mediante muestras de excrementos de murciélagos recogidas en sitios de reposo de estos animales dentro de las casas.

Métodos: Se llevó a cabo un ensayo de PCR TaqMan[®] en tiempo real dirigido a la región ITS1 del ADN ribosómico de *H. capsulatum*.

Resultados: Se muestrearon 59 puntos en el municipio de São Paulo, todos ubicados en lugares habitados. *H. capsulatum* se aisló en nueve de estos lugares.

Conclusiones: Una rápida identificación y control de los sitios donde se encuentra *H. capsulatum* contribuiría a la creación de una base de datos más sólida en lo referente a la distribución de este hongo.

© 2019 Asociación Española de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Histoplasmosis

Murciélagos

Interiores

Vigilancia

qPCR

* Corresponding author.

E-mail address: adrianamenezes@prefeitura.sp.gov.br (A.A. Reis-Menezes).

<https://doi.org/10.1016/j.riam.2019.01.002>

1130-1406/© 2019 Asociación Española de Micología. Published by Elsevier España, S.L.U. All rights reserved.

Histoplasmosis, caused by the dimorphic fungus *Histoplasma capsulatum*, is an important public health problem in Brazil as information about its epidemiology is scarce.^{5,7} The fungus grows in soil enriched with birds or bats excrements, and infects humans during activities of soil disturbance.¹⁰ Bats acquire the fungus through inhalation and can develop a systemic disease, releasing spores in their feces.⁴ Therefore, the exposure to bat excrements represents an important source of infection, especially in urban environments as these animals are adapted to roosting in roofs, ceilings and other man-made shelters.^{6,9} In order to address the issue of whether or not the fungus is present inside houses, fecal samples were collected in roost sites directly into clean containers avoiding cross contamination. On ceilings with easy access, great extension and different environmental conditions, such as excess of humidity or heat, the samples were collected from different points according to these conditions. These sampling points were selected based on complaints through the System of Assistance to the Citizen of the municipality of São Paulo.

DNA extraction was performed using the PowerSoil[®] Isolation Kit (MO BIO, USA) following the manufacturer's instructions with some modifications. A Real-Time TaqMan PCR assay (TaqMan[®] Gene Expression Assay, Applied Biosystems[™], South San Francisco, CA, USA) was carried out following the manufacturer's instructions, and targeting the ITS1 region of the ribosomal DNA of *H. capsulatum*. The primers HcITS-54F (5'-ACCCTTGCTACCGGACCTGTT-3') and HcITS-204R (5'-TTTGGACTGGATTATTATCGCTCTCA-3'), and the probe HcITS-155 (5'-FAM-CGGTGAACGATTGGCGTCTGAGC-QSY 3') were used as described by Buitrago et al.² The target sequence was amplified in a final reaction of 25 µl containing 2.5 µl DEPC treated-water (Applied Biosystems[™]), 12.5 µl TaqMan[®] Environmental Master Mix 2.0 (Applied Biosystems[™]), 2.5 µl probe at 2.5 µM (Applied Biosystems[™]), 2.5 µl of each primer (0.6 µM) (Applied Biosystems[™]) and 2.5 µl of DNA. DNA extracted from the *H. capsulatum* isolate MIC 1408/14 – culture collection – Mycology Section-ZCD/SP was used as the positive control, and DEPC treated-water was the negative control. The concentrations of the probe and primers were optimized, and those with the highest normalized reporter (ΔRn) and threshold cycle (Ct) were used to prepare the standard curve generated with serial 10-fold dilutions, wherein a 97% efficiency was obtained. Horse blood was used as an exogenous control, with the β -actin gene being the external target.

Fifty-nine sampling points were inspected, all of them located at inhabited places. *H. capsulatum* was isolated from 9 samples. The first positive samples, FZ20 and FZ23, showed Ct values of 38.26 and 37.1, respectively (Fig. 1). As the Ct obtained was higher than 37 cycles, the qPCR amplification products of both samples were submitted to electrophoresis in a 1.5% agarose gel, yielding a

Table 1
Positive fecal samples and threshold cycle (Ct) values.

Fecal samples	Ct mean
FZ 20/15	36.98
FZ 23/15	36.1
FZ 07/16	36.85
FZ 02/17	37.77
FZ 09/17	38.3
FZ 10/17	37.01
FZ 14/17	36.99
FZ 9/18	38.65
FZ11/18	37.26

180-bp band, as expected. The test was repeated in sextuplicate for the same samples (Fig. 2). The qPCR products were purified using QIAquick[®] PCR purification kit (Qiagen, Hilden, Germany) following manufacturer's instructions, and sent for sequencing to the Human Genome Research Centre at the University of São Paulo. DNA analysis was performed in the equipment ABI 3730 DNA Analyser (Applied Biosystems[™]), the sequencing reactions used BigDye[™] Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems[™]) and the sequences were analyzed by Sequencing Analysis 5.3.1 software, using Base Caller KB. The sequences were compared to other sequences deposited at GenBank, showing 99% identity with *H. capsulatum*. The qPCR, electrophoresis and sequencing results obtained suggest that Ct values close to 38.5 indicate the presence of *H. capsulatum*, and that was the value used as cut-off. Table 1 shows those Ct values considered positive in the samples analyzed.

The World Organization for Animal Health (OIE) highlights the need for detecting diseases that affect the wild animal population as a tool for a better understanding of the epidemiology of these diseases and their relation to the infection in humans.⁸ Yet this strategy is regrettably not implemented by public health systems in Brazil, as there is scarce reliable information to subsidize effective measures regarding the health surveillance of histoplasmosis. Additionally, the standard method for the detection of *H. capsulatum* is its isolation in culture. Although the latter being an easy method, the fungus is slow-growing, taking several days, even months, before a result is available. Moreover, feces are a heavily contaminated material, with the presence of fast-growing bacteria that can inhibit fungal growth, and there is a necessity of a confirmatory test after isolation.¹¹ Due to these limitations, public health surveillance is impaired, as control actions regarding this disease cannot be taken in a timely manner. In this study, the use of a Real-Time TaqMan assay proved to be a very promising tool for histoplasmosis surveillance. It can also enable the improvement of control actions as infection sites are




Fig. 1. Amplification plot of the first positive qPCR results of feces samples: FZ20 Ct = 38.26; FZ 23 Ct = 37.1; positive control (C+) Ct = 12.6; exogenous internal positive control (IPC): FZ20 Ct = 28.56; FZ23 Ct = 34.64; ΔRn : normalized reporter; Ct: threshold cycle.




Fig. 2. (A) Amplification plot of a qPCR reaction from a sextuplicate of the feces sample FZ20. Three replicates showed undetermined Ct and three showed the following Ct values: Ct1 = 36.9; Ct2 = 37.1 and Ct3 = 37.72; positive control (C+) Ct = 12.6; exogenous internal positive control (IPC) Ct = 28.75. (B) Amplification plot of a qPCR from a sextuplicate of the feces sample FZ23. One replicate showed undetermined Ct and five showed the following Ct values: Ct1 = 36.62; Ct2 = 35.47; Ct3 = 36.94; Ct4 = 36.56 and Ct5 = 37.91; positive control (C+) Ct = 12.6; exogenous internal positive control (IPC) Ct = 34.98; ΔRn : normalized reporter; Ct: threshold cycle.




Fig. 3. Population density of the municipality of São Paulo and locations in which *H. capsulatum* was found (QGIS version 2.18).

detected more rapidly, suggesting a potential reduction of clinical cases of histoplasmosis. Other studies demonstrated that some PCR protocols were successful when used to find the fungus in environmental sources,^{3,4,12} but they require a gel electrophoresis analysis, making the process more laborious and time-consuming. Real-time PCR for the detection of fungal DNA has been applied in many contexts and samples, especially for diagnostic purposes, and is specific and sensitive.¹ This technique, compared to other PCR protocols, is more accurate and faster when applied to DNA extracted from clinical strains or culture extracts.^{2,11}

The sampling points where positive samples were identified corresponded to populated areas (Fig. 3). This finding is of great importance for epidemiological studies. The rapid identification and subsequent monitoring of sites within urban areas where the fungus was isolated contribute to draft a more reliable database of *H. capsulatum* distribution, as more samples can be processed in a shorter period of time, promoting a better understanding of histoplasmosis epidemiology.

Funding

This work was supported by the São Paulo Research Foundation (Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP; Grant number 2014/06571-2).

Conflict of interest

None.

Acknowledgements

The authors thank Rosane Correa de Oliveira, from Zoonoses Surveillance Division, and Dr. Ana Paula de Arruda Geraldês Kataoka, Laboratory of Zoonoses and Vector-borne Diseases, for the

institutional support and incentive to the development of the study. The authors also thank Dr. Gilda Maria B. del Negro, from Institute of Tropical Medicine – University of São Paulo-SP, and Dr. Mauro Cintra Giudice, from Oswaldo Cruz Colleges, for their technical support and expertise.

References

1. Buitrago MJ, Berenguer J, Mellado E, Tudela-Rodríguez JL, Cuenca-Estrella M. Detection of imported histoplasmosis in serum of HIV-infected patients using a real-time PCR-based assay. *Eur J Clin Microbiol Infect Dis.* 2006;25:665–8.
2. Buitrago MJ, Canteros CE, Frías de León G, Gonzales A, Marques-Evangelista MO, Muñoz CO, et al. Comparison of PCR protocols for detecting *Histoplasma capsulatum* DNA through a multicenter study. *Rev Iberoam Micol.* 2013;30:256–60.
3. De León MGF, López GA, Taylor ML, Altamirano GA, Reyes-Montes MR. Development of specific sequence-characterizes amplified region markers for detecting *Histoplasma capsulatum* in clinical and environmental samples. *J Clin Microbiol.* 2012;50:673–9.
4. Dos Santos B, Langoni H, Silva RC, Menozzi BD, Bosco SMG, Paiz LM, et al. Molecular detection of *Histoplasma capsulatum* in insectivorous and frugivorous bats in Southeastern Brazil. *Med Mycol.* 2017:1–4.
5. Ferreira BS, Araújo Filho JA, Pereira NM, Godoy LM, Lamounier BB, Nunes ED, et al. Disseminated histoplasmosis in AIDS patients: an urban disease. Experience in a metropolis in the middle east of Brazil. *Le Infez Med.* 2017;3:258–62.
6. Galvão Dias MA, Zancopé Oliveira RM, Giudice MC, Montenegro Netto H, Jordão LR, Grigorio IM, et al. Isolation of *Histoplasma capsulatum* from bats in the urban area of São Paulo State, Brazil. *Epidemiol Infect.* 2010:1–3.
7. Gómez BL. Histoplasmosis: epidemiology in Latin America. *Curr Fungal Infect Rep.* 2011;5:199–205.
8. Jebara KB, Ripani A, Popovic M, Forcella S. WAHIS-Wild and its interference: the OIE worldwide monitoring system for wild animal diseases. *Veter Ital.* 2016;52:91–100.
9. Jung K, Kalko EKV. Adaptability and vulnerability of high flying Neotropical aerial insectivorous bats to urbanization. *Divers Distrib.* 2011:1–13.
10. Kauffman CA. Histoplasmosis: a clinical and laboratory update. *Clin Microbiol Rev.* 2007;20:115–32.
11. Martagon-Villamil J, Shrestha N, Sholtis M, Isada CM, Hall GS, Bryne T, et al. Identification of *Histoplasma capsulatum* from culture extracts by Real-Time PCR. *J Clin Microbiol.* 2003;41:1295–8.
12. Norkaew T, Ohno H, Sriburee P, Tanabe K, Tharavichitkul P, Takarn P, et al. Detection of environmental sources of *Histoplasma capsulatum* in Chiang Mai, Thailand, by nested PCR. *Mycopathologia.* 2013;176:395–402.