

Pacifier and swaddling effective in impeding premature infant's pain score and heart rate

Defi Efendi*, Yeni Rustina and Dewi Gayatri

Faculty of Nursing, Universitas Indonesia, Depok, Jawa Barat, Indonesia

KEYWORDS

Premature;
Pacifier;
Swaddling;
Invasive procedure;
Pain

Abstract

Objective: To assess the effectiveness of pacifier and swaddling on premature infant's pain score, hearthrate, and oxygen saturation during an invasive procedure.

Method: This randomized control trial involv 30 premature infants who were randomly assigned into control (n = 15) and intervention (n = 15) groups using parallel design. Infants in the intervention group received pacifier and swaddling when they were undergoing invasive procedures. The outcome indicators of the two-day intervention were pain score, hearth rate, and oxygen saturation. The Premature Infant Pain Profile (PIPP) was used in this study to measure infants' pain.

Results: The paired t-test results showed that the pain score and heart rate were significantly increased following the procedure in the control group ($p = 0.003$; $p = 0.013 < 0.05$); meanwhile, there was no significant increase in the intervention group ($p = 0.256$; $p = 0.783 > 0.005$). There was no significant different in oxygen saturation in the control group ($p = 0.270$) and in the intervention ($p = 0.370$) group before and after the procedure.

Conclusions: Providing pacifier and swaddling can impede the increase of premature infants' pain score and hearth rate during an invasive procedures, therefore it can be implemented as an alternative to pain management in premature infants.

© 2018 Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Premature infants receive invasive procedures for 93 times during hospitalization^{1,2}. These invasive procedures can cause stress to the infants and lead to frequent stress response^{3,4}. The Synactive Theory of Development proposed by Als in 1982 was relevant to this phenomenon. The theory consists of five subsystems: autonomic stability (physiological parameters), motor control (body movement); state control (sleep cycle), interaction and social responsiveness,

and self-regulation⁵. During the invasive procedure, there will be over stimulation that influences all subsystems in infants. Those influences include instability in heart rate, oxygen saturation, and respiration rate. The changes in the motor subsystem are manifested by hand and leg agitation activities; while the changes in state subsystem can be manifested by infant sleep disturbances⁶⁻⁸. Those changes can influence the process of brain nerve myelinization that can cause development disturbances in the short and long term periods⁹.

*Corresponding author.

Email: defiefendi86@gmail.com (D. Efendi).

The non pharmacologic pain management has been developed years ago. Previous studies had been analyzed the effectiveness of swaddling in pain controlling¹⁰, physiological function¹¹, safety¹², and sleep¹³. Meanwhile, the use of pacifier had been proved to prevent heart variability¹⁴, and as a means of pain control^{15,16}. A numbers of previous studies have used combination between pacifier and sucrose¹⁷⁻²¹, and facilitated tucking^{6,8} in dealing with premature infants' pain who undergoing several invasive procedures such as heel-stick procedure, intramuscular injection of Hepatitis Vaccine, or intra venous insertion. However, there are limited studies in using combination between pacifier and swaddling in different pain procedures in preterm infants.

Method

This study was a prospective randomized controlled trial with parallel design approach. Each sample received one type of invasive procedure such as hill-prick or venous access which was randomly allocated to the infants receiving pacifier and swaddling (Group A) or routine care (Group B). The blinding process was carried out by two experts to evaluate pain scores²².

The study sample was selected based on the convenience sampling from the special care unit and Neonatal Intensive Care Unit at Dr. Cipto Mangunkusumo Hospital, Jakarta. There were 30 preterm infants who met the criterias: (1) gestational age 26-36 weeks; (2) 2-30 days of post birth age, and (3) birth weight < 2500 g. The exclusion criterias-were: (1) infants on sedative treatment, muscle relaxant, antiepileptic or analgesic drugs; (2) have congenital anomalies; (3) have neurological problems and history of neonatal seizure, and (4) experience severe physiological alteration.

Pain scores and physiological functions were measured 15 minutes before invasive procedure (baseline/T1). The intervention (pacifier and swaddling) was applied three minutes before the invasive procedure started (T2). The next measurements were carried out at the beginning of invasive procedure (T3), three minutes (T4) and 15 minutes after the procedure ending (T5). These measurements conducted in two days consequently in both groups.

Infants' pain was measured using the Premature Infant Pain Profile (PIPP) which was developed by Steven, Jhonston, Pethrysen, and Taddion²³. The PIPP is reported to have a moderate internal consistency (0.59-0.76 item total correlation), high interrater reliability (0.95-0.97) and intrarater reliability (0.89-0.91)^{7,24}.

The Ethical clearance was obtained from the Ethical Research Committee, Faculty of Nursing Universitas Indonesia (0244/UN2.F12.D/HKP.02.04/2015) and the Ethical Research Committee, Faculty of Medicine Universitas Indonesia (0244/UN2.F12.D/HKP.02.04/2015). Written informed consent was obtained from mothers.

Pain score and physiological responses were measured five times (T1, T2, T3, T4, and T5). Data were rated by two experts. A Bland Altman test was carried out to test the interrater reliability of two experts and the result were 0.632 and 1.99 in the range of -5 to +5. It means that there was a similar perception between experts.

Chi-square test was used to test the homogeneity of data in both control and intervention groups. The hypothesis was

tested using dependent and independent t-test with the level of significance was 0.05.

Results

The result of the homogeneity test showed that there were no significant differences in infants' characteristics between control and intervention groups. The characteristics were gestational age, chronological age, the length of stay, birth weight, body weight, length of the invasive procedure, gender, type of invasive procedure, and Apgar score.

The output of this study was the comparison of pain score, saturation, heart rate differences before and after invasive procedures between intervention and control groups (Table 1). There were no significant differences in all variables. However, there were significant differences in several spots of measurements between two groups (Figure 1).

The peak score of pain was during the invasive procedure. The score increased 6.67 point in the control group compared to 2.34 point in the intervention group. The increase of pain score was significant in the control group ($p = 0.003$) (Figure 2).

The heart rate increased in both groups during the procedure, and it decreased significantly in the intervention group after the procedure ($p = 0.013$).

The oxygen saturation was decreased in both groups and relieved three minutes after the procedure. There was no significant difference in oxygen saturations before, during, and after procedures between two groups.

Discussion

The low score of pain and stability of physiological function are the main objectives of non-pharmacologic pain management. The main objectives of this study were to measure the differences in pain score, saturation, and heart rate before, during, and after procedures between infants who received pacifier and swaddling, and the control group who received routine care. Based on those indicators, there were no significant effect of pacifier and swaddling in reducing pain score, stabilizing heart rate, and oxygen saturation. This result was relevant with previous studies that providing pacifier were not significantly reducing the pain score in preterm infants²⁵ and preventing alteration in heart rate¹⁴. Other studies also found that using pacifier and swaddling was not effective in preventing oxygen saturation variability²⁶. Longer stimulation is needed to give effect on the oxygen saturation changes²⁶.

On the other hand, several studies had different finding with this study. The previous study identified that pacifier with sucrose can decrease infant's pain¹⁷. A meta-analysis of 10 articles identified that providing pacifier can decrease pain responses¹⁶ and decrease pain score in term and preterm infants^{8,20}. Review articles also found that pacifier can decrease infants' pain score^{27,28}.

This study was different with previous studies in term of infant gestational age. In the previous studies, gestational age of the infants involved was more than 37 weeks^{17,20}; meanwhile, in this study the gestational age of the infant was less than 37 weeks. Preterm infants are more sensitive

Table 1 Comparison of treatment effects on pain score, hearth rate and oxygen saturation (n = 30)

Measurement indicator	Group	N (%)	Mean ± SD	Within group	Between group
				t (p) ^a	t (p) ^b
Pain score					
	<i>Intervention</i>	15 (50)			
	Before		5.93 ± 2.22	2.81 (0.783)	
	After		6.13 ± 2.00		
	Difference		-0.33 ± 2.90		
	<i>Control</i>	15 (50)			
	Before		5.40 ± 1.84	3.56 (0.003) ^c	
	After		7.67 ± 2.74		
	Difference		-2.33 ± 2.58		2.00 (0.056)
Hearth rate					
	<i>Intervention</i>	15 (50)			
	Before		139.40 ± 13.69	1.18 (0.256)	
	After		144.33 ± 20.19		
	Difference		-4.93 ± 16.15		
	<i>Control</i>	15 (50)			
	Before		149.63 ± 11.45	2.84 (0.013) ^c	
	After		158.57 ± 18.90		
	Difference		-8.93 ± 12.20		0.77 (0.450)
Oxygen saturation					
	<i>Intervention</i>	15 (50)			
	Before		96.80 ± 3.05	1.16 (0.267)	
	After		96.13 ± 2.85		
	Difference		0.8 ± 2.21		
	<i>Control</i>	15 (50)			
	Before		96.33 ± 4.62	0.92 (0.370)	
	After		97.13 ± 4.66		
	Difference		0.8 ± 3.35		1.48 (0.150)

Difference: the result from mean after invasive procedure minus mean before invasive procedure.
 Between groups: the comparison of mean differences between intervention and control groups.
^aCalculated by dependent t-test.
^bCalculated by independent t-test.
^cSignificant at $\alpha < 0.05$.




Figure 1 Pain score of the preterm infant (n = 30).




Figure 2 Heart rate of the preterm infant (n = 30).




Figure 3 The mean of oxygen saturation of the preterm infant (n = 30).

to stimuli which characterized by physiological changes, ineffective regulation, and negative cues³⁰⁻³². Prematurity can lead the increase of hearth rate, respiration rate, and oxygen saturation compared to term infants. Besides that, our study did not measure baseline data of infant state which was done in the previous study that may influence the difference of physiological function indicators.

The pain observation in this study was two days, meanwhile the observation in the previous studies was carried out in three and five days^{8,20}. Furthermore, the previous studies measure the mean score of the whole observation period^{17,20}, while this study using different score between before and after the procedure. This study is also different with other previous study in regard to the intervention used. This study used a combination between pacifier and swaddling, yet other studies used a combination of pacifier-facilitated tucking⁸, also pacifier and sucrose^{17,20}.

However, our study found that pacifier and swaddling impeding the increase of pain score during acute period (during procedure-three minutes after the procedure). The frequency of heart rate also decreased faster in the pacifier and swaddling group compared to infants in the routine care group. The decrease of heart rate was faster significantly in the pacifier and swaddling group ($p = 0.013$) (Figure 3).

Physiologically, heart rate, respiration rate, blood pressure, intracranial pressure of infants who receive pain procedure will increase. This response due to the release of adrenal stress hormones associated with pain⁷. The pain was impeded by the antinociceptive effect of application pacifier and swaddling that can reduce pain sensation¹⁷.

Based on Gate Control Theory, applying pacifier and swaddling can impede the transfer of pain impulse along the spinothalamic tract that can reduce the pain sensation^{28,33}. The combination of pacifier and swaddling can increase the stimuli that sift transferring pain stimuli which can generate stronger analgesic effect^{28,34}. Pacifier and swaddling are simple tools that are easy to find in the developing country.

Conflicts of interest

The authors declare that they have no conflicts interest.

References

- Newnham CA, Inder TE, Milgrom J. Early human development measuring preterm cumulative stressors within the NICU: The neonatal infant stressor scale. *Early Hum Dev* [Internet]. 2009;85(9):549-55. doi: <http://dx.doi.org/10.1016/j.earlhumdev.2009.05.002>.
- Hunt KN. The NICU: Environmental effects of the neonatal intensive care unit on infants and caregivers. In Unpublished; 2011.
- Kenner C, McGrath jacqueline. *Developmental care of newborn & preterm infants a guide for health professional*. St. Louis Missouri: Mosby, Inc.; 2004.
- Kenner C, Lott JW. *Comprehensive neonatal nursing care*. 5th ed. Zuccarini M, editor. New York: Springer Publisher Company; 2014. 78 p.
- Als H. Individualized developmental care for preterm infants. *Encyclopedia on Early Childhood Development*. 2004. p. 1-7.
- Liaw JJ, Yang L, Lee CM, Fan H-C, Chang Y-C, Cheng L-P. Effects of combined use of non-nutritive sucking, oral sucrose, and facilitated tucking on infant behavioural states across heel-stick procedures: A prospective, randomised controlled trial. *Int J Nurs Stud* [Internet]. 2013 Jul [cited 2015 Jan 31];50(7):883-94. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23068310>
- Mitchell A, Brooks S, Roane D. The premature infant and painful procedures. *Pain Manag Nurs*. 2000;1(2):58-65.
- Yin J jjuan L, Ying-ti SC, Yang L, Katherine Wang K-W, Chen C-M, Chang Y-C, et al. Non-nutritive sucking and facilitated tucking relieve preterm infant pain during heel-stick procedures: A prospective, randomised controlled crossover trial. *Int J Nurs Stud* [Internet]. 2012 Mar [cited 2015 Feb 17];49(3):300-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22001561>
- Westrup B, Sizun J, Lagercrantz H. Family-centered developmental supportive care: A holistic and humane approach to reduce stress and pain in neonates. *J Perinatol* [Internet]. 2007 May [cited 2014 May 5];27:512-8. Available at: <http://www.nature.com/doi/10.1038/sj.jp.7211724>
- Ho S, Ho L. Effects of facilitated swaddling for controlling procedural pain in premature neonates: A randomized controlled trial. *J Pain*. 2012;13(4):59.
- Gerard CM, Harris KA, Thach BT. Physiologic studies on swaddling: An ancient child care practice. *J Pediatr*. 2002;141(2):399-404.
- Mcdonnell E, Moon RY. Infant deaths and injuries associated with wearable blankets, swaddle wraps, and swaddling. *J Pediatr* [Internet]. 2014;164(5):1152-6. doi: <http://dx.doi.org/10.1016/j.jpeds.2013.12.045>.
- Richardson HL, Walker AM, Horne RSC. Influence of swaddling experience on spontaneous arousal patterns and autonomic control in sleeping infants. *J Pediatr* [Internet]. 2010;157(1):85-91. doi: <http://dx.doi.org/10.1016/j.jpeds.2010.01.005>.
- Lappi H, Valkonen-korhonen M, Georgiadis S, Tarvainen MP, Tarkka IM, Karjalainen PA, et al. Effects of nutritive and non-nutritive sucking on infant heart rate variability during the first 6 months of life. *Infant Behav Dev*. 2007;30:546-56.
- Phillips RM, Chantry CJ, Gallagher MP. Analgesic effects of breast-feeding or pacifier use with maternal holding in term infants. *Ambul Pediatr*. 2005;5(6):359-64.
- Edwards L. Non-nutritive sucking and the preterm infant. *UMI Diss Publ*. 2011;1-7.
- Akman I, Eren O. Sweet solutions and pacifiers for pain relief in newborn infants. *J Pain*. 2002;3(3):199-202.
- Carbajal R, Lenclen R, Gajdos V, Jugie M, Paupe A, Objective A. Preterm neonates during subcutaneous injections. *Pediatrics*. 2002;110(2):389-94.
- Liaw J, Yang L, Ti Y, Blackburn ST, Chang Y, Sun L. Non-nutritive sucking relieves pain for preterm infants during heel stick procedures in Taiwan. *J Clin Nurs*. 2010;19:2741-51.

20. Liaw J, Zeng W, Yang L. Nonnutritive sucking and oral sucrose relieve neonatal pain during intramuscular injection of hepatitis vaccine. *J Pain Symptom Manage* [Internet]. 2011;42(6):918-30. doi: <http://dx.doi.org/10.1016/j.jpainsymman.2011.02.016>.
21. Miller HD. Nonnutritive suckling and sucrose-induced analgesia: Effect on heart rate, oxygen saturation, and pain in intubated infants. *UMI Diss Publ.* 2009;
22. Sastroasmoro S, Ismael S. *Dasar-dasar metodologi penelitian klinis*. Jakarta: sagung seto; 2011.
23. Steven B, Jhonston C, Pethrysen P, Taddion A. Premature infant pain profile: Development and initial validation. *Clin J Pain.* 1996;12(1):13-22.
24. Larson CW. The efficacy of facilitated tucking for relieving procedural pain of endotracheal suctioning in very low birth weight infant. *UMI Diss Publ.* 2002;1-109.
25. Carbajal R, Lenclen R, Gajdos V, Jugie M, Paupe A, Objective A. Crossover trial of analgesic efficacy of glucose and pacifier in very preterm neonates during subcutaneous injections. *Pediatrics.* 2002;110(2):389-94.
26. Hashemi F, Taheri L, Ghodsbin F, Vossoughi M. Comparing the effect of swaddling and breastfeeding and their combined effect on the pain induced by BCG vaccination in infants. *Appl Nurs Res* [Internet]. 2015;1-17. doi: <http://dx.doi.org/10.1016/j.apnr.2015.05.013>.
27. Hall RW. Pain management in newborns analgesia sedation pain stress NICU infant-newborn. *Clin Perinatol* [Internet]. 2014;41(4):895-924. Available at: <http://dx.doi.org/10.1016/j.clp.2014.08.010>
28. Johnston CC, Fernandes AM, Campbell-yeo M. Pain in neonates is different. *Pain* [Internet]. 2011;152(3):S65-73. Available at: <http://dx.doi.org/10.1016/j.pain.2010.10.008>
29. Sexton S, Natale R. Risk and benefit of pacifier. *Am Fam Physician.* 2009;79(8):682-7.
30. Pineda RG, Otr L, Tjoeng TH, Vavasseur C, Kidokoro H, Neil JJ, et al. Patterns of altered neurobehavior in preterm infants within the Neonatal Intensive Care Unit. *J Pediatr* [Internet]. 2013;162(3):470-6.e1. doi: <http://dx.doi.org/10.1016/j.jpeds.2012.08.011>.
31. Ranganna R, Bustani P. Reducing noise on the neonatal unit. *Infant.* 2011;7(1):25-8.
32. Boxwell G. *Neonatal intensive care nursing*. 2nd ed. New York: Routledge; 2010.
33. Melzack R, Wall PD. Pain mechanisms: A new theory. *J Storl.* 1965;150(3699):971-9.
34. Ramachandran S, Dutta S. Early developmental care interventions of preterm very low birth weight infants. *Indian Pediatr.* 2013;50:765-71.