



## Letters to the editor

**Lactate-dehydrogenase associated with mortality in hospitalized patients with COVID-19 in Mexico**

Dear Editor,

Recently, an attractive study entitled “Lactate-dehydrogenase associated with mortality in hospitalized patients with COVID-19 in Mexico: a multi-centre retrospective cohort study” was published in *Annals of Hepatology* [1]. In this study, the authors investigated the risk factors associated with mortality in hospitalized patients with coronavirus disease 2019 (COVID-19). Undoubtedly, this is a study with essential clinical guiding value. However, the following concerns need to be noted.

First of all, it is apparent that the purpose of this study is to explore the risk factors related to mortality in hospitalized patients with COVID-19 and several potential variables were screened in this study. However, some variables that are easily obtained in clinical practice and have been clarified to be related to mortality in COVID-19 patients should not be neglected. For instance, previous studies have exhibited that obesity is closely related to the prognosis and disease severity in COVID-19 patients [2,3], and it is recommended that clinicians should pay enough attention to obese patients in order to improve the short- and long-term prognosis. On the other hand, considering that weight and height are currently used to define obesity [4], and these two variables can be easily extracted from medical records. Therefore, the influence of obesity

on the prognosis of COVID-19 patients should not be unnoticed. It is recommended that obesity should be adjusted as a potential confounding variable in the multivariate analysis to ensure the reliability of the final conclusions.

Second, it is common knowledge that the severity of the disease is associated with the prognosis. The higher the severity of the disease, the worse the prognosis. Similarly, this regular pattern also applies to COVID patients. Nowadays, the severity of COVID could be divided into the following four types: mild, moderate, severe, and critical [5]. Additionally, prior study has demonstrated that the severity of disease in COVID patients is associated with a higher risk of mortality [6]. Unfortunately, this study only described the recruitment of COVID patients, yet does not clarify the severity of the patients' disease, which could easily lead to misinterpretations. One possible hypothesis is that deaths mainly occur in severe/critical patients, and are caused by the severity of the disease, not by lactate dehydrogenase level.

Third, it is undeniable that treatment strategies are related to the prognosis of COVID patients, and appropriate treatment strategies can remarkably improve the prognosis of patients [7]. Nevertheless, this study did not describe in detail the treatment drugs and strategies between the two groups. How many patients need to be treated in the intensive care unit (ICU)? How many patients have received mechanical ventilation treatment? Obviously, Table 1 describes that 3.4% of patients in the survivor group have a history of dialysis, while 10.1% of the patients in the non-survivor group have a history of dialysis ( $P=0.012$ ). Nonetheless, it is indistinct that how many

**Table 1**  
Demographic characteristics and laboratory findings on admission.

Variable	Survivors (n = 298)	Non-survivors (n = 79)	P value
Age, mean (range), years	41.8 ( $\pm 22.7$ )	46.72 ( $\pm 25.7$ )	0.016
Male, n (%)	215 (72.1%)	62 (78.5%)	0.257
Diabetes mellitus, n (%)	64 (21.5%)	24 (30.4%)	0.096
Hypertension, n (%)	56 (18.8%)	20 (25.3%)	0.199
Cardiovascular disease, n (%)	6 (2%)	9 (11.4%)	0.000
COPD, n (%)	8 (2.7%)	8 (10.1%)	0.013
Dialysis, n (%)	10 (3.4%)	8 (10.1%)	0.012
Haemoglobin g/dl, median (IQR)	14.8 (13.3–15.8)	14 (12.2–15.5)	0.022
Leukocytes $\times 10^3/\mu\text{l}$ , median (IQR)	7.3 (5.4–10.8)	9.4 (8.2–13.1)	0.000
Platelets $\times 10^3/\mu\text{l}$ , median (IQR)	219 (171.7–301.5)	243 (180–310)	0.454
Creatinine mg/dl, median (IQR)	0.86 (0.7–1)	0.95 (0.7–1.7)	0.005
Total bilirubin mg/dl, median (IQR)	0.77 (0.51–1)	0.97 (0.6–2.2)	0.000
Alkaline phosphatase U/l, median (IQR)	76 (58–102.5)	102 (68–124)	0.001
ALT U/l, median (IQR)	36 (26–56)	46 (24–100)	0.040
AST U/l, median (IQR)	43.5 (31–61.5)	51 (36–142)	0.000
CRP mg/dl, median (IQR)	121 (48–197)	232 (126.5–321.5)	0.000
INR, median (IQR)	1 (1–1.2)	1 (1–1.2)	0.888
Albumin g/dl, median (IQR)	3.5 (3–3.9)	3.3 (3–3.7)	0.283
Sodium mmol/l, median (IQR)	138 (136–140)	137 (135–140)	0.421
LDH U/l, median (IQR)	380.5 (265.7–523.2)	536 (419–776)	0.000
MELD score, median (IQR)	8 (6–10)	9 (8–16)	0.000

COPD (Chronic Obstructive Pulmonary Disease), ALT (alanine aminotransferase), AST (aspartate aminotransferase), CRP (C reactive protein), INR (international normalized ratio), LDH (lactate dehydrogenase), MELD (Model of End-Stage Liver Disease).

patients receive continuous renal replacement therapy (CRRT) in ICU for patients with a history of dialysis?

### Declaration of funding interests

None.

### Conflict of interest

The authors have no conflicts of interest to declare.

### References

- [1] Vidal-Cevallos P, Higuera-De-La-Tijera F, Chávez-Tapia NC, Sanchez-Giron F, Reyes EC, Rosales-Salyano VH, et al. Lactate-dehydrogenase associated with mortality in hospitalized patients with COVID-19 in Mexico: a multi-centre retrospective cohort study. *Ann Hepatol* 2021;100338, <http://dx.doi.org/10.1016/j.aohep.2021.100338>.
- [2] Sattar N, McInnes IB, McMurray JJV. Obesity is a risk factor for severe COVID-19 infection: multiple potential mechanisms. *Circulation* 2020;142(1):4–6.
- [3] Cai Q, Chen F, Wang T, Luo F, Liu X, Wu Q, et al. Obesity and COVID-19 severity in a designated hospital in Shenzhen, China. *Diabetes Care* 2020;43(7):1392–8.
- [4] Compston JE, Flahive J, Hosmer DW, Watts NB, Siris ES, Silverman S, et al. Relationship of weight, height, and body mass index with fracture risk at different sites in postmenopausal women: the Global Longitudinal study of Osteoporosis in Women (GLOW). *J Bone Miner Res* 2014;29(2):487–93.
- [5] Zhang J, Wang X, Jia X, Li J, Hu K, Chen G, et al. Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect* 2020;26(6):767–72.
- [6] Velavan TP, Meyer CG. Mild versus severe COVID-19: laboratory markers. *Int J Infect Dis* 2020;95:304–7.
- [7] Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the treatment of Covid-19. *N Engl J Med* 2020;383(19):1813–26.

Feiyan Zhu<sup>a</sup>

Xiaofei Li<sup>b,\*</sup>

<sup>a</sup> Laboratory Department, YiWu Central Hospital, Zhejiang 322000, China

<sup>b</sup> Department of Infectious Diseases, YiWu Central Hospital, Zhejiang 322000, China

\* Corresponding author.

E-mail address: [icqwc46@163.com](mailto:icqwc46@163.com) (X. Li)

3 March 2021

Available online 14 April 2021