Contents lists available at ScienceDirect





Annals of Hepatology

journal homepage: www.elsevier.es/annalsofhepatology

Liquid biopsy: New opportunities for precision medicine in hepatocellular carcinoma care



Inah Marie C. Aquino^{a,b}, Devis Pascut^{b,*}

^a College of Medicine, University of the Philippines Manila, Ermita, Manila, Metro Manila 1000, Philippines ^b Liver Cancer Unit, Fondazione Italiana Fegato – ONLUS, Basovizza, Trieste 34149, Italy

ARTICLE INFO

Article History: Received 25 August 2023 Accepted 18 October 2023 Available online 14 November 2023

Keywords: Liquid biopsy ctDNA cfDNA Artificial intelligence HCC CDSS

ABSTRACT

Liquid biopsy, specifically the analysis of circulating tumor DNA (ctDNA), offers a non-invasive approach for hepatocellular carcinoma (HCC) diagnosis and management. However, its implementation in the clinical setting is difficult due to challenges such as low ctDNA yield and difficulty in understanding the mutation signals from background noise. This review highlights the crucial role of artificial intelligence (AI) in addressing these limitations and in improving discoveries in the field of liquid biopsy for HCC care. Combining AI with liquid biopsy data can offer a promising future for the discovery of novel biomarkers and an AI-powered clinical decision support system (CDSS) can turn liquid biopsy into an important tool for personalized management of HCC. Despite the current challenges, the integration of AI shows promise to significantly improve patient outcomes and revolutionize the field of oncology.

© 2023 Fundación Clínica Médica Sur, A.C. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)

1. Introduction

Liver cancer is the sixth most common cancer, and it is now the third leading cause of cancer-related death worldwide. In 2020, more than 900,000 new cases were diagnosed globally, and more than 800,000 liver cancer-related deaths were reported [1]. Hepatocellular carcinoma (HCC) represents about 75 %-85 % of primary liver cancers being a major health burden worldwide [2]. In Europe, the age-stan-dardized mortality rate is around 3.8–4.0 per 100,000 population [2]. While in the US, the survival rates for HCC patients remain low with only a 3–34 % 5-year survival rate [3].

HCC can arise from various causes of chronic liver injury but regardless of etiology, it is suggested that a common mechanism leading to it is the continuous cycle of cell death and regeneration [4]. This can result in accumulation of multiple genetic events such as somatic mutations, copy number alterations, pathway alterations, and epigenetic changes that can lead to HCC initiation, progression, and metastasis [4]. Advancements in sequencing techniques allowed molecular characterization of tumors which can provide an understanding of the genomic landscape of HCC and help identify driver

Corresponding author.

which there is a shift from a "one-size fits all" approach to an individualized management approach however, challenges are met in the collection and analysis of tumor tissue [6,7]. Archived tissues can be used to analyze mutations, but it may not contain enough tumor content to satisfy test sensitivity and may be degraded to provide any pertinent information [5]. It should also be highlighted that in the past, tissue biopsy for liver tumors was discouraged due to the socalled "seeding" problem. In addition, liver cancer exhibits significant intratumoral heterogeneity, making tissue biopsies less representative. In this regard, liquid biopsy offers great potential to overcome these limitations since it consists of a minimally-invasive analysis of biological fluids, most commonly blood, that can be used to detect cancer at early stages, thus giving patients good chances for effective treatment [8].

mutations [4-6]. This supports the rise of precision medicine in

In recent years, a wide range of liquid-biopsy tests has been developed primarily using circulating tumor proteins, nucleic acids, such as circulating tumor DNA (ctDNA), circulating tumor cells (CTCs), and tumor-derived extracellular vesicles (EVs). The use of liquid biopsy, particularly ctDNA analysis, has emerged as a promising approach in the field of liver cancer diagnostics and monitoring [8]. However, challenges such as low DNA yield [9,10] and the difficulty in deciphering mutation signals against background noise [11] have limited its widespread clinical implementation. In this minireview, we emphasize the crucial role of artificial intelligence (AI) in addressing these limitations and accelerating discoveries in the field of liquid biopsy for liver cancer.

https://doi.org/10.1016/j.aohep.2023.101176

1665-2681/© 2023 Fundación Clínica Médica Sur, A.C. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)

Abbreviations: HCC, hepatocellular carcinoma; ctDNA, circulating tumor DNA; cfDNA, cell-free DNA; miRNA, microRNA; CTCs, circulating tumor cells; MAFs, mutant allele frequencies; AI, artificial intelligence; CDSS, clinical decision support system; AUC, area under the curve; OS, overall survival; PFS, progression-free survival; RFS, recurrence-free survival; HR, hazard ratio

E-mail address: devis.pascut@fegato.it (D. Pascut).

2. Liquid biopsy

Interest in liquid biopsy has continued to grow over the years and has led to approval for blood-based tests for precision cancer care [12]. In 2013, the US Food and Drug Administration (FDA) approved the CellSearchTM CTC enumeration platform, which enumerates epithelial circulating tumor cells (CTCs) (Table 1) [12]. This groundbreaking approval paved the way for the implementation of liquid biopsy in clinical trials such as the METABREAST trial, where CTC counts were used as a criterion for selecting first-line treatment in meta-static breast cancer (Table 1) [8,13].

Subsequently, in 2016, the first ctDNA-based companion diagnostic test, known as the Cobas EGFR Mutation Test v2, was developed to detect epidermal growth factor receptor (EGFR)-sensitizing mutations in non-small cell lung cancer (NSCLC) patients (Table 1). This test played a crucial role in guiding the use of EGFR-tyrosine kinase inhibitors as a targeted therapy [14]. With the rise of next-generation sequencing techniques, multigene panels were developed to detect target mutations in several cancer-related genes in advanced cancer. In 2019, FDA approved some liquid biopsy tests. One of these is the Guardant360 CDx (Table 1), which analyzes a 73-gene panel of cell-free DNA (cfDNA) to guide treatment for patients with NSCLC. Another is CancerSEEK, which identifies eight prevalent cancer types through the analysis of eight protein biomarkers combined with tumor-specific mutations in cfDNA of cancer patients [12]. Additional FDA-approved liquid biopsy tests are outlined in Table 1. These examples highlight how these tests have improved the selection of patients who could derive greater benefit from targeted therapies, signifying a noteworthy advancement in personalized medicine within the field of oncology.

HCC in particular presents considerable challenges in its management and treatment due to pronounced tumor heterogeneity, both in terms of intra- and inter-tumor molecular diversity [15]. Multinodular HCCs, diagnosed in approximately 41 % to 75 % of patients, limit the availability of curative treatment options and often lead to unfavorable prognosis [16]. The extensive tumor heterogeneity, particularly in multicentric HCC cases, could be underestimated when assessed through a single tumor biopsy, posing significant obstacles to precision medicine and the development of biomarker-based solutions in HCC [16,17]. Advancements in liquid biopsy, particularly the analysis of ctDNA, have revolutionized oncology by enhancing early detection and improving monitoring and targeted treatments through the evaluation of genomic and molecular profiles of tumors [17]. In contrast to various other cancers (as shown in Table 1), the field of liquid biopsy for HCC has been characterized by a lack of extensive and comprehensive studies able to provide sufficient and consistent results. This dearth of robust data has contributed to challenges in attaining FDA approvals for HCC-related applications.

3. Circulating tumor DNA in HCC

cfDNA, first reported in human peripheral blood in 1948 by Mandel and Metais, exists as double-stranded fragments of approximately 150 to 200 base pairs in length. In healthy individuals, cfDNA derived from apoptotic myeloid and lymphoid cells is present at low levels (about 10 to 15 ng per milliliter) [18]. However, in the presence of tumors, inflammation, tissue damage, or after surgery, the concentration of cfDNA can increase in the bloodstream. ctDNA, a subset of cfDNA, refers specifically to fragmented DNA originating from tumor cells and it accounts for a fluctuating proportion of cfDNA, ranging from less than 0.1 % to over 90 % [9]. Generally, cfDNA levels are elevated in patients with carcinoma compared to healthy individuals. Significant amounts of ctDNA are released into the circulatory system through tumor cell apoptosis or necrosis, and its quantity can reflect the tumor burden in patients with cancer [9]. Table 2 provides a comprehensive overview of some studies exploring the potential of liquid biopsy tests for diagnosis, prognosis and treatment monitoring of HCC.

3.1. ctDNA for early HCC detection

cfDNA and ctDNA generally hold substantial diagnostic potential in HCC; both can provide heightened sensitivity and improved clinical correlation, enhancing their value as diagnostic tools [10]. The methylation profile of cfDNA is particularly intriguing, given that epigenetic modifications have been identified as significant contributors to tumor initiation and progression [3,31]. Several studies have reported that changes in methylation patterns of different genes can

Table 1

FDA approved liquid biopsy tests.

Liquid Biopsy Assay	Disease Type	Mutation	Manufacturer	Year approved
Cobas [®] EGFR mutation test	NSCLC	EGFR mutation (exon 19 deletions, L858R mutation)	Roche Molecular Diagnostics	2016
Guardant 360 CDx	NSCLC	ctDNA test mutations of 73 genes	Guardant Health	2022
		Some examples: EGFR mutation (exon 19 deletions, L858R, and T790M)		
		ERBB2/HER2 activating mutations (SNVs and exon 20 inser- tions)		
		KRAS G12C		
Guardant 360 CDx (newly approved)	Breast cancer	ESR1 missense mutations between codons 310-547	Guardant Health	2023
FoundationOne [®] Liquid CDx	Ovarian cancer	BRCA1, BRCA2	Foundation Medicine, Inc	2020
	NSCLC	ALK rearrangement		
	Breast Cancer	РІКЗСА		
	mCRPC	BRCA1,BRCA2,ATM		
	Lung cancer	EGFR mutations (exon 19 deletions, L858R mutation)		
	NSCLC	MET (exon 14 mutations)		
therascreen [®] KRAS RGQ PCR Kit	NSCLC	KRAS wild-type(absence of mutations in codons 12 and 13)	QIAGEN	2021
	CRC	KRAS G12C		
InVisionFirst [®] -Lung	NSCLC	37 genes NSCLC	NeoGenomics, Inc	2020
Epi proColon®	Colorectal cancer	SEPT9	Geneststr.5 Berlin	2016
Oncomine [™] Dx Target Test	NSCLC	BRAF, EGFR and IDH1 mutations	Life Technologies Corporation	2022
	cholangiocarcinoma	Chromosome abnormalities by rearrangement in <i>ROS1</i> and <i>RET</i>		
TS0500 ctDNA®	Pan-cancer	500+ genes	Illumina	2019
	Breast Cancer			
	Lung Cancer			
	Colorectal cancer			
	Castric cancer			

Table 2

Liquid biopsy studies based on ctDNA analysis in HCC.

Churcher	Trated Drawns	Number of Data	Commenter	Mala Pia dia an
Study	Tested Parameter	Number of Patients	Comparator	Main Findings
Hu et al. 2017 [19]	UBE2Q1 gene promoter methylation	80 HCC 40 LC 40 CHB	AFP (cut-off 200 ng/mL) HCC vs. LC and CHB: Sensitivity=53.8 %Specificity=87.5 % AUC _{AFP} =0.668	 a) UBE2Q1 methylation (M) performed similarly to AFP in differentiating HCC vs. LC and CHB Sensitivity= 66.3 %; Specificity= 57.5 % AUC_M = 0.619 b) M combined with AFP performed bet- ter than AFP alone and M alone in dif- ferentiating HCC vs. LC and CHB, by using different cut-off points Sensitivity= 58.8 %; Specificity= 75.0 % AUC_{M+AFP20} = 0.720 Sensitivity= 53.8 %; Specificity= 87.5 % AUC_{M+AFP200} = 0.760 Sensitivity= 37.5 %; Specificity= 88.7 %
Dong et al. 2017 [20]	RASSF1A, APC, BVES, TIMP3, GSTP1, HOXA9 methylation	98 HCC 90 CHB	AFP (cut-off 20 ng/mL) HCC vs. CHB: Sensitivity= 48 % Specificity=73.9 % AUC _{AFP} =0.609	 AUC_{M+AFP400} = 0.694 a) HCC vs. CHB Only sensitivity of <i>RASSF1A</i> methylation was greater than AFP (52%) For all the genes, specificities of methylation were better than AFP: <i>RASSF1A</i> (91.5%), <i>APC</i> (96.4%), <i>BVES</i> (97.65%), <i>TIMP3</i> (98.8%), <i>GSTP1</i> (98.7%), <i>HOXA9</i> (95.8%) b) Combination of methylation status of <i>RASSF1A</i>, <i>BVES</i>, <i>HOXA9</i> performed better than AFP in differentiating HCC vs. CHB AUC_{3gene}= 0.834 (95 %CI 0.774-0.894, p-value = 0.031) c) Combination methylation status of <i>RASSF1A</i>, <i>BVES</i>, <i>HOXA9</i> and AFP performed better than AFP alone in differentiating HCC vs. CHB AUC_{3gene+AFP}= 0.852 (95 %CI 0.796-0.908, ADC)
Kisiel <i>et al.</i> , 2019 [21]	Six-marker methylated DNA markers (MDMs)(HOXA1, EMX1,AK055957,ECE1, PFKP and CLEC11A)	95 HCC 51 LC 98 HC	AFP (cutoff 10 ng/mL) HCC vs. LC and HC Sensitivity= 60 % Specificity= 91 % AUC _{AFP} =0.80	 p-value = 0.028) a) Higher sensitivity compared to AFP alone (HCC vs. LC and HC): Sensitivity= 95 % Specificity= 92 % b) Performed better than AFP in differen- tiating HCC vs. LC and HC AUC_{6MDMspanel}= 0.96 c) 6-MDM panel demonstrated high detection rates for different BCLC stages of HCC: Stage 0: 75 %(3/4) Stage A: 93 % (39/42) Stage B: 93 % (13/14) Stage C: 100 % (28/28) Stage D: 100 % (7/7)
An et al. 2019 [22]	Mutation number of cfDNA	26 HCC 10 H 10 LC	AFP (cut-off 400 ng/mL) (a) Higher in HCC vs. H/LC, sta- tistically significant (S) p-value = 0.0368 (b) ROC curve analysis distin- guishing HCC vs. H/LC: AUC _{AFP} = 0.7827	 a) Higher in HCC patients, statistically not significant (NS) p-value = 0.0547 b) Performed better than AFP differenti- ating HCC vs. H/LC
	Maximal VAF (%) in cfDNA			a) Higher in HCC patients, NS p-value = 0.2238 b) Performed better than AFP differenti- ating HCC vs. H/LC AUC _{mayVAE} = 0.8019
	cfDNA Concentration (ng/mL)			 a) Higher in HCC patients, S p-value = 0.0249 b) Performed better than AFP differenti- ating HCC vs. H/LC AUC_{cfDNAconc} = 0.9173
	ctDNA concentration (ng/mL) (Maximal VAF x cfDNA con- centration)			a) Higher in HCC patients, NS p-value = 0.1919 b) Performed better than AFP differenti- ating HCC vs. H/LC AUC _{ctDNAconc} = 0.8712

Table 2 (Continued)

Diagnosis					
Study	Tested Parameter	Number of Patients	Comparator	Main Findings	
Lin et al., 2023 [23]	ARID1A, CTNNB1, TP53 mutations	21 HCC 14 CHB	 (a) AFP (cut-off not mentioned) Higher in HCC vs. CHB, S p-value=0.0001 (b) Clinical Factors alone (age, gender, and cirrhosis) ROC curve analysis distinguish- ing HCC vs. CHB AUC_{clinicalfactors}= 0.741 (95 % CI 0.5546-0.9284) 	a) All mutations (<i>ARID1A</i> , <i>CTNNB1</i> , <i>TP53</i>) genes higher in HCC vs. CHB <i>ARID1A</i> mutation: 85.7 % vs. 42.9 %, S p- value = 0.011; <i>CTNNB1</i> mutation: 42.9 % vs. 0 %, S p- value = 0.005; <i>TP53</i> mutation: 100 % vs. 71.4 %, S p- value = 0.019 b) Combining three mutations with the clinical factors performed better differ- entiating HCC vs. CHB AUC _{mutations+clinicalfactors} = 0.935 (95 % CI 0.8547-1.0000, p-value = 0.041)	
Prognosis and Monitoring Treatment Response					

Study	ctDNA property	Number of Patients	Prognostic Factor/Treatment	Main Findings
Oversoe et al. 2020 [24]	TERT C228T mutation	95 HCC	Overall Mortality (OM)	Presence of plasma TERT C228T mutation significantly increased overall mortal- ity:
				 a) crude HR= 3.01 (95 %CI 1.85-4.90, p-value<0.000) b) adjustment for potential confounders: TNM stage, vascular invasion, sex, age and presence of cirrhosis adjusted HR = 2.16 (95 %CI 1.20–3.88, p-
		55 treatment naïve HCC	ОМ	value= 0.01) Presence of plasma TERT C228T mutation significantly increased overall mortal- ity:
				a) crude HR= 3.88 (95 %CI 2.02–7.45, p- value<0.000)
				b) adjustment for potential confounders adjusted HR = 4.11 (95 %CI 1.73-9.76, p- value= 0.001)
		95 HCC	Advanced TNM stage	Presence of plasma TERT C228T mutation positively correlated with advanced TNM stage (p-value<0 0001)
		95 HCC	Vascular Invasion	Presence of plasma TERT C228T mutation positively correlated with vascular invasion (p.v.)ue= 0.005)
Zhao et al. 2020 [25]	ctDNA abundance	39 HCC	Maximum Tumor Diameter	a) Better correlation with tumor size than AFP
				son r=0.4, p-value=0.0091 b) Performed better in predicting tumor size, particularly tumor size with 3.5
				cm AUC _{ctDNAabundance} = 0.82, sensitivity 0.75 and specificity 1.
				AUC for AFP not reported
	ctDNA abundance	39 HCC	BCLC staging	a) ctDNA abundance higher in BCLC C patients than BCLC A and B median 25 % vs. 2 % n-value= 0.0005
	TP53 mutation	39 HCC	Overall Survival (OS)	a) Patients with <i>TP53</i> mutation had the worse OS curve, p-value= 0.0589
				b) Death in group with TP53 mutation higher than without mutation, p-value = 0.0365
	TP53 mutation	39 HCC	Progression-free survival (PFS)	Patients with TP53 mutation had the
lkeda <i>et al</i> . 2018 [26]	Characterized genomic alter- ation in 18 genes including TP53, CTNNB1, ARID1A	26 HCC	N/A	worse Pr5 curve, p-value = 0.0411 88.5 % (23/26) patients had at least one characterized genomic alteration, sug- gesting potential clinical utility for ctDNA scorement in LICC
lkeda <i>et al.</i> 2018 [27]	CDKN2A R80* nonsense muta- tion and CTNNB1 G34V mis- sense mutation	1 BCLC stage B HCV-related HCC	des-gamma carboxy prothrom- bin (DCP) levels	Patient treated with combination treat- ment with Palbociclib (CDK4/6 inhibi- tor) and celecoxib (COX-2 inhibitor). DCP levels declined by 84 % from base- line after 8 works of treatment

(continued)

Table 2 (Continued)

Diagnosis					
Study	Tested Parameter	Number of Patients	Comparator	Main Findings	
				Patient developed complications, stopped the treatment for comfort care and died.	
	MET Y501C missense mutation, TP53 R273C missense muta- tion, and PTEN L139* nonsense mutation	1 BCLC stage A HCV-related HCC	AFP levels	Patient treated with sirolimus (mTOR inhibitor) and cabozantinib (MET inhibitor), AFP levels decreased by 63 % from baseline after 4 weeks of treat- ment.	
				ctDNA analysis repeated after 8 weeks of treatment, <i>MET</i> Y501C mutation disappeared.	
An et al. 2019 [22]	Mutation number	26 HCC	Largest Tumor Diameter	Linear regression analysis: R ² = 0.1682, p-value = 0.0375	
	Maximal VAF (%)			Linear regression analysis: R ² = 0.4974, p-value <0.0001	
	ctDNA concentration (ng/mL)		Disease free survival (DEC)	Linear regression analysis: R ² = 0.2676, p-value = 0.0068	
	postoperative	 (postoperative vs. preoperative), 2 cohorts: (a) Total Clearance cohort: no somatic mutations in postop- erative blood (b) Mutational Residual Cohort: somatic mutation present 	Disease-free Survival (DFS)	word to be a solution of the solution of th	
	Clearance rate	Mutational Residual Cohort, 2 groups: (a) High clearance rate (-59.3 % to -98.5 %)	DFS	Within the Mutational Residual Cohort, those with high clearance rate of ctDNA from the bloodstream, had improved DFS than with low clearance	
		(b) Low clearance rate (265 % to -42.2 %)		rate group Median DFS 17.5 months vs. 6.7 months, log-rank HR= 3.164, p-value= 0.0195	
Alunni-Fabbroni, 2019 [28]	cfDNA concentration at different treatment points (every 8 weeks)	13 HCC grouped high or low cfDNA levels based on median values depending on time- points	OS	Patients with high cfDNA vs. low cfDNA found to had worse OS with borderline significance at latest timepoints T3, p- value= 0.057 and T4, p-value= 0.095. NS for T1 and T2 timepoints, p>0.1 for both	
	CYP2B6 mutation at different time points	13 male unresectable, advanced HCC	OS	Patients carrying the CYP2B6 mutation had worse OS in time frame T1-T2, p- value=0.01, NS for T3-T4 time points, p-value= 0.360	
Ge et al., 2021 [29]	Maximal VAF (%)	26 HCC patients	Largest tumor size	Maximal ctDNA VAF was positively cor- related with the size of the largest tumor; Linear regression analysis: R = 0.44, p-value = 0.024	
	TERT C228T Mutation	26 HCC patients	Largest tumor size	Presence of this mutation significantly correlated with largest tumor diame- ter; Linear regression analysis: R = 0.41, p-value = 0.037	
	TERT C228T Mutation	26 HCC patients	Macrovascular invasion (MVI)	Presence of this mutation correlated with MVI (p-value = 0.004)	
	TERT C228T Mutation	26 HCC patients	OS	Patients with this mutation and other ctDNA mutation not reported had sig- nificantly worse OS than patients with- out median OS 3 vs. 17.5 months, p-value =0.016	
Fu et al., 2022 [30]	Number of mutant genes in ctDNA	258 HCC patients	Recurrence free survival (RFS)	The number of total mutant genes in ctDNA was associated with early tumor recurrence cut-off at median number=6, HR =2.2,	
	Presence of high-risk genes (HRG) present in ctDNA: APC, ARID1A, CDKN2A,FAT1,LRP1B, MAP3K1, PREX2,TERT, TP53	 258 HCC patients Divided into three risk levels according to the number of mutant HRGs detected in preoperative ctDNA: (a) Low-risk: no mutant HRGs; (b) Median-risk: 1-2; (c) High-risk group: >2 mutant HRCs 	RFS	 p<0.001 Patients at the high-risk group had the lowest 1 year RFS rate while the patients at the low-risk had the highest. High-risk: 31.0 %, HR=7.1, 95 % CI 3.2-16.0, p < 0.001; Median-risk: 68.6 %, HR= 2.9, 95 % CI 1.4-6.0, p < 0.001 Low-risk: 88.2 % 	

(continued)

Table 2 (Continued)

Diagnosis					
Study	Tested Parameter	Number of Patients	Comparator	Main Findings	
FAT1, LRP1B and TP53 mutation	29 HCC patients with recurrence who received systemic ther- apy (Lenvatinib with ICIs) (preoperative and postoperative samples)	Progression Free survival (PFS)	Detection of FAT1 and LRP1B var- iants in preoperative ctDNA associated with worse PFS after systemic therapies, p- value<0.0001; p-value=0.034, respectively Presence of FAT1 or LRP1B var- iants and without TP53 var- iants had the worst PFS with median PFS time of 2.8 months, HR=17.1, p < 0.001		

HCC, hepatocellular carcinoma; LC, liver cirrhosis; CHB, chronic hepatitis B; M, UBE2Q1 methylation; H, hepatitis; HC, healthy controls; BL, Benign Lesions; NS, not statistically significant; S, statistically significant; ROC, receiver operating characteristic; AUC, area under the curve; MDM, methylated DNA markers; VAF, variant allele frequency; OM, overall mortality; OS, overall survival; PFS, progression-free survival; DFS, disease-free survival; MVI, macrovascular invasion; RFS, recurrence free survival.

distinguish HCC from controls [19,21]. An example is the study by Dong et al. in 2017, in which they found that a combination of methylation of Ras association domain family 1A (RASSF1A), blood vessel epicardial substance (BVES) and homeobox (HOXA9) gene promoters in serum and AFP outperformed AFP alone in distinguishing between HCC and chronic hepatitis B patients without HCC (CHB). Indeed, the Area Under the Curve (AUC) increased from 0.609 to 0.852 (pvalue=0.028) [20] (Table2). cfDNA and ctDNA can also harbor certain key driver mutations in common genes in HCC and their detection can serve as potential biomarkers for early detection of HCC [3]. In a study by Lin et al. in 2023, the mutations on three gene AT-rich interaction domain 1A (ARID1A), catenin beta 1 (CTNNB1), and tumor protein p53 (TP53) were detected in ctDNA. These mutations were found more prevalent in HCC patients compared to CHB patients (85.7 % vs. 42.9 %, p-value = 0.011; 42.9 % vs. 0 %, p-value = 0.005; 100 % vs. 71.4 %, p-value = 0.019, respectively). This study also demonstrated that combining the mutation profile of the three genes with clinical factors, such as age, gender, and presence of cirrhosis, improved the performance of the model for the detection of early-stage HCC in patients with HBV with the AUC increasing from 0.741 to 0.935 (pvalue=0.041) [23] (Table 2).

In another study, An *et al.* in 2019 compared different cfDNA parameters such as mutation number, maximal variant allele frequency (VAF), cfDNA concentration and ctDNA concentration between non-HCC patients, specifically hepatitis/cirrhotic patients and HCC patients. They found cfDNA concentration (*p-value* =0.0249) was significantly higher in HCC patients compared to the controls. In addition, this study also showed that all the cfDNA parameters performed equally or better than AFP in differentiating between HCC and hepatitis/cirrhosis patients [22] (Table 2).

Hence, cfDNA and ctDNA hold a potential diagnostic value in HCC, showcasing high sensitivity and robust clinical relevance. The methylation profile of ctDNA and mutations within specific genes emerge as promising tools demonstrating improved results in HCC detection, opening new possibilities for better patient management.

3.2. ctDNA to monitor HCC progression and treatment response

The analysis of ctDNA finds multiple applications in the clinical management of HCC patients [29]. Oversoe *et al.* in 2020, detected the presence of telomerase reverse transcriptase (*TERT*) C228T mutation in the ctDNA derived from 44 % of the HCC patients enrolled in the study (n=96), while it was absent in patients with cirrhosis (n=45) [24]. After adjustment for potential confounders, the presence of *TERT* mutation in plasma was associated with a higher mortality rate (adjusted HR 2.16, 95 % CI 1.20-3.88, p=0.010). On the other hand, in treatment-naive HCC patients, the presence of *TERT* C228T mutation was strongly associated with mortality (adjusted HR 4.11,

95 % CI 1.73-9.76, p=0.001). Additionally, the researchers established a positive correlation between the presence of TERT mutation in plasma and advanced TNM stage (p-value<0.0001) or vascular invasion (p-value=0.005) [24] (Table 2). Thus, highlighting the possible role of TERT C228T mutation detected in ctDNA as a prognostic marker in HCC. ctDNA has been used as a reliable biomarker to monitor tumor progression and assess treatment efficacy [28,30]. Due to the invasiveness of tissue biopsy, this procedure is highly discouraged to monitor the tumor progression, thus liquid biopsy provides an ideal option for real-time monitoring of the disease. A prospective study conducted by Zhao et al. in 2020 enrolled 42 patients with unresectable liver cancer, 39 of which had HCC [25]. The primary objective of the study was to investigate the relationship between ctDNA abundance and tumor characteristics, as along with the significance of TP53 mutations on response to interventional therapy. The results of the study revealed a strong correlation between ctDNA abundance and tumor size, highlighting the potential of ctDNA as a potential marker for quantifying tumor burden [25]. In addition, increased ctDNA levels were associated with the BCLC staging, indicating its potential as a prognostic biomarker [25]. Furthermore, patients without TP53 mutations showed improved overall survival (OS) and progression-free survival (PFS) compared to patients with TP53 mutations (OS: p-value=0.0589; PFS: p-value=0.0411) (Table 2). These findings suggest a possible association of this mutation with a less favorable response to interventional therapy, but further research with larger sample size is needed to confirm the results [25].

Indeed, the incorporation of ctDNA analysis into clinical practice enhances precision medicine approaches and improves patient outcomes in the management of HCC. Another example is a recent program called the TARGET (Tumour chARacterisation to Guide Experimental Targeted Therapy) study that aimed to assess the feasibility of using ctDNA to identify actionable mutations in early-phase clinical trials in a range of advanced-stage cancers [32]. In the initial phase, the study enrolled 100 patients, and 41 had actionable alterations. Among them, 11 received matched molecular therapies that led to a stabilization of the disease or to a partial response in patients with advanced cancers [32]. Similarly in HCC, another study identified potentially actionable alterations in ctDNA of 26 patients with advanced HCC involving 18 genes with the most common genetic alterations found in TP53, CTNNB1, and ARID1A genes [26] (Table 2). In a separate study, two HCC patients with specific somatic alterations in ctDNA received targeted drugs corresponding to their mutations, resulting in a decline in tumor marker (Alpha-Fetoprotein (AFP) and des-gamma-carboxy prothrombin (DCP)) during treatment [27]. In particular, one patient harboring a cyclin-dependent kinase inhibitor 2A (CDKN2A)-inactivating and a CTNNB1-activating mutation, received matched treatments consisting of palbociclib, a



Fig. 1. Challenges in liquid biopsy tests and offered solutions by AI in Clinical Oncology. (A): Despite the potential of liquid biopsy, particularly ctDNA, still some limitations exist that limit its clinical use, (B): AI can play a crucial role in addressing these limitations, during the study design, during the analytical phase, and (C) in the interpretation of the results thought AI-powered clinical decision support systems (CDSS). CDSS can aid in early detection, treatment selection, and monitoring of the response to treatment leading to personalized management of the patient. The Figure was created on Biorender.com, licensed under Biorender's Academic License Terms. (https://www.biorender.com/academic-license)

cyclin-dependent kinase (*CDK*)4/6 inhibitor, and celecoxib, a cyclooxygenase (*COX*)–2/*WNT* inhibitor. Within 2 months DCP levels were reduced by 84 % [27] (Table 2). Another patient harboring phosphatase and tensin homolog deleted on chromosome 10 (*PTEN*)–inactivating and a mesenchymal-epithelial transition (*MET*)–activating mutations was treated with sirolimus (mechanistic target of rapamycin (*mTOR*) inhibitor) and cabozantinib (*MET* inhibitor) [27]. Imaging conducted 6 weeks after treatment indicated stable disease with central necrosis evident within the tumors, accompanied by a reduction in AFP levels within a month. Subsequent ctDNA analysis performed after 8 weeks of cabozantinib treatment revealed the absence of the *MET* mutation (*MET* Y501C), aligning with a positive molecular response to therapy [27] (Table 2). These studies evidenced the feasibility of a liquid biopsy-driven targeted therapy that can improve patient outcome.

4. Artificial intelligence can improve ctDNA-based liquid biopsy tests

Thus far, the accumulating evidence from the literature holds promise for the prospective application of liquid biopsy in clinical oncology.

However, it should be considered that ctDNA constitutes only a minor fraction of the total cfDNA present in the plasma of cancer patients [9,10] and in the case of early or very early cancer stage it is even less abundant. For example, data derived from different studies in NSCLCs estimated the clonal mutant allele frequencies (MAFs) of ctDNA in the total cfDNA ranging from 0.01 % to 9.3 % (median 0.31 %) in patients with stage I [33]. Additionally, the "background noise " derived from white blood cells (WBC) makes the detection of

the ctDNA mutations even more difficult and less reliable [11] possibly resulting in a misinterpretation of the results or a misclassification of the patient.

New technical and analytical approaches should be adopted to address these limitations. The availability of more sensitive techniques, such as ultra-deep sequencing, that enables the identification of target ctDNA at very low concentrations [31,20], opened new opportunities in this field. This technique became cost-benefit advantageous in the case of targeted sequencing, where specific amplicons and genes are analyzed to detect known or novel mutations. However, many cancers, including HCC, lack a set of known mutations that can identify a tumor. Thus, the fundamental step that will improve liquid biopsy tests based on ctDNA is the identification of a specific mutational panel for HCC. In this regard, Artificial Intelligence (AI) can offer great help to analyze and identify specific combinations of mutations that "label" specifically the cancer, possibly using data available in genomic databases [34].

Recently, several works demonstrated the benefits derived from the application of machine learning algorithms to liquid biopsy tests. For example, Roth *et al.* used the filter-based feature method and Support Vector Machine (SVM), a supervised learning method for solving data mining problems, to filter 1158 miRNAs collected from blood to determine a suitable subset of biomarkers in glioblastoma [35]. They achieved the best performance by combining 180 miRNAs with an accuracy of 81 %, specificity of 79 %, and sensitivity of 83 % [34,35]. Another example is the combination of Single strand Adaptor Library Preparation (SALP)-sequence and SVM in the pursuit of uncovering novel cfDNA-based biomarkers for esophageal cancer, as demonstrated by Liu *et al.* [36]. In this study, the analysis of 30 cfDNA samples derived from 26 esophageal cancer (ESCA) patients and 4 healthy individuals resulted in the identification of 103 epigenetic markers and 37 genetic markers [36], thus proving the advantage conferred by AI technologies within the context of liquid-based tests.

Enhancements can be further obtained by combining the information derived from ctDNA with additional multi-omics data, including, transcriptomic, and proteomic information, all sourced from a single blood sample. Here AI methodologies play a pivotal role in seamlessly integrating these diverse layers of molecular data, illuminating concealed associations among them. By combining AI algorithms with the reservoir of information derived from liquid biopsy, researchers can gain a deeper understanding of the underlying biology of liver cancer, identify novel biomarkers, and develop more effective treatment strategies. An example of this concept is the study conducted by Chen et al. wherein they introduced an innovative integrated approach called HIFI (5-Hydroxymethylcytosine/motIf/Fragmentation/nucleosome footprint) [37]. This method was developed to diagnose HCC by analyzing cfDNA genomic features. Impressively, HIFI outperformed the conventional AFP biomarker in terms of sensitivity and specificity. This novel method emerges as a promising non-invasive and accurate strategy for early HCC diagnosis and surveillance in high-risk populations, highlighting the importance of cfDNA analysis in cancer detection [37].

5. Liquid biopsy and AI boosts the development of clinical decision support systems

AI-based clinical decision support systems (CDSS) have the potential to transform the field of liquid biopsy for liver cancer (Fig. 1). By analyzing large-scale patient data, including ctDNA profiles, treatment responses, and clinical outcomes, AI algorithms can generate predictive models and treatment recommendations [38]. CDSS powered by AI can aid clinicians in making informed decisions, selecting appropriate therapies, and monitoring treatment responses. Several groups have explored this potential and have reported promising results. An example is a study by Visser et al. which evaluated the performance of liquid biopsy-based decision support algorithms for the diagnosis and subtyping of lung cancer. They found that multiparametric models combining assessment of protein tumor markers (cancer antigen 125 (CA125), cancer antigen 15-3 (CA15.3), carcinoembryonic antigen (CEA), cytokeratin 19 fragment 21-1 (CYFRA 21-1), human epididymis protein 4 (HE4), neuron-specific enolase (NSE), pro gastrin-releasing peptide (proGRP), and squamous cell carcinoma antigen (SCCA)) and ctDNA tumor markers (ctDNA mutations in epidermal growth factor receptor (EGFR), Kirsten rat sarcoma virus (KRAS) and b-raf proto-oncogene (BRAF)) were able to identify twothirds of all lung cancer and NSCLC patients, as well as half of the small-cell lung cancer (SCLC) patients [39]. These findings underscore the specific significance of the models in scenarios where traditional lung tissue biopsies prove impractical or provide inconclusive outcomes. Furthermore, the study revealed the utility of baseline ctDNA measurements for monitoring treatment responses [39].

Now, despite these potential clinical implications, there are still some discussion on how it can be optimally integrated in the clinical decision-making process [39]. The use of AI to analyze circulating biomarkers, such as cell-free DNA (cfDNA) for early detection has become increasingly important in clinical practice as this has been credited to reduce morbidity and mortality of cancer patients. Efforts to develop and validate multi-cancer early detection (MCED) tests were made by the Circulating Cell-free Genome Atlas (CCGA), a prospective, multicenter, case-control, observational study. CCGA is funded by GRAIL, Inc and involves collaboration with various researchers and institutions. In this study, participants with and without cancer are enrolled and cellfree DNA (cfDNA) sequencing is done in combination with machine learning to detect cancer signals across multiple cancer types and predict cancer signal origin (CSO) with high accuracy [40]. The CCGA was divided into three substudies which include a comprehensive comparison of genomic sequencing approaches, refinement of the selected assay and classifiers and a large clinical validation study of the MCED test [40].

The first substudy focused on evaluating features related to cfDNA in experimental tests and prototype machine-learning classifiers [41]. It aimed to identify the most promising approach for a MCED test with a low false-positive rate and sufficient sensitivity to improve outcomes. The results showed that cfDNA methylation was the most promising genomic feature for cancer signal detection and the whole-genome methylation-based approach was selected for further development and improvement into a targeted methylation assay, and machine learning classifier for cancer detection and CSO prediction in the second substudy [41]. The third and last CCGA substudy reported on a large clinical validation study of this refined MCED test, GalleriTM MCED test. This test can detect more than 50 different cancer types and has shown high specificity for cancer signal detection (99.5 %) and accurate prediction of cancer signal origin with 4,077 study participants involving cancer and non-cancer individuals [40]. In addition, this test correctly predicted cancer tissue of origin in 88.7 % of truepositive cases. Validation in an independent set further confirmed its efficacy in detecting cancer signals [40]. These findings indicate its potential as a complementary screening tool for detecting multiple cancer types and improving clinical outcomes. Follow-up for all participants for clinical outcome of the CCGA study is still ongoing and the estimate completion is the first guarter of 2024 [42].

Several CDSS have already been developed and some are available in the market and have undergone thorough reviews [43]. While critiques have been raised about their performance due to the limitations associated with the detection and interpretation of ctDNA, as discussed in this review, AI has the potential to overcome these challenges and strengthen CDSS capabilities. Such as the case of the GalleriTM MCED blood test as previously discussed, designed to detect more than 50 different cancer types [40]. Recently concerns have been raised about its sensitivity in detecting cancers. However, a certain commitment in addressing these limitations and advancing towards more sensitive tests became evident in the recent partnership with an AI company, which aims to harness the power of AI to enhance sensitivity and improve the performance of liquid biopsy tests [44].

Recently some AI-based CDSS have been developed also for the management of HCC patients. Choi et al., developed a CDSS algorithm that can recommend the optimal initial treatment of patients with HCC and predict their OS after treatment by the use of AI and machine learning techniques to integrate various patient and tumorrelated variables [38]. This study discussed the importance of a goodquality database and selecting pretreatment variables to yield clinically meaningful results. It also highlighted the potential benefits of incorporating genetic information and imaging data to enable more personalized treatment approaches for individual patients [38]. Overall, the study was able to demonstrate the potential of AI and machine learning in improving decision-making in HCC. Another study that highlighted the use of AI in HCC was the study by Oestmann et al., which demonstrated that AI-based image analysis has the potential to expand the role of imaging-based diagnosis in primary liver cancer [45]. In this study, they used a deep learning model, specifically a convolutional neural network (CNN) to differentiate between histopathological validated HCC and non-HCC lesions on multi-phasic contrast-enhanced MRI, including lesions with atypical imaging features. Their model was able to achieve an overall accuracy of 87.3 % [45].

As far as we know, there is no information on the use of AI-based CDSS related to ctDNA or other omic data. This represents the future development for clinical hepatic oncology having the great potential to improve patient outcomes, optimize treatment strategies, and personalize liver cancer management (Fig. 1).

6. Conclusions

In conclusion, ctDNA analysis holds solid potential in HCC care. The use of ctDNA as a non-invasive alternative presents an effective solution to the limitations of tissue biopsy, including invasiveness, tumor heterogeneity, and limited tumor content. The diagnostic potential of ctDNA becomes evident in its capability to detect earlystage HCC through the analysis of methylation patterns and specific gene mutations. Additionally, ctDNA has proven to be a valuable prognostic indicator, correlating with tumor burden, disease progression, and treatment response.

Al holds the transformative potential to revolutionize the field of liquid biopsy for liver cancer. By addressing challenges like low DNA yield and the difficulty in deciphering mutation signals, AI algorithms can enhance sensitivity, improve signal-to-noise ratios, integrate multiomics data, and develop clinical decision support systems. This convergence of AI and liquid biopsy expedites discoveries and advancements within liver cancer diagnostics and treatment strategies.

Author contributions

Inah Marie Aquino reviewed the literature and drafted the manuscript; Devis Pascut reviewed the literature and made critical revisions related to the important intellectual content of the manuscript; Devis Pascut made the final approval of the manuscript submitted.

Funding

Inah Marie Aquino was supported by a scholarship from the DOST-PCHRD of the Philippines. Devis Pascut was supported by abd intramural grant from Fondazione Italiana Fegato.

Declaration of interests

None.

Acknowledgments

The authors would like to express our gratitude to Professor Claudio Tiribelli for his valuable insights and critical reading of the manuscript.

References

- [1] Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 2021;71(3):209–49. https://doi.org/10.3322/caac.21660.
- [2] Singal AG, Lampertico P, Nahon P. Epidemiology and surveillance for hepatocellular carcinoma: new trends. J Hepatol 2020;72(2):250–61. https://doi.org/ 10.1016/j.jhep.2019.08.025.
- [3] Adeniji N, Dhanasekaran R. Current and emerging tools for hepatocellular carcinoma surveillance. Hepatol Commun 2021;5(12):1972–86. https://doi.org/ 10.1002/hep4.1823.
- [4] Dhanasekaran R, Bandoh S, Roberts LR. Molecular pathogenesis of hepatocellular carcinoma and impact of therapeutic advances. F1000Research 2016;5:879. https://doi.org/10.12688/f1000research.6946.1.
- [5] Malone ER, Oliva M, Sabatini PJB, Stockley TL, Siu LL. Molecular profiling for precision cancer therapies. Genome Med 2020;12(1):8. https://doi.org/10.1186/ s13073-019-0703-1.
- [6] El-Deiry WS, Goldberg RM, Lenz H, Shields AF, Gibney GT, Tan AR, et al. The current state of molecular testing in the treatment of patients with solid tumors, 2019. CA Cancer J Clin 2019;69(4):305–43. https://doi.org/10.3322/caac.21560.
- [7] Cheng ML, Pectasides E, Hanna GJ, Parsons HA, Choudhury AD, Oxnard GR. Circulating tumor DNA in advanced solid tumors: clinical relevance and future directions. CA Cancer J Clin 2021;71(2):176–90. https://doi.org/10.3322/caac.21650.
- [8] De Rubis G, Rajeev Krishnan S, Bebawy M. Liquid biopsies in cancer diagnosis, monitoring, and prognosis. Trends Pharmacol Sci 2019;40(3):172–86. https://doi. org/10.1016/j.tips.2019.01.006.
- [9] Corcoran RB, Chabner BA. Application of cell-free DNA analysis to cancer treatment. Obstet Gynecol Surv 2018;379(18):1754–65.

- [10] Ye Q, Ling S, Zheng S, Xu X. Liquid biopsy in hepatocellular carcinoma: circulating tumor cells and circulating tumor DNA. Mol Cancer 2019;18(1):114. https://doi. org/10.1186/s12943-019-1043-x.
- [11] Chan HT, Chin YM, Nakamura Y, Low SK. Clonal hematopoiesis in liquid biopsy: from biological noise to valuable clinical implications. Cancers 2020;12(8). https://doi.org/10.3390/cancers12082277.
- [12] Arechederra M, Ávila MA, Berasain C. Liquid biopsy for cancer management: a revolutionary but still limited new tool for precision medicine. Adv Lab Med Av En Med Lab 2020;1(3):20200009. https://doi.org/10.1515/almed-2020-0009.
- [13] de Wit S, van Dalum G, Terstappen LWMM. Detection of circulating tumor cells. Scientifica 2014;2014:1–11. https://doi.org/10.1155/2014/819362.
- [14] Kwapisz D. The first liquid biopsy test approved. Is it a new era of mutation testing for non-small cell lung cancer? Ann Transl Med 2017;5(3):46. -46. https://doi. org/10.21037/atm.2017.01.32.
- [15] Kalasekar SM, VanSant-Webb CH, Evason KJ. Intratumor heterogeneity in hepatocellular carcinoma: challenges and opportunities. Cancers 2021;13(21):5524. https://doi.org/10.3390/cancers13215524.
- [16] Coto-Llerena M, Benjak A, Gallon J, Meier MA, Boldanova T, Terracciano LM, et al. Circulating cell-free DNA captures the intratumor heterogeneity in multinodular hepatocellular carcinoma. JCO Precis Oncol 2022(6):e2100335. https://doi.org/ 10.1200/PO.21.00335.
- [17] Weng J, Atyah M, Zhou C, Ren N. Prospects and challenges of circulating tumor DNA in precision medicine of hepatocellular carcinoma. Clin Exp Med 2020;20 (3):329–37. https://doi.org/10.1007/s10238-020-00620-9.
- [18] Wu X, Li J, Gassa A, Buchner D, Alakus H, Dong Q, Ren N, et al. Circulating tumor DNA as an emerging liquid biopsy biomarker for early diagnosis and therapeutic monitoring in hepatocellular carcinoma. Int J Biol Sci 2020;16(9):1551–62. https://doi.org/10.7150/ijbs.44024.
- [19] Hu N, Fan XP, Fan YC, Chen LY, Qiao CY, Han LY, et al. Hypomethylated ubiquitinconjugating Enzyme2 Q1 (UBE2Q1) gene promoter in the serum is a promising biomarker for hepatitis B virus-associated hepatocellular carcinoma. Tohoku J Exp Med 2017;242(2):93–100. https://doi.org/10.1620/tjem.242.93.
- [20] Dong X, Hou Q, Chen Y, Wang X. Diagnostic value of the methylation of multiple gene promoters in serum in hepatitis B virus-related hepatocellular carcinoma. Dis Markers 2017;2017:1–6. https://doi.org/10.1155/2017/2929381.
- [21] Kisiel JB, Dukek BA, V S R Kanipakam R, Ghoz HM, Yab TC, Berger CK, et al. Hepatocellular carcinoma detection by plasma methylated DNA: discovery, phase I pilot, and phase II clinical validation. Hepatology 2019;69(3):1180–92. https:// doi.org/10.1002/hep.30244.
- [22] An Y, Guan Y, Xu Y, Han Y, Wu C, Zhou B, et al. The diagnostic and prognostic usage of circulating tumor DNA in operable hepatocellular carcinoma. Am J Transl Res 2019;11(10):6462–74.
- [23] Lin D, Luo R, Ye Z, Wei Q, Bae H, Juon H, et al. Genomic characterization of earlystage hepatocellular carcinoma patients with Hepatitis B using circulating tumor DNA. Clin Res Hepatol Gastroenterol 2023;47(7):102161. https://doi.org/ 10.1016/j.clinre.2023.102161.
- [24] Oversoe SK, Clement MS, Pedersen MH, Weber B, Aagaard NK, Villadsen GE, et al. TERT promoter mutated circulating tumor DNA as a biomarker for prognosis in hepatocellular carcinoma. Scand J Gastroenterol 2020;55(12):1433–40. https:// doi.org/10.1080/00365521.2020.1837928.
- [25] Zhao W, Qiu L, Liu H, Xu Y, Zhan M, Zhang W, et al. Circulating tumor DNA as a potential prognostic and predictive biomarker during interventional therapy of unresectable primary liver cancer. J Gastrointest Oncol 2020;11(5):1065–77. https://doi.org/10.21037/jgo-20-409.
- [26] Ikeda S, Lim JS, Kurzrock R. Analysis of tissue and circulating tumor DNA by next generation sequencing of hepatocellular carcinoma: implications for targeted therapeutics. Mol Cancer Ther 2018;17(5):1114–22. https://doi.org/10.1158/ 1535-7163.MCT-17-0604.
- [27] Ikeda S, Tsigelny IF, Skjevik ÅA, et al. Next-generation sequencing of circulating tumor DNA reveals frequent alterations in advanced hepatocellular carcinoma. The Oncologist 2018;23(5):586–93. https://doi.org/10.1634/theoncologist.2017-0479.
- [28] Alunni-Fabbroni M, Rönsch K, Huber T, Cyran CC, Seidensticker M, Mayerle J, et al. Circulating DNA as prognostic biomarker in patients with advanced hepatocellular carcinoma: a translational exploratory study from the SORAMIC trial. J Transl Med 2019;17(1):328. https://doi.org/10.1186/s12967-019-2079-9.
- [29] Ge Z, Helmijr JCA, Jansen MPHM, Boor PPC, Noordam L, Peppelenbosch M, et al. Detection of oncogenic mutations in paired circulating tumor DNA and circulating tumor cells in patients with hepatocellular carcinoma. Transl Oncol 2021;14 (7):101073. https://doi.org/10.1016/j.tranon.2021.101073.
- [30] Fu Y, Yang Z, Hu Z, Yang Z, Pan Y, Chen J. Preoperative serum ctDNA predicts early hepatocellular carcinoma recurrence and response to systemic therapies. Hepatol Int 2022;16(4):868-78. https://doi.org/10.1007/s12072-022-10348-1.
 [31] Yang JC, Hu JJ, Li YX, Luo W, Liu JZ, Ye DW. Clinical applications of liquid biopsy in
- [31] Yang JC, Hu JJ, Li YX, Luo W, Liu JZ, Ye DW. Clinical applications of liquid biopsy in hepatocellular carcinoma. Front Oncol 2022;12:781820. https://doi.org/10.3389/ fonc.2022.781820.
- [32] Rothwell DG, Ayub M, Cook N, Thistlethwaite F, Carter L, Dean E, et al. Utility of ctDNA to support patient selection for early phase clinical trials: the TARGET study. Nat Med 2019;25(5):738–43. https://doi.org/10.1038/s41591-019-0380-z.
 [33] Abbosh C, Birkbak NJ, Swanton C. Early stage NSCLC challenges to implementing
- [33] Abbosh C, Birkbak NJ, Swanton C. Early stage NSCLC challenges to implementing ctDNA-based screening and MRD detection. Nat Rev Clin Oncol 2018;15(9):577– 86. https://doi.org/10.1038/s41571-018-0058-3.
- [34] Liu L, Chen X, Petinrin OO, Zhang W, Rahaman S, Tang ZR, et al. Machine learning protocols in early cancer detection based on liquid biopsy: a survey. Life 2021;11 (7):638. https://doi.org/10.3390/life11070638.

- [35] Roth P, Wischhusen J, Happold C, Chandran PA, Hofer S, Eisele G. A specific miRNA signature in the peripheral blood of glioblastoma patients: Glioblastoma-associated miRNA profile in peripheral blood. J Neurochem 2011;118(3):449–57. https://doi.org/10.1111/j.1471-4159.2011.07307.x.
- [36] Liu S, Wu J, Xia Q, Liu H, Li W, Xia X, et al. Finding new cancer epigenetic and genetic biomarkers from cell-free DNA by combining SALP-seq and machine learning. Comput Struct Biotechnol J 2020;18:1891–903. https://doi.org/10.1016/ j.csbj.2020.06.042.
- [37] Chen L, Abou-Alfa GK, Zheng B, Liu JF, Bai J, Du LT, et al. Genome-scale profiling of circulating cell-free DNA signatures for early detection of hepatocellular carcinoma in cirrhotic patients. Cell Res 2021;31(5):589–92. https://doi.org/10.1038/ s41422-020-00457-7.
- [38] Choi GH, Yun J, Choi J, Lee D, Shim JH, Lee HC, et al. Development of machine learning-based clinical decision support system for hepatocellular carcinoma. Sci Rep 2020;10(1):14855. https://doi.org/10.1038/s41598-020-71796-z.
- [39] Visser E, Genet SAAM, De Kock RPPA, Van Den Borne BEEM, Youssef-El Soud M, Belderbos HNA, et al. Liquid biopsy-based decision support algorithms for diagnosis and subtyping of lung cancer. Lung Cancer 2023;178:28–36. https://doi.org/ 10.1016/j.lungcan.2023.01.014.
- [40] Klein EA, Richards D, Cohn A, Tummala M, Lapham R, Cosgrove D, et al. Clinical validation of a targeted methylation-based multi-cancer early detection test

using an independent validation set. Ann Oncol 2021;32(9):1167-77. https://doi. org/10.1016/j.annonc.2021.05.806.

- [41] Jamshidi A, Liu MC, Klein EA, Venn O, Hubbell E, Beausang JF. Evaluation of cellfree DNA approaches for multi-cancer early detection. Cancer Cell 2022;40 (12):1537–49 e12. https://doi.org/10.1016/j.ccell.2022.10.022.
- [42] National Library of Medicine. The Circulating Cell-free Genome Atlas Study (CCGA). ClinicalTrials.gov; 2022. Published August 3,Accessed July 22, 2023 https://clinicaltrials.gov/study/NCT02889978.
- [43] Fernandez-Uriarte A, Pons-Belda OD, Diamandis EP. Cancer screening companies are rapidly proliferating: are they ready for business? Cancer Epidemiol Biomarkers Prev 2022;31(6):1146–50. https://doi.org/10.1158/1055-9965.EPI-22-0102.
- [44] Pons-Belda OD, Fernandez-Uriarte A, Diamandis EP. Multi cancer early detection by using circulating tumor DNA—the galleri test. reply to klein et al. the promise of multicancer early detection. comment on "pons-Belda et al. can circulating tumor DNA support a successful screening test for early cancer detection? The grail paradigm. diagnostics 2021, 11, 2171.". Diagnostics 2022;12(5):1244. https://doi.org/10.3390/diagnostics12051244.
- [45] Oestmann PM, Wang CJ, Savic LJ, Hamm CA, Stark S, Schobert I, et al. Deep learning-assisted differentiation of pathologically proven atypical and typical hepatocellular carcinoma (HCC) versus non-HCC on contrast-enhanced MRI of the liver. Eur Radiol 2021;31(7):4981–90. https://doi.org/10.1007/s00330-020-07559-1.