

REVIEW ARTICLE

25 years of Eye Movement Desensitization and Reprocessing (EMDR): The EMDR therapy protocol, hypotheses of its mechanism of action and a systematic review of its efficacy in the treatment of post-traumatic stress disorder[☆]



Patricia Novo Navarro^{a,b,c}, Ramón Landin-Romero^{d,e,f,g},
Rocio Guardiola-Wanden-Berghe^{b,c}, Ana Moreno-Alcázar^{c,d}, Alicia Valiente-Gómez^{c,d},
Walter Lupo^h, Francisca Garcíaⁱ, Isabel Fernández^j, Víctor Pérez^{b,c},
Benedikt L. Amann^{c,d,*}

^a *Departamento de Psiquiatría y Medicina Legal, Doctorado de Psiquiatría y Psicología Clínica, Universidad Autónoma de Barcelona, Barcelona, Spain*

^b *Instituto de Neuropsiquiatría y Adicciones (INAD), Consorci Parc de Salut Mar, Barcelona, Spain*

^c *Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Barcelona, Spain*

^d *FIDMAG Research Foundation, Germanes Hospitalàries, Barcelona, Spain*

^e *Neuroscience Research Australia, Sydney, Australia*

^f *School of Medical Sciences, University of New South Wales, Sydney, Australia*

^g *ARC Centre of Excellence in Cognition and its Disorders, Sydney, Australia*

^h *Clínica Logos, Barcelona, Spain*

ⁱ *EMDR España, Madrid, Spain*

^j *Psychotraumatology Research Center, Milan, Italy*

Received 9 October 2015; accepted 21 December 2015

Available online 17 March 2018

KEYWORDS

Eye movement
desensitization and
reprocessing;

Abstract Eye movement desensitization and reprocessing (EMDR) is a relatively new psychotherapy that has gradually gained popularity for the treatment of post-traumatic stress disorder. In the present work, the standardised EMDR protocol is introduced, along with current hypotheses of its mechanism of action, as well as a critical review of the available literature

[☆] Please cite this article as: Novo Navarro P, Landin-Romero R, Guardiola-Wanden-Berghe R, Moreno-Alcázar A, Valiente-Gómez A, Lupo W, et al. 25 años de Eye Movement Desensitization and Reprocessing: protocolo de aplicación, hipótesis de funcionamiento y revisión sistemática de su eficacia en el trastorno por estrés posttraumático. Rev Psiquiatr Salud Ment (Barc.). 2018;11:101–114.

* Corresponding author.

E-mail address: benedikt.amann@gmail.com (B.L. Amann).

Posttraumatic stress disorder;
Systematic review;
Psychotherapy;
Mechanism of action

PALABRAS CLAVE

Desensibilización y reprocesamiento por movimiento ocular;
Trastorno por estrés postraumático;
Revisión sistemática;
Psicoterapia;
Mecanismo de acción

on its clinical effectiveness in adult post-traumatic stress disorder. A systematic review of the published literature was performed using PubMed and PsycINFO databases with the keywords "eye movement desensitization and reprocessing" and "post-traumatic stress disorder" and its abbreviations "EMDR" and "PTSD". Fifteen randomised controlled trials of good methodological quality were selected. These studies compared EMDR with unspecific interventions, waiting lists, or specific therapies. Overall, the results of these studies suggest that EMDR is a useful, evidence-based tool for the treatment of post-traumatic stress disorder, in line with recent recommendations from different international health organisations.

© 2016 SEP y SEPB. Published by Elsevier España, S.L.U. All rights reserved.

25 años de Eye Movement Desensitization and Reprocessing: protocolo de aplicación, hipótesis de funcionamiento y revisión sistemática de su eficacia en el trastorno por estrés postraumático

Resumen El tratamiento de desensibilización y reprocesamiento por movimiento ocular (EMDR de sus siglas en inglés: eye movement desensitization and reprocessing) es una terapia relativamente novedosa que de forma progresiva ha ido ganando popularidad en el tratamiento del trastorno por estrés postraumático. El objetivo de este trabajo es introducir el protocolo estándar EMDR, ofrecer una revisión de las hipótesis actuales sobre su mecanismo de acción y analizar la evidencia científica disponible sobre su eficacia clínica en pacientes adultos con diagnóstico de trastorno por estrés postraumático. Se realizó una revisión sistemática de la literatura publicada en las bases de datos PubMed y PsycINFO con los términos «eye movement desensitization and reprocessing» y «posttraumatic stress disorder» y sus contracciones en inglés «EMDR» y «PTSD». Se obtuvieron como resultado 15 ensayos controlados aleatorizados de elevada calidad metodológica que compararon EMDR con tratamientos no específicos, lista de espera y con tratamientos específicos. Los resultados de estos estudios permiten concluir que EMDR es una herramienta útil y basada en evidencia científica, tal y como refleja su reciente recomendación como tratamiento de elección en el trastorno por estrés postraumático por parte de distintas organizaciones internacionales de salud.

© 2016 SEP y SEPB. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Post traumatic stress disorder (PTSD) is a psychiatric disorder classified within the group of anxiety disorders. PTSD may develop following exposure to a traumatic event in which there is a real risk of death, a threat or suffering severe injuries. PTSD has a highly significant negative repercussion on patients' lives, and its prevalence stands at from 7% to 8% of the population.^{1,2} According to the fourth edition of the Diagnostic and statistical manual of mental disorders, DSM, this disorder includes 4 main groups of symptoms: (i) re-experiencing the traumatic event in the form of nightmares, flashbacks and distressing thoughts; (ii) physiological hyperactivation that manifests as increased irritability, hypervigilance, sleep alterations and/or difficulty in concentrating; (iii) behaviour that avoids people or places associated with the traumatic events, developing feelings of detachment and/or making efforts to avoid certain thoughts, and (iv) persistent negative alterations in thoughts as well as in mood.³

There are currently several non-pharmacological therapeutic approaches in the treatment of PTSD⁴: cognitive-behavioural therapy (CBT) that centres on the trauma, exposure therapy and treatment by eye movement desensitization and reprocessing (EMDR). In recent years EMDR has attained a significant popularity, with increasing numbers of therapists being trained in the technique and the subsequent considerable increase in its clinical use in different European countries.⁵ There has also been a significant increase in the number of scientific publications indexed in PubMed about the efficacy of using PTSD for other indications too, together with hypotheses about its mechanism of action (Fig. 1).

Due to the growing popularity of EMDR and the arrival of the 25th anniversary of its introduction as a treatment, this work contains a review of what EMDR consists of as well as its standard protocol for use and the current hypotheses on the mechanisms of action that underlie the therapy. Finally, it will systematically evaluate the literature which has examined the efficacy of EMDR treatment in cases of PTSD in comparison with non-specific treatments or remaining in a




Figure 1 Evolution of the number of publications on eye movement desensitization and reprocessing indexed in PubMed in the period 1989–2014.

waiting list, other specific interventions and pharmacological treatment.

EMDR: basic concepts and application protocol

EMDR is a psychotherapy that was developed in the 1980s by the North American psychologist Francine Shapiro.⁶ Shapiro originally described a standard intervention protocol that centred on working on traumatic memories and the stress symptoms associated with them. The protocol has been refined over time and by the addition of new knowledge until the standard protocol now in use was obtained.^{7–11} The most striking and controversial aspect of EMDR may be the use of bilateral stimulation, in the majority of cases by horizontal saccadic eye movements, to desensitise the discomfort caused by traumatic memories. This aims to reprocess these memories and include them within the normalised biographical memories of the patient.¹¹

The standard protocol now includes the 8 phases which are described below:

- (1) Patient history: in this first phase the therapist centres on evaluating the attachment history of the patient and on identifying traumatic episodes and how they connect with their current symptoms. The patient's medical history and current physical problems are also recorded. The bases and foundations of the therapy are then explained and a treatment plan is prepared. The traumatic memories to be worked on are selected, always following the temporal pattern of past-present-future.
- (2) Patient preparation: the different forms of bilateral stimulation are tested in the second phase:
 - Horizontal eye movements: the therapist moves his fingers within the horizontal plane of the patient's visual field, so that the latter follows them with his

eyes, keeping the head immobile, at a speed of 2 movements per second and up to a total of approximately 30–40 movements.

- "Tapping": this consists of gently and alternately tapping the backs of the patient's hands, which have to be resting on his knees. This too should be at a speed of 2 taps per second, although the speed and intensity may vary, depending on the patient.
 - Bilateral auditory stimulation: this may consist of tones or bilateralised music played through headphones given to the patient.
- (3) Evaluation of the primary aspects of the memory: this step basically consists of accessing the traumatic memory and identifying the thoughts, emotions and bodily physical sensations associated with it. The therapist asks the patient which image represents the worst part of the traumatic memory and which negative cognition about the self that is a statement that can be generalised and pronounced in the present time they associate with that image (for example, "I am a failure"). The patient is then asked to identify the most suitable positive cognition to replace the negative one. This new cognition must express what the patient would like to believe about himself when thinking of the image (for example, "I am a valid person"). The patient evaluates the degree to which the positive cognition is true when thinking of the traumatic memory image on a scale of from 1 to 7, where 1 is completely false and 7 is completely true. Finally, the patient identifies which emotions he feels when bringing the traumatic experience and negative cognition to mind, as well as the level of alteration in units of subjective discomfort. The Subjective Units of Distress scale is used for this, with scores of from 0 to 10, where 10 is the maximum level of discomfort and 0 is the minimum.

- (4) Desensitisation of the memory: the therapist asks the patient to bring the traumatic image and negative cognition to mind, together with the associated emotion and physical discomfort which this causes at the present time. The therapist then starts bilateral stimulation (eye movement, "tapping" or auditory stimulation) for 30–40 s. Eye movements are general advisable, although if these are not tolerated well by the patient then the other 2 forms of bilateral stimulation may be used. During the periods of stimulation the therapist informs the patient that they should be a mere observer of what happens without making any type of judgement. After each round of stimulation the patient is given time to express what they have thought or felt, without the therapist making and type of comment, judgement or interpretation. Once new material ceases to emerge and the traumatic image no longer causes discomfort in the present (obtaining a score on the Subjective Units of Distress scale of 0 or 1 at most) the next phase commences.
- (5) Installation of the positive cognition: the therapist now asks the patient to bring the positive cognition to mind and associated it with the original experience. Further periods of bilateral stimulation then take place to link the positive cognition with the original memory.
- (6) Body scan: the patient is asked to close their eyes and concentrate on the original experience and the positive cognition, observing their body in case any type of feeling arises. If the patient expresses any negative or unpleasant sensation, the therapist will commence bilateral stimulation until it disappears. If the sensation is positive it is reinforced using periods of bilateral stimulation lasting for 10–12 s.
- (7) Debriefing: the possible effects that may occur after the EMDR session are explained, such as the emergence of "insights", new thoughts, memories or even dreams and nightmares, and recommendations about what to do in each cases are made.
- (8) Re-evaluation: this has the aim of evaluating the effect that the patient may have experienced after the last session, re-evaluating the memory that was worked on to check whether it has been functionally processed.

The number of sessions dedicated to each phase as well as the number of phases worked on in each session may vary from one patient to another.

Hypothesis on how EMDR works

EMDR is an eclectic psychotherapy which contains elements that are compatible with the majority of psychological approaches.¹¹ For example, the importance of childhood memories clearly fits with the psychodynamic model, while the exercise of keeping the image of the trauma in mind is similar to the exposure in imagination used in trauma-centred CBT.¹² As well as being an experience-based client-centred method, EMDR also fits with the concept of

positive and negative judgements about oneself that has deep roots in the field of phenomenological and humanist therapy.¹³ Due to all of this, some authors have suggested that the effectiveness of EMDR basically arises thanks to the common effects of different therapies.¹⁴ Nevertheless, Shapiro and Maxfield¹⁵ propose a theoretical model in which they offer data indicating that rhythmic bilateral stimulation is a fundamental and indispensable part of EMDR. The theoretical framework proposed by Shapiro to explain the observed clinical efficacy of EMDR is the Adaptive Information Processing model, AIP-model, which involves recovering the information about past negative experiences to re-include them in a positive emotional and cognitive framework.¹¹ The core component of EMDR consists of keeping the attention focused on two different things, the traumatic memory and the rhythmic bilateral stimulation. This state of attention is fundamental as it is able to induce certain physiological conditions that activate information processing. Hypothetically bilateral stimulation is able to unblock brain processing centres, creating a connection between the previous adverse information and a response to a normal non-traumatic stimulation. Bilateral stimulation causes a response of relaxation and a series of physiological responses that, when they connect with stored information on previous adverse experiences, generate new information that is functionally reintegrated.

Given the theoretical framework proposed by Shapiro, study of the role of bilateral stimulation and of eye movements most of all, has aroused the curiosity of many researchers who are interested in EMDR. The first studies of the effect of eye movements were undertaken by Wilson et al.¹⁶ and Andrade et al.¹⁷ Based on the results of these studies, 3 main hypotheses have been suggested to explain the effect of eye movements.

The first hypothesis is that eye movements increase the connection between hemispheres. According to this model, PTSD symptoms are due to a fault in the processing of episodic memories. Bilateral eye movements would facilitate interaction between the hemispheres by increasing neuronal activation and connectivity to promote processing of the information. The result would be an improvement in the capacity to remember all of the elements of the traumatic event, facilitating the inclusion of the episodic memory in semantic memory.^{18–20} However, some recent electroencephalogram studies do not support this first theory, as they show that eye movements do not aid communication between the 2 hemispheres of the brain.^{20,21} This has led to other mechanisms being proposed within this model, in connection with changes in functional brain connectivity.²²

The second hypothesis states that eye movements lead to physiological changes similar to those that occur during the REM (rapid eye movement) sleep phase.²³ This model holds that the integration of episodic memory within the semantic memory occurs during sleep. Repeated bilateral eye movements would activate neurological mechanisms similar to those of the REM phase through an orientation response, thereby making it possible to include traumatic memories. There is now one controlled study which shows an improvement in sleep alterations in PTSD patients who receive EMDR in comparison with a control group.²⁴ Within the physiological model, some authors have also suggested

that bilateral stimulation would activate an exploratory reflex which places the organism in a state of alert; in the context of the therapy, as there are no real dangers, a relaxation response would arise that would involve a reduction of arousal, thereby facilitating the reprocessing and integration of the traumatic memory.^{25–28}

The third and last hypothesis is that eye movements induce a demanding attentional task which has the effect of “exhausting” the working memory.²⁹ As the working memory has a limited capacity, the dual attentional task of (i) keeping the traumatic content and its associated characteristics “online”, while at the same time, (ii) concentrating on bilateral eye movements may alter the quality of storage of traumatic memories, reducing the intensity of episodic memories and, with this, the symptoms of PTSD.³⁰ Some authors, such as Gunter and Bodner,²⁹ consider that the unpleasant memories worked on during EMDR sessions are kept in the visual-spatial agenda, as one of the components of the working memory that is in charge of maintaining and handling visual or spatial information. According to these authors, as the therapy progresses the stored content gradually becomes less vivid and emotional as the eye movements exhaust processing resources.

Recommendation by international clinical associations

The scientific evidence for the efficacy of EMDR in the treatment of PTSD has been recognised by several international associations. The American Psychiatric Association has awarded it a level of confidence II, equivalent to a “recommendation with moderate clinical reliability”.³¹ The American Psychological Association considers this treatment to have “strong research support”.³² According to the National Institute for Care Excellence (NICE),³³ EMDR may be recommended with degree of evidence A for the treatment of PTSD in adults, with the backing of 11 controlled randomised studies. Additionally, in 2013 EMDR was included in the guide of the World Health Organisation as the first option for the treatment of PTSD in adults, children and adolescents.³⁴

Systematic review of studies of the clinical efficacy of EMDR in PTSD

The scientific evidence for treatment using EMDR has been taken from different reviews and meta-analyses. The first meta-analysis of the efficacy of EMDR was published by Davidson and Parker in 2001,¹⁴ and it included 34 studies. The results showed that EMDR was equally effective as other exposure techniques. In another meta-analysis performed in 2006, Seidler and Wagner³⁵ reported that trauma-centred EMDR and CBT tended to be equally effective in the treatment of PTSD. Similarly, the systematic review by Bisson et al.⁴ in 2013 concluded that EMDR and CBT centred on trauma have very similar effects, and they stated that both treatments have stronger empirical backing than stress-management programmes. Finally, a meta-analysis published in 2014 that included 26 controlled randomised

trials found that EMDR therapy significantly reduces the symptoms of PTSD, depression and anxiety.³⁶

Objectives

The aim of this review is to identify, synthesise and critically evaluate research into EMDR treatment in adults with PTSD diagnosed according to the DSM-IV. The increasing popularity of EMDR, the increase in the number of available studies in databases and the recent publication of protocols adapted to other mental illnesses (by Amann et al.,³⁷ for example) show that this review is timely. Although previous reviews in this field exist,^{4,35} they include studies of doubtful methodological quality and have a predominantly narrative approach. This review is an advance as it identifies and includes 15 controlled randomised studies that were evaluated using the Jadad scale³⁸ to determine their methodological quality. This scale covers distortions due to randomisation, masking and losses during follow-up. The scale runs from 0 to 5 points, where 5 points is considered to refer to a rigorous randomised clinical trial and where less than 3 points indicates a low quality clinical trial. For this study works that obtained a score of 3 or more were selected.

Lastly, this work was undertaken using a systematic review methodology that performs a critical evaluation of research techniques and the results of the studies included, adapting the Preferred Reporting Items for Systematic Reviews and Meta-Analyses^{39,40} into Spanish. This is a set of 27 items based on the evidence established by minimum criteria for the presentation of reports in systematic reviews and meta-analyses.

Method

A systematic review was undertaken of scientific papers that evaluated the efficacy of EMDR in the treatment of PTSD. These papers were identified in a search of the bibliography using systematic review techniques. Following the Thesaurus of the National Medicine Library (Medical Subject Heading Terms, MeSH) and the Psychological Index Terms of the American Psychological Association it was decided to use the following as the most appropriate search terms: “eye movement desensitization and reprocessing” and “post-traumatic stress disorder”, as well as their acronyms in English “EMDR” and “PTSD”. The final search equation was defined using the Boolean connector OR in each one of the 2 equations. The databases consulted were Medline (via PubMed) and PsycINFO. The search was restricted to from 1/1/1984 to 31/12/2014. It was restricted to human beings and no subheadings or tags were used, so that the search took place in all fields. A search was also carried out of the secondary literature, with the aim of covering other forms of publication. In this way the bibliographies of the papers selected in the main search were examined to detect studies that had not been identified in the computerised search.

The final selection of the papers was undertaken using the following inclusion criteria: (i) original papers published in peer-reviewed journals; (ii) adult population (over the age of 16 years old); (iii) patients diagnosed PTSD according to DSM criteria, when the version used corresponded to the




Figure 2 Search methodology (flow chart).

year the original work was published: DSM-III-R (APA, 1987), DSM-IV (APA, 1994) or DSM-IV-TR (APA, 2000), and (iv) a trial design with a control group.

The exclusion criteria were: (i) studies that were not original (reviews, meta-analyses, other types of experiment); (ii) studies of the basic working mechanisms of EMDR, and (iii) quasi-experimental studies (of a single group or a single case). Studies were selected by the authors PNN and BLA. Disagreements were resolved by RGWB.

After selecting the 18 papers that fulfilled the inclusion and exclusion criteria described above the Jadad scale was used to evaluate the methodological quality of each clinical trial.³⁸ Absolute and relative frequencies were calculated for statistical analysis using SPSS 16.0 for Windows software (SPSS, Inc., Chicago, IL, U.S.A.).

Results

Fig. 2 shows a flow chart that describes the selection of the studies. A total of 852 original works were identified, of which 266 were found in Medline and 586 in PsycINFO. When reviewing the bibliographies of the papers found, one work was selected that had not been detected in the primary search. When both searches were compared 140 duplicated works were found, as well as 390 papers that were not original which were eliminated, leaving 323 original papers. Of the resulting 323 papers, 305 were excluded as they did not use an adult population ($n = 47$), as they did not use a clinical sample of PTSD ($n = 88$), as they studied the mechanism of action of EMDR ($n = 82$) and as their clinical trials were neither randomised nor controlled ($n = 88$). After applying the Jadad scale questionnaire 3 works were excluded due to

their poor technical quality, resulting in the final selection of the 15 papers included in this review, with an average methodological quality of 4 points (Jadad). In general the methodological quality of the clinical trials was found to be appropriate, and it was also found to have improved over time. As well as the Jadad criteria, the number of sessions was reviewed, together with the replicability of the treatments and/or whether they were manual, the training of therapists and supervision of the therapy by an independent evaluator. The work also fulfilled the majority of the criteria in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (19 of 27 criteria), excepting those on the quantitative study of results.

Analysis of current scientific production gives rise to the following statistics: the average age of the papers was 11.73 ± 1.37 years (CI 95% 8.79–14.67), with a minimum of 2 years and a maximum of 20 years. The obsolescence of the papers analysed using the median (Burton–Kebler index) was of 12 years, and the Price index stands at 13.33 (the percentage of papers less than 5 years old).

Clinical trials of EMDR in post-traumatic stress disorder

EMDR versus waiting list group and/or non-specific treatments

Of the 15 clinical trials included (Table 1), 4 of them compared the efficacy of EMDR with non-specific treatments.^{41–44} All of these studies showed a significant clinical improvement with EMDR in comparison with other treatments. The average methodological quality of these works was 4.5 points (Jadad).

More specifically, EMDR was compared with waiting list,^{41,44} with active listening according to Rogers' model⁴³ and with counselling.⁴² In the studies that evaluated the symptoms of PTSD by self-reporting and standardised clinical interviews, EMDR showed a significant reduction in the symptoms of PTSD, as well as the symptoms of depression and anxiety in comparison with the non-specific treatment group. Some studies also showed a significant reduction in dissociative symptoms⁴¹ and an improvement in self-concept⁴³ when EMDR was used in comparison with patients in waiting lists or those who received no specific treatment.

In 3 of the 4 studies included the therapeutic effect of EMDR remained after 3 months,⁴² after 6 months⁴¹ and after 35 months⁴⁴ of follow-up. Some studies observed an association between the improvement in symptoms and increased adherence to the treatment^{41,44} and improved training for the therapists.^{42–44}

EMDR versus specific treatments

11 clinical trials were found that compared the efficacy of EMDR with other specific treatments.^{45–55} EMDR was compared to long-term imaginary exposure,^{45,49,51} muscle relaxation,^{45,51} biofeedback assisted muscle relaxation,⁴⁶ trauma-centred CBT,⁵⁰ prolonged exposure with cognitive restructuring,⁴⁸ a stress inoculation programme,⁵⁰ pharmacological treatment,⁵³ emotional expression techniques⁵⁴ and short eclectic therapy.⁵⁵

EMDR was found to be equally effective as other specific techniques in 5 studies (Jadad 3.8 points). The first study

Table 1 Studies of the efficacy of eye movement desensitization and reprocessing in post-traumatic stress disorder in adults (n = 15).

	Study name	Jadad: 3–5	No.	Interventions compared	Population treated (n)	Number of sessions, EMDR experience, evaluator/constancy	Follow-up	Results	Main conclusions
1	Vaughan et al. (1994) ⁴⁵	3	36	EMDR Imagined exposure Muscle relaxation used Waiting list	12 13 11 17 initial phase (subsequently randomised between interventions)	EMDR: 3–5 sessions Imagined exposure: 3–5 sessions Muscle relaxation used: 3–5 sessions Waiting list: 2–3 weeks Experience: NR Evaluator/constancy: NR	3 months	Scales administered by others: SI-PTSD Self-administered scales: IES, STAI, BDI, HRSD	All of the groups improved significantly in comparison with the Waiting list and benefits were maintained in the follow-up at 3 months. The results indicate that EMDR is more effective for intrusive images after the end of the treatment
2	Rothbaum (1997) ⁴¹	3	18	EMDR Waiting list	10 8	EMDR: 3 sessions. Waiting list: NR. Experience: NR. Evaluator/constancy: yes	3 months	Self-administered scales: PSS-SR, IES, BDI, STAI	EMDR significantly improved PTSD symptoms and depressive symptoms after treatment
3	Marcus et al. (1997) ⁴²	5	67	EMDR Conventional care	33 34	EMDR: 3 sessions. Conventional care: 3 sessions. Experience: level I and II (3–5 years). Evaluator/constancy: no	Follow-up work; Marcus et al. (2004). 3 and 6 months	Scales administered by others: GAF. Self-administered scales: M-PTSD, PSS-SR, IES, STAI, SLC-90, SUD	EMDR showed a significant improvement and with greater rapidity than the control group in measurements of PTSD, depression, anxiety and general symptoms. In the follow-up the results remain stable
4	Scheck et al. (1998) ⁴³	5	60	EMDR Active listening	30 30	EMDR: 2 sessions. Active listening: 2 sessions. Experience: level I and II. Evaluator/constancy: no	3 months	Scales administered by others: PTSD-I. Self-administered scales: BDI, STAI, PENN, IES, TSCS	EMDR showed a significant improvement in anxiety, depression, self-concept and PTSD symptoms. In follow-up at 3 months the difference between groups reduced

Table 1 (Continued)

Study name	Jadad: 3-5	No.	Interventions compared	Population treated (n)	Number of sessions, EMDR experience, evaluator/constancy	Follow-up	Results	Main conclusions
5 Carlson et al. (1998) ⁴⁶	3	35	EMDR Biofeedback assisted relaxation Waiting list	10 13 12	EMDR: 12 sessions. Biofeedback assisted relaxation: 12 sessions. Habitual treatment: 6 weeks. Experience: NR. Evaluator/constancy: yes	3 months	Scales administered by others: CAPS Self-administered scales: SI-PTSD-SR, Mississippi, IES, STAI, BDI	EMDR showed a significant improvement in comparison with the other 2 conditions in anxiety, depression and PTSD symptoms after treatment and in the follow-up
6 Devilly y Spence (1999) ⁴⁷	3	23	EMDR Trauma-centred CBT	12 11	EMDR: 8 sessions. Trauma-centred CBT: 9 sessions. Experience: level I and II. Evaluator/constancy: yes	3 months	Scales administered by others: PTSD-I. Self-administered scales: STAI, BDI, SLC-90, SUD, PPD, CMS, IES, PSS-SR, CEQ, DEVS-T	Trauma-centred CBT is better than EMDR in anxiety and PTSD symptoms after treatment and in the follow-up
7 Power et al. (2002) ⁴⁸	4	72	EMDR Prolonged exposure with cognitive restructuring Waiting list	27 21 24	EMDR: 10 sessions. Prolonged exposure with cognitive restructuring: 10 sessions. Waiting list: 10 weeks. Experience: level I and II. Evaluator/constancy: yes	15 months	Scales administered by others: CAPS, MADRS, HAM-A Self-administered scales: SI-PTSD-SR, IES, BDI, STAI, HADS	EMDR and prolonged exposure are more effective than the Waiting list. EMDR shows greater reduction in depression and better social functioning after treatment. After follow-up: a slight gain in depressive symptoms
8 Ironson et al. (2002) ⁴⁹	3	22	EMDR Prolonged exposure	10 12	EMDR: 6 sessions. Prolonged exposure (12): 6 sessions. Experience: level I and II. Evaluator/constancy: yes	3 months	Self-administered scales: PSS-SR, BDI, SUD, DES	Both are equally effective in PTSD symptoms and depression after treatment and in the follow-up. EMDR is faster and with a lower rate of abandonment

Table 1 (Continued)

	Study name	Jadad: 3–5	No.	Interventions compared	Population treated (n)	Number of sessions, EMDR experience, evaluator/constancy	Follow-up	Results	Main conclusions
9	Lee et al. (2002) ⁵⁰	3	24	EMDR Stress inoculation programme with prolonged exposure	12 12	EMDR: 7 sessions. Stress inoculation programme: 7 sessions. Experience: level I and II. Evaluator/constancy: yes	3 months	Scales administered by others: SI-PTSD Self-administered scales: MMPI-K, IES, BDI	EMDR is more effective in intrusive symptoms after treatment and with more gain in depressive and PTSD symptoms in the follow-up
10	Taylor et al. (2003) ⁵¹	4	60	EMDR Prolonged exposure Relaxation training	19 22 19	EMDR: 8 sessions. Prolonged exposure: 8 sessions. Relaxation training: 8 sessions. Experience: level I and II. Evaluator/constancy: yes	No follow-up.	Scales administered by others: SCID-IV, CAPS Self-administered scales: PSS-SR, BDI	Prolonged exposure produces significant improvements in avoidance symptoms and re-experiencing. EMDR and relaxation do not differ in efficacy or speed
11	Rothbaum et al. (2005) ⁵²	4	60	EMDR Prolonged exposure Waiting list	20 20 20	EMDR: 9 sessions. Prolonged exposure: 9 sessions. Waiting list: 4–5 weeks. Experience: NR. Evaluator/constancy: NR	6 months	Scales administered by others: CAPS All, SLESQ, SCID-NP. Self-administered scales: PSS-SR, IES, BDI, DES, STAI	EMDR and prolonged exposure are equally effective in depression, dissociation and anxiety after treatment and in the follow-up. Both are superior to the Waiting list
12	Van der Kolk et al. (2007) ⁵³	5	88	EMDR Prozac Placebo	29 19 29	EMDR: 8 sessions. Prozac: 8 sessions. Placebo: 8 weeks. Experience: level I and II. Evaluator/constancy: yes	6 months	Scales administered by others: CAPS, SCID-I, SCID-II. Self-administered scales: BDI	Data showing the superiority of EMDR after treatment were found and also in the follow-up in reducing PTSD and depression symptoms

Table 1 (Continued)

	Study name	Jadad: 3–5	No.	Interventions compared	Population treated (n)	Number of sessions, EMDR experience, evaluator/constancy	Follow-up	Results	Main conclusions
13	Hogberg et al. (2007) ⁴⁴	5	24	EMDR Waiting list	13 11	EMDR: 5 sessions. Waiting list: 8 weeks. Experience: level I and II. Evaluator/constancy: yes	Follow-up work; Hogberg et al. (2008), ⁷² 35 months	Scales administered by others: SCID-I, GAF, HAMA-A, HAMA-D. Self-administered scales: IES, BAI, SDI, WHO-10	EMDR shows a significant improvement in PTSD criteria after treatment, as well as for anxiety and depression. In Hogberg et al. (2008), the results remain stable after 35 months
14	Karatzias et al. (2011) ⁵⁴	5	46	EMDR Emotional expression techniques	23 23	EMDR: 8 sessions. Emotional expression techniques: 8 sessions. Waiting list: 8 weeks. Experience: NR. Evaluator/constancy: yes	3 months	Scales administered by others: CAPS. Self-administered scales: PCL-C, HADS-D, HADS-A, SWLS	Both interventions showed significant therapeutic benefits after intervention and in the follow-up, although some patients with EMDR experienced more pronounced clinical changes
15	Nijdam et al. (2012) ⁵⁵	5	140	EMDR Short eclectic therapy	70 70	EMDR: 15 sessions. Short eclectic therapy: 15 sessions. Experience: level I. Evaluator/constancy: yes	No follow-up	Scales administered by others: SI-PTSD, SCID-I. Self-administered scales: IES, HADS	Both treatments are equally effective in reducing PTSD symptoms, indicating that the effects of EMDR are faster

All: Assault Information Interview; BAI: Beck Anxiety Inventory; BDI: Beck Depression Inventory; CAPS: Clinician-Administered PTSD Scale; CEQ: Credibility/Expectancy Questionnaire; CMS: Mississippi Scale for Civilian PTSD; DES: Dissociative Experiences Scale; DEVS-T: Distress Evaluation Scale for Treatment; EMDR: eye movement desensitization and reprocessing; GAF: Global Assessment of Functioning; HADS-A: The Hospital Anxiety and Depression Scale-Anxiety; HADS-D: The Hospital Anxiety and Depression Scale-Depression; HAMA-A: Hamilton Anxiety Rating Scale-Anxiety; HAMA-D: Hamilton Anxiety Rating Scale-Depression; HRSD: Hamilton Rating Scale for Depression; IES: Impact of Event Scale; Jadad: scale that measures the methodological quality of studies; MADRS: The Montgomery-Asberg Depression Rating Scale; Mississippi: Mississippi Scale for Combat-Related PTSD; MMPI-K: Minnesota Multiphasic Personality Inventory K-Scale; M-PTSD: Modified PTSD Scale; NR: not reported; PCL-C: Posttraumatic Stress Disorder Checklist, civilians version; PENN: Penn Inventory for Posttraumatic Stress Disorder; PPD: Personal Problem Definition Questionnaire; PSS-SR: PTSD Symptom Scale Self-Report; PTSD: post-traumatic stress disorder; PTSD-I: PTSD Interview; SCID-I: Structured Clinical Interview for DSM-IV Axis I Disorders; SCID-II: Structured Clinical Interview for DSM-IV Axis II Disorders; SCID-IV: Structured Clinical Interview for DSM Disorders IV; SCID-NP: Structured Clinical Interview for DSM-IV Non-Patients Version; SDI: Social Disability Index; SI-PTSD: Davidson's Structured Interview for PTSD; SLC-90: Symptom Checklist-90; SLESQ: Stressful Life Events Screening Questionnaire; SR: self-report; STAI: State-Trait Anxiety Inventory; SUD: Subjective Units of Distress; SWLS: Satisfaction with Life Scale; CBT: cognitive-behavioural therapy; PTSD: post-traumatic stress disorder; TSCS: Tennessee Self-Concept Scale; WHO-10: World Health Organization Ten Well-Being Scale.

compared EMDR with muscle relaxation and imaginary exposure without finding any significant differences between the groups.⁴⁵ Similarly, 2 clinical trials found that there were no significant differences when EMDR was compared to prolonged exposure,^{49,52} as both interventions were found to be equally useful in treating PTSD. Nevertheless, the study by Ironson et al.⁴⁹ found that EMDR was faster in reducing symptoms and that it was tolerated better by patients, given that a lower rate of abandonment was recorded. Karatzias et al.⁵⁴ showed that EMDR and emotional liberation techniques produced significant therapeutic benefits, although they state that there was a slightly higher proportion of patients with clinical improvement in the group that received EMDR. The authors hypothesise that these results were due to the fact that both therapies share fundamental elements, such as desensitisation, distraction and exposure. Lastly, Nijdam et al.⁵⁵ compared EMDR with short eclectic therapy and found that both treatments were equally effective in reducing the symptoms of PTSD, and they indicate that the effects of EMDR were faster.

Four clinical trials with an average quality of 3.8 points (Jadad) obtained favourable results with EMDR in comparison with specific techniques. The comparison of EMDR with biofeedback-assisted relaxation training⁴⁶ showed results in favour of EMDR in reducing the symptoms of anxiety, depression and PTSD after treatment and after a follow-up period of 3 months. The work by Power et al.⁴⁸ found that EMDR was superior to prolonged exposure with cognitive restructuring in reducing depressive symptoms and improving social functioning after treatment, and these effects were maintained after a 15 month follow-up. EMDR was also found to be more effective than the stress inoculation programme with gradual exposure⁵⁰ for the symptoms of avoidance and PTSD flashbacks after treatment and after a 3 month follow-up. Only one study compared the efficacy of EMDR with pharmacological treatment. Van der Kolk et al.⁵³ compared a group of patients with PTSD who were treated using EMDR with another group of patients treated with a placebo and another group treated using antidepressive drugs. The results indicated that the intervention with EMDR was more effective than the pharmacological therapy in achieving a substantial and sustained reduction at 6 months of the PTSD symptoms as well as depression in the majority of the patients.

Lastly, 2 trials were found with unfavourable results for EMDR, with an average methodological quality of 3.5 points (Jadad). Devilly and Spence (1999) compared the efficacy of EMDR with trauma-centred CBT. Their results indicated that CBT was more effective and better tolerated than EMDR after treatment and after a 3 month follow-up. In the second study Taylor et al.⁵¹ showed that prolonged exposure was more effective than EMDR in reducing avoidance and flashback symptoms, and that it was also faster in reducing avoidance symptoms.

Discussion

As it is the 25th anniversary of the introduction of EMDR, this work has the aim of briefly illustrating the history and development of this treatment. It reviews its basic administration protocol, shows the results of recent studies of

its mechanism of action and reviews its clinical efficacy. A systematic review was carried out controlled randomised trials with appropriate methodological quality and in adult populations diagnosed PTSD according to DSM-IV criteria. The review was performed according to the standardised methodological quality criteria for conducting systematic reviews.

Since it was introduced EMDR has generated a considerable amount of debate in the clinical and scientific communities. This mainly centres on the role of bilateral stimulation (and more specifically of eye movements) as an active part of treatment, and the uncertainty about whether its efficacy is due to overlap of its elements with other psychotherapeutic techniques. Eye movements were originally described as the "crucial" part of EMDR.⁶ This argument was subsequently revised,⁹ concluding that dual attention is the causal mechanism for the effect of EMDR. Some studies have found that eye movements make a unique contribution to the success of the intervention,^{13,17,56,57} while others have found no differences in the result when the same procedure with and without eye movements is compared.^{14,58} These studies are difficult to compare, as they differ widely in methodological terms. However, a recent meta-analysis found that eye movements make a significant contribution in processing emotional memories.⁵⁹ The results of this study were not free of controversy, as there were accusations of bad praxis in the development of the research⁶⁰ and a subsequent refutation.⁶¹

On the other hand, the debate on the degree to which the causal mechanisms of its effect differ substantially or not from those that function in trauma-centred CBT or standard exposure is sterile from a methodological standpoint. This is due to lack of knowledge and agreement on which biological mechanisms function in general in all psychotherapies. Trauma-centred CBT consists of a combination of exposure techniques and cognitive interventions, while EMDR is an eclectic psychotherapy which includes a set of structured procedures and protocols. Although many procedures seem to overlap with trauma-centred CBT, the National Institute for Care Excellence has stated that the 2 approaches are different as they require specific training programmes.³³ In a paper published in 2002 Rogers and Silver⁶² describe the differences between EMDR protocols and those involving exposure in detail, the latter being a key component of trauma-centred CBT.

The basic protocol for administering EMDR is a parsimonious and painstakingly detailed procedure, facilitating its clinical use as well as scientific scrutiny. The majority of the studies that have examined its mechanism of action have centred on studying one part of EMDR therapy, bilateral stimulation, and more specifically on eye movement. 3 hypotheses have fundamentally emerged from these studies on the possible effect of bilateral stimulation: memory resource overload, a fall in physiological arousal or improved connectivity between the hemispheres of the brain. These theoretical models, which often overlap each other, point to the directions of future research which aims for increased conceptual clarity and consistency.

The level of sophistication of studies of the mechanism of action of EMDR has increased exponentially in recent years. A study undertaken by Landin-Romero et al.⁶³ using functional neuroimaging techniques stands out. This showed for

the first time that EMDR is able to modulate patterns of neuronal net activation and deactivation by default. This brain system has chiefly been linked with autobiographical and introspective thought by an individual. Nevertheless, with very few exceptions the majority of these studies have major methodological limitations. Their results should therefore be considered to be preliminary, so that hypotheses about the underlying biological effects of EMDR are currently mainly speculative.

Finally, this work presented a systematic review of 15 randomised controlled studies which compared the efficacy of EMDR in the treatment of PTSD with that of other psychotherapeutic interventions. According to the recommendations of different international psychological and psychiatric associations, and in spite of the fact that this work lacks quantitative analysis of the results, it can be said that there are sufficient high-methodological quality studies which indicate that EMDR is at the least equally effective as a therapy as trauma-centred CBT. Additionally, some studies seem to indicate that EMDR is more effective at reducing symptoms in a shorter period of time, and that it is a therapy that is tolerated better by patients.^{5,49,51}

It should be pointed out that the first studies of EMDR were criticised because of aspects linked to the methodology used.⁶⁴ These were mainly the absence of a control group to isolate the possible placebo effect of the intervention,^{25,65} the lack of double blind evaluations,⁴⁵ exclusive use of measurements supplied by self-reports and ambiguity about the severity and diagnosis of PTSD. Clinicians who specialise in EMDR¹¹ also pointed out some factors that may have hindered the observation of positive effects of the treatment. These included the insufficient number of EMDR sessions^{25,43,66} and/or the administration of psychotherapy by professionals who lack training⁴¹ or who had not properly followed the protocol. More recent studies can be seen to have attempted to overcome these methodological problems, and the methodological quality can be seen to have improved in general, as have the criteria proposed in increasingly clinical areas for EMDR. Branches of comparison with specific techniques are now included, together with clinical interviews and standardised self-completed reports to evaluate symptoms,^{51,53} evaluators who are blind regarding the intervention conditions^{49,50} and qualified professionals with training and years of experience in the use of EMDR.⁴⁴

It is important to point out that EMDR is gaining ground in the treatment of other psychiatric diseases that are comorbid with PTSD or traumatic events, such as addictions,⁶⁷ bipolar disorder,^{37,68} unipolar depression⁶⁹ or psychosis.^{70,71} There is now preliminary scientific evidence for the technique in controlled randomised studies with promising results.

To conclude, there is a growing number of clinical studies, reviews and meta-analyses which supply data on the effectiveness and safety of EMDR in the treatment of PTSD. They place it at the same level as other treatments of choice, such as trauma-centred CBT. Nevertheless, research into EMDR is still at an early stage. Further studies are still necessary to gain greater understanding of its mechanism of action and the neurobiological foundations underlying the therapy. Finally, more clinical trials with an appropriate methodological quality are needed to strengthen the results obtained to

date on the efficacy of EMDR in the treatment of patients diagnosed PTSD, and to confirm the first positive results of using it to treat other mental illnesses which have trauma as a comorbidity.

Conflict of interests

Francisca García is the President of the Spanish EMDR Association, and Isabel Fernández is the President of the European EMDR Association. Benedikt L. Amann, Ramón Landin-Romero and Walter Lupo have been guest speakers in several national and international EMDR conferences.

Acknowledgements

This review was supported by the European and Spanish EMDR Association, the Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), several grants from the National Plan for R&D&I, the Instituto de Salud Carlos III-Subdirección General de Evaluación y Fomento de la Investigación, Plan Nacional 2008–2011 and 2013–2016, and The European Regional Development Fund (FEDER): stabilisation contract (CES 12/024) and grants FIS (PI07/1278, PI10/02622 and PI/15/02242) for Benedikt L. Amann.

References

1. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Post-traumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry*. 1995;52:1048–60.
2. Vieweg WV, Julius DA, Fernandez A, Beatty-Brooks M, Hettema JM, Pandurangi AK. Posttraumatic stress disorder: clinical features, pathophysiology, and treatment. *Am J Med*. 2006;119:383–90.
3. American Psychiatric Association. Manual diagnóstico y estadístico de los trastornos mentales, cuarta edición (DSM-IV). 4th ed. Barcelona: Masson; 1995.
4. Bisson JI, Roberts NP, Andrew M, Cooper R, Lewis C. Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *Cochrane Database Syst Rev*. 2013;12:CD003388.
5. McGuire TM, Lee CW, Drummond PD. Potential of eye movement desensitization and reprocessing therapy in the treatment of post-traumatic stress disorder. *Psychol Res Behav Manag*. 2014;7:273–83.
6. Shapiro F. Eye movement desensitization: a new treatment for post-traumatic stress disorder. *J Behav Ther Exp Psychiatry*. 1989;20:211–7.
7. Shapiro F. EMDR: in the eye of a paradigm shift. *Behavior Therapist*. 1994;17:153–7.
8. Shapiro F. Eye movement desensitization and reprocessing (EMDR): evaluation of controlled PTSD research. *J Behav Ther Exp Psychiatry*. 1996;27:209–18.
9. Shapiro F. Desensitization and reprocessing: basic principles, protocols and procedures. 2nd ed. New York: Guilford Press; 2001.
10. Shapiro F. EMDR 12 years after its introduction: past and future research. *J Clin Psychol*. 2002;58:1–22.
11. Shapiro F. Desensibilización y reprocesamiento por movimiento ocular. 2.ª ed. México: Pax México; 2005.
12. Cusack K, Spates CR. The cognitive dismantling of eye movement desensitization and reprocessing (EMDR) treatment of posttraumatic stress disorder (PTSD). *J Anxiety Disord*. 1999;13:87–99.

13. Van den Hout M, Muris P, Salemink E, Kindt M. Autobiographical memories become less vivid and emotional after eye movements. *Br J Clin Psychol.* 2001;40:121–30.
14. Davidson PR, Parker KC. Eye movement desensitization and reprocessing (EMDR): a meta-analysis. *J Consult Clin Psychol.* 2001;69:305–16.
15. Shapiro F, Maxfield L. Eye movement desensitization and reprocessing (EMDR): information processing in the treatment of trauma. *J Clin Psychol.* 2002;58:933–46.
16. Wilson DL, Silver SM, Covi WG, Foster S. Eye movement desensitization and reprocessing: effectiveness and autonomic correlates. *J Behav Ther Exp Psychiatry.* 1996;27:219–29.
17. Andrade J, Kavanagh D, Baddeley A. Eye-movements and visual imagery: a working memory approach to the treatment of post-traumatic stress disorder. *Br J Clin Psychol.* 1997;36 Pt 2:209–23.
18. Christman SD, Garvey KJ, Propper RE, Phaneuf KA. Bilateral eye movements enhance the retrieval of episodic memories. *Neuropsychology.* 2003;17:221–9.
19. Parker A, Relph S, Dagnall N. Effects of bilateral eye movements on the retrieval of item, associative, and contextual information. *Neuropsychology.* 2008;22:136–45.
20. Propper RE, Pierce J, Geisler MW, Christman SD, Bellorado N. Effect of bilateral eye movements on frontal interhemispheric gamma EEG coherence: implications for EMDR therapy. *J Nerv Ment Dis.* 2007;195:785–8.
21. Samara Z, Elzinga BM, Slagter HA, Nieuwenhuis S. Do horizontal saccadic eye movements increase interhemispheric coherence? Investigation of a hypothesized neural mechanism underlying EMDR. *Front Psychiatry.* 2011;2:4.
22. Nieuwenhuis S, Elzinga BM, Ras PH, Berends F, Duijs P, Samara Z, et al. Bilateral saccadic eye movements and tactile stimulation, but not auditory stimulation, enhance memory retrieval. *Brain Cogn.* 2013;81:52–6.
23. Stickgold R. EMDR: a putative neurobiological mechanism of action. *J Clin Psychol.* 2002;58:61–75.
24. Raboni MR, Alonso FF, Tufik S, Suchecki D. Improvement of mood and sleep alterations in posttraumatic stress disorder patients by eye movement desensitization and reprocessing. *Front Behav Neurosci.* 2014;8:209.
25. Wilson SA, Becker LA, Tinker RH. Eye movement desensitization and reprocessing (EMDR) treatment for psychologically traumatized individuals. *J Consult Clin Psychol.* 1995;63:928–37.
26. Barrowcliff ALGN, MacCulloch S, Freeman TCA, MacCulloch MJ. Eye-movements reduce the vividness, emotional valence and electrodermal arousal associated with negative autobiographical memories. *J Forens Psychiatry Psychol.* 2004;15:325–45.
27. MacCulloch MJ, Feldman P. Eye movement desensitisation treatment utilises the positive visceral element of the investigatory reflex to inhibit the memories of post-traumatic stress disorder: a theoretical analysis. *Br. J. Psychiatry.* 1996;169:571–9.
28. Sack M, Lempa W, Steinmetz A, Lamprecht F, Hofmann A. Alterations in autonomic tone during trauma exposure using eye movement desensitization and reprocessing (EMDR)—results of a preliminary investigation. *J Anxiety Disord.* 2008;22:1264–71.
29. Gunter RW, Bodner GE. How eye movements affect unpleasant memories: support for a working-memory account. *Behav Res Ther.* 2008;46:913–31.
30. Smeets MA, Dijks MW, Pervan I, Engelhard IM, van den Hout MA. Time-course of eye movement-related decrease in vividness and emotionality of unpleasant autobiographical memories. *Memory.* 2012;20:346–57.
31. Practice guidelines for the treatment of patients with acute stress disorder and posttraumatic stress disorder. Arlington, VA: American Psychiatric Association; 2004.
32. APA Presidential Task Force on Evidence-Based Practice. Evidence-based practice in psychology. *Am Psychol.* 2006;61:271–85.
33. National Institute for Clinical Excellence. Posttraumatic stress disorder (PTSD): the management of adults and children in primary and secondary care. London: NHS; 2005.
34. World Health Organization. Guidelines for the management of conditions specifically related to stress. Geneva: WHO; 2013.
35. Seidler GH, Wagner FE. Comparing the efficacy of EMDR and trauma-focused cognitive-behavioral therapy in the treatment of PTSD: a meta-analytic study. *Psychol Med.* 2006;36:1515–22.
36. Chen YR, Hung KW, Tsai JC, Chu H, Chung MH, Chen SR, et al. Efficacy of eye-movement desensitization and reprocessing for patients with posttraumatic-stress disorder: a meta-analysis of randomized controlled trials. *PLOS ONE.* 2014;9:e103676.
37. Amann BL, Batalla R, Blanch V, Capellades D, Carvajal MJ, Fernández I, et al. The EMDR therapy protocol for bipolar disorder. In: Luber M, editor. Eye movement desensitization and reprocessing (EMDR) scripted protocols and summary sheets: treating trauma, anxiety and mood-related conditions. New York: Springer; 2015. p. 223–87.
38. Jadad AR, Enkin M, editors. Randomized controlled trials: questions, answers and musings. Oxford: Blackwell; 2007.
39. Urrútia G, Bonfill X. [PRISMA declaration: a proposal to improve the publication of systematic reviews and meta-analyses] Spanish. *Med Clin (Barc).* 2010;135:507–11.
40. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA Statement. *Open Med.* 2009;3:e123–30.
41. Rothbaum BO. A controlled study of eye movement desensitization and reprocessing in the treatment of posttraumatic stress disorder sexual assault victims. *Bull Menninger Clin.* 1997;61:317–34.
42. Marcus SV, Marquis P, Sakai C. Controlled study of treatment of PTSD using EMDR in an HMO setting. *Psychother Theor Res Pract Train.* 1997;34:307–15.
43. Scheck MM, Schaeffer JA, Gillette C. Brief psychological intervention with traumatized young women: the efficacy of eye movement desensitization and reprocessing. *J Trauma Stress.* 1998;11:25–44.
44. Hogberg G, Pagani M, Sundin O, Soares J, Aberg-Wistedt A, Tärnell B, et al. On treatment with eye movement desensitization and reprocessing of chronic post-traumatic stress disorder in public transportation workers—a randomized controlled trial. *Nord J Psychiatry.* 2007;61:54–61.
45. Vaughan K, Armstrong MS, Gold R, O'Connor N, Jenneke W, Tarrier N. A trial of eye movement desensitization compared to image habituation training and applied muscle relaxation in post-traumatic stress disorder. *J Behav Ther Exp Psychiatry.* 1994;25:283–91.
46. Carlson JG, Chemtob CM, Rusnak K, Hedlund NL, Muraoka MY. Eye movement desensitization and reprocessing (EDMR) treatment for combat-related posttraumatic stress disorder. *J Trauma Stress.* 1998;11:3–24.
47. Devilly GJ, Spence SH. The relative efficacy and treatment distress of EMDR and a cognitive-behavior trauma treatment protocol in the amelioration of posttraumatic stress disorder. *J Anxiety Disord.* 1999;13:131–57.
48. Power KG, McGoldrick T, Brown K, Buchanan R, Sharp D, Swanson V, et al. A controlled comparison of eye movement desensitisation and reprocessing versus exposure plus cognitive restructuring versus waiting list in the treatment of posttraumatic stress disorder. *J Psychol Psychother.* 2002;9:299–318.
49. Ironson G, Freund B, Strauss JL, Williams J. Comparison of two treatments for traumatic stress: a community-based study of EMDR and prolonged exposure. *J Clin Psychol.* 2002;58:113–28.
50. Lee C, Gavriel H, Drummond P, Richards J, Greenwald R. Treatment of PTSD: stress inoculation training with prolonged exposure compared to EMDR. *J Clin Psychol.* 2002;58:1071–89.
51. Taylor S, Thordarson DS, Maxfield L, Fedoroff IC, Lovell K, Ogradniczuk J. Comparative efficacy, speed, and adverse

- effects of three PTSD treatments: exposure therapy, EMDR, and relaxation training. *J Consult Clin Psychol.* 2003;71:330–8.
52. Rothbaum BO, Astin MC, Marsteller F. Prolonged exposure versus eye movement desensitization and reprocessing (EMDR) for PTSD rape victims. *J Trauma Stress.* 2005;18:607–16.
 53. Van der Kolk BA, Spinazzola J, Blaustein ME, Hopper JW, Hopper EK, Korn DL, et al. A randomized clinical trial of eye movement desensitization and reprocessing (EMDR), fluoxetine, and pill placebo in the treatment of posttraumatic stress disorder: treatment effects and long-term maintenance. *J Clin Psychiatry.* 2007;68:37–46.
 54. Karatzias T, Power K, Brown K, McGoldrick T, Begum M, Young J, et al. A controlled comparison of the effectiveness and efficiency of two psychological therapies for posttraumatic stress disorder: eye movement desensitization and reprocessing vs. emotional freedom techniques. *J Nerv Ment Dis.* 2011;199:372–8.
 55. Nijdam MJ, Gersons BP, Reitsma JB, de Jongh A, Olff M. Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: randomised controlled trial. *Br J Psychiatry.* 2012;200:224–31.
 56. Kavanagh DJ, Freese S, Andrade J, May J. Effects of visuospatial tasks on desensitization to emotive memories. *Br J Clin Psychol.* 2001;40:267–80.
 57. Lee CW, Drummond PD. Effects of eye movement versus therapist instructions on the processing of distressing memories. *J Anxiety Disord.* 2008;22:801–8.
 58. Cahill SP, Carrigan MH, Frueh BC. Does EMDR work? And if so, why?: A critical review of controlled outcome and dismantling research. *J Anxiety Disord.* 1999;13:5–33.
 59. Lee CW, Cuijpers P. A meta-analysis of the contribution of eye movements in processing emotional memories. *J Behav Ther Exp Psychiatry.* 2013;44:231–9.
 60. Devilly GJ, Ono M, Lohr JM. The use of meta-analytic software to derive hypotheses for EMDR. *J Behav Ther Exp Psychiatry.* 2014;45:223–5.
 61. Lee CW, Cuijpers P. What does the data say about the importance of eye movement in EMDR? *J Behav Ther Exp Psychiatry.* 2014;45:226–8.
 62. Rogers S, Silver SM. Is EMDR an exposure therapy? A review of trauma protocols. *J Clin Psychol.* 2002;58:43–59.
 63. Landin-Romero R, Novo P, Vicens V, McKenna PJ, Santed A, Pomarol-Clotet E. EMDR therapy modulates the default mode network in a subsyndromal, traumatized bipolar patient. *Neuropsychobiology.* 2013;67:181–4.
 64. Lohr JM, Kleinknecht RA, Tolin DF, Barrett RH. The empirical status of the clinical application of eye movement desensitization and reprocessing. *J Behav Ther Exp Psychiatry.* 1995;26:285–302.
 65. Grainger RD, Levin C, Allen-Byrd L, Doctor RM, Lee H. An empirical evaluation of eye movement desensitization and reprocessing (EMDR) with survivors of a natural disaster. *J Trauma Stress.* 1997;10:665–71.
 66. Lytle RA, Hazlett-Stevens H, Borkovec TD. Efficacy of eye movement desensitization in the treatment of cognitive intrusions related to a past stressful event. *J Anxiety Disord.* 2002;16:273–88.
 67. Perez-Dandieu B, Tapia G. Treating trauma in addiction with EMDR: a pilot study. *J Psychoactive Drugs.* 2014;46:303–9.
 68. Novo P, Landin-Romero R, Rada J, Vicens V, Fernandez I, Garcia F, et al. Eye movement desensitization and reprocessing therapy in subsyndromal bipolar patients with a history of traumatic events: a randomized, controlled pilot-study. *Psychiatry Res.* 2014;219:122–8.
 69. Hase M, Balmaceda UM, Hase A, Lehnung M, Tumani V, Huchzermeyer C, et al. Eye movement desensitization and reprocessing (EMDR) therapy in the treatment of depression: a matched pairs study in an inpatient setting. *Brain Behav.* 2015;5:e00342.
 70. Van den Berg DP, De Bont PA, Van der Vleugel BM, De Roos C, De Jongh A, Van Minnen A, et al. Prolonged exposure vs eye movement desensitization and reprocessing vs waiting list for posttraumatic stress disorder in patients with a psychotic disorder: a randomized clinical trial. *JAMA Psychiatry.* 2015;72:259–67.
 71. Van den Berg DP, De Bont PA, Van der Vleugel BM, De Roos C, De Jongh A, Van Minnen A, et al. Trauma-focused treatment in PTSD patients with psychosis: symptom exacerbation, adverse events, and revictimization. *Schizophr Bull.* 2015, pii: sbv172.
 72. Hogberg G, Pagani M, Sundin O, Soares J, Aberg-Wistedt A, Tärnell B, et al. Treatment of post-traumatic stress disorder with eye movement desensitization and reprocessing: outcome is stable in 35-month follow-up. *Psychiatry Res.* 2008;159:101–8.