

Original Article

Laparoscopic Left Lateral Sectionectomy. Presentation of Our Technique[☆]

E. Herrero Fonollosa,^{a,*} E. Cugat Andorra,^a M.I. García-Domingo,^a J. Rivero Deniz,^a
J. Camps Lasa,^a A. Rodríguez Campos,^b M. Riveros Caballero,^a C. Marco Molina^a

^aUnidad de Cirugía HBP, Servicio de Cirugía, General Hospital Universitari Mútua de Terrassa, Terrassa, Barcelona, Spain

^bServicio de Anestesia y Reanimación, Hospital Universitari Mútua de Terrassa, Terrassa, Barcelona, Spain

ARTICLE INFO

Article history:

Received 11 January 2011

Accepted 19 June 2011

Available online 20 January 2012

Keywords:

Hepatectomy

Laparoscopy

Sectionectomy

A B S T R A C T

Introduction: Laparoscopy has not been as widely used in hepatic surgery as in other areas due to the complexity of this type of surgery and the lack of surgical teams with experience in both fields.

Material and methods: The aim of this work is to present the technique used in our centre to perform left lateral sectionectomy (LLS) using laparoscopy. A total of 70 patients have been operated on using laparoscopy due to both benign and malignant liver between February 2000 and July 2010. An LLS was performed on twenty-one cases using the technique described. The surgical technique is described, highlighting aspects such as, the arrangement of the trocars, the mobilisation of the liver or hepatic transection. The morbidity and mortality associated with the procedure are analysed.

Results: LLS was performed on 12 women and nine men, with ages between 35 and 89 years. The mean number of lesions was 1.4 (between 1 and 4), with a mean size of 3.5 cm. The mean surgical time was 142 min (between 90 and 210). There was one conversion to laparotomy. Complications were recorded in 3 (14%) patients. There were no repeat surgery, and one patient required a transfusion. The mean hospital stay was 4.3 days.

Conclusions: The best techniques and the wide experience in laparoscopy have enabled this technique to become established, with a morbidity of less than 15% and zero mortality. LLS is a safe and effective technique in selected patients. The detailed description of this procedure may stimulate other surgery groups to perform this approach.

© 2011 AEC. Published by Elsevier España, S.L. All rights reserved.

Seccionectomía lateral izquierda por laparoscopia. Presentación de nuestra técnica

R E S U M E N

Introducción: En cirugía hepática, la laparoscopia no ha conseguido la difusión obtenida en otras áreas debido a la complejidad de este tipo de cirugía y a la falta de equipos quirúrgicos con experiencia en ambos campos.

Palabras clave:

Hepatectomía

[☆] Please, cite this article as: Herrero Fonollosa E, et al. Seccionectomía lateral izquierda por laparoscopia. Presentación de nuestra técnica. Cir Esp. 2011;89:650-6.

* Corresponding author.

E-mail address: eherrero@mutuaterrassa.es (E. Herrero Fonollosa).

2173-5077/\$ – see front matter © 2011 AEC. Published by Elsevier España, S.L. All rights reserved.

Laparoscopia Seccionectomía

Material y métodos: El objetivo de este trabajo es presentar la técnica utilizada en nuestro centro para realizar la seccionectomía lateral izquierda por laparoscopia (SLI). Entre febrero de 2000 y julio de 2010, 70 pacientes han sido intervenidos por laparoscopia por patología hepática, tanto benigna como maligna. En veintidós casos se realizó una SLI según la técnica descrita. Se describe la técnica quirúrgica, destacando aspectos como la disposición de los trocares, la movilización del hígado o la transección hepática. Se analiza la morbilidad relacionada con el procedimiento.

Resultados: Se ha realizado la SLI en 12 mujeres y 9 hombres con edades comprendidas entre los 35 y los 89 años. El número de lesiones fue de 1,4 (entre 1 y 4), con un tamaño de 3,5 cm. El tiempo operatorio fue de 142 min (entre 90 y 210). Hubo una conversión a laparotomía. Se registraron complicaciones en 3 pacientes (14%). No hubo reintervenciones y un paciente requirió una transfusión. La estancia media hospitalaria fue de 4,3 días.

Conclusiones: Las mejoras técnicas y la mayor experiencia en laparoscopia han permitido plantear la realización de este procedimiento con una morbilidad inferior al 15% y una mortalidad nula. La SLI es una técnica segura y efectiva en pacientes seleccionados. La descripción detallada de este procedimiento puede estimular a otros grupos de cirugía hepática a realizar este abordaje.

© 2011 AEC. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

The success obtained by laparoscopic abdominal surgery is unquestionable. However, hepatic surgery has not seen a widespread use of laparoscopic techniques, mostly due to the complexity inherent in these procedures and the lack of surgical teams specialised in both fields (hepatic and laparoscopic surgery).¹⁻⁵

The first laparoscopic anatomical liver resection was performed in 1996, and in Spain the first procedure was performed in 2000.⁶ From its onset, the majority of authors have coincided in defining certain favourable liver segments that are more accessible for a laparoscopic approach, these being the left lateral and right anterior segments.⁷⁻⁹

The anatomy of the left lateral sector and the configuration of the portal and supra-hepatic pedicles facilitate a feasible and safe laparoscopic approach in this area, with oncological results similar to those achieved using open surgery when performing R0 resections with tumour-free margins. The laparoscopic approach for lesions on the left lateral segment should be considered the method of choice for experienced centres,¹⁰⁻¹² with superior results in terms of blood losses, operational time, and hospital stay, when compared to the laparotomy approach.¹³⁻¹⁹

The aim of our study was to describe the surgical technique used for a left lateral sectionectomy (LLS) as performed by our surgical team, and to analyse the morbidity and mortality rates associated with this procedure.

Patients and Method

Patients

In 2000, our institution started performing laparoscopic liver surgery. The first cases were for benign pathologies, but indications were later expanded to include laparoscopic

resections for malignant ones. A total of 70 procedures have been performed using laparoscopic techniques for hepatic pathologies: 17 cases of cystic pathologies (6 simple cysts, 4 hydatid cysts, 6 cases of polycystic liver disease, and 1 cystadenoma), 10 cases of solid benign tumours (8 adenomas, 1 sclerosing cholangitis, and 1 haemangioma), and 43 cases of malignant tumours (21 metastases from colorectal cancer, 10 hepatocellular carcinomas [HCC], 5 metastases from breast cancer, 2 metastases from lung cancer, 2 cholangiocarcinomas, 1 lymphoma, 1 metastasis from a haemangioendothelioma, and 1 metastasis from a melanoma). Since 2005, laparoscopic resection of lesions in segments II and III has been well established.

Surgical Technique

Our surgical protocol involved balanced general anaesthesia, using propofol for induction and sevoflurane and remifentanyl on continuous perfusion as maintenance therapy. Blood pressure and central venous pressure were monitored throughout the procedure. During the operation, we also administered the necessary liquids to maintain diuresis at 0.5 mg/kg/h, without exceeding a central venous pressure (CVP) of 6 mm Hg, administering the necessary vasoactive drugs according to patient tolerance to clamping and unclamping of the hepatic hilum.

The patient was placed in the supine decubitus position with the legs spread ("French position"). The operating table was placed in the anti-Trendelenburg position. We used six trocars (three 10-12 mm and three 5 mm). Pneumoperitoneum was achieved using a Veress needle at the umbilical level. We used a pressure of 10-12 mm Hg, and CO₂ for the pneumoperitoneum.

We used a 30° optic placed at the umbilical level. The 10-12 mm working trocars were placed on each side of the median line, approximately 5 cm above the navel. We then placed a 5 mm trocar in the left hypochondrium and another in a very lateral position on the right hypochondrium, which was then used for the Pringle manoeuvre.




Fig. 1 – Dissection of the suprahepatic veins.




Fig. 2 – Intraoperative ultrasound.

Finally, we placed the last 5 mm trocar in the epigastric region for the traction of the round ligament of the liver during the transection. The surgeon was situated between the patient's legs, the first assistant to the left of the patient, and the second assistant to the right.

The abdominal cavity was then examined to rule out the presence of widespread disease in the case of malignant pathology.

We proceeded with a dissection of the round ligament using Ligasure[®] V (Valleylab, Tyco Healthcare) or Ultracision[®] (Ethicon EndoSurgery, Johnson & Johnson Ltd., Cincinnati, OH, USA), cutting as close as possible to the abdominal wall for traction during the transection and to prevent it from obstructing the view. Then we dissected the falciform ligament until nearing the suprahepatic veins, since dissection of these vessels is favoured by the pneumoperitoneum (Fig. 1). This manoeuvre facilitates quick access to the suprahepatic veins for clamping, should it be necessary.

We then made an incision in the left triangular ligament, taking special care to avoid the left diaphragmatic vein, which passes very close to the ligament. Damage to this vein produces bleeding similar to the suprahepatic vein, which is very difficult to control and causes the risk of gas embolism. Complete mobilisation of segments II and III is essential since this allows for separating them from the diaphragm, facilitating the hepatic transection and the placement of endostaples at the proper angle.

The use of intraoperative ultrasound (IOUS) (Fig. 2) is a standard procedure of this surgery since it allows us to: (a) rule out lesions that were not diagnosed before the procedure; (b) ensure a transection line with proper margins, and (c) diagnose variants of the primary pedicles in order to prevent haemorrhage during the transection.

The traction of the round ligament and the elevation of segment IV using a clamp allow for the exposure of the hepatic hilum and pars flaccida of the lesser omentum to be dissected. Using a clamp through the trocar situated on the lateral side of the right hypochondrium, a 10 cm non-elastic band is passed under the hilum, through the foramen of Winslow, and out the opening we have made in the lesser omentum, encircling

the hepatic hilum. The two ends are passed through a 2 cm-long silastic tube that will serve to clamp the hilum using grasping forceps situated in the right lateral trocar. If the patient had undergone a previous operation on the hepatic hilum, passing the band through the foramen of Winslow can be impeded, and the surgeon can opt for the use of a laparoscopic clamp on the anterior face of the hepatic hilum.

It is of paramount importance during the hepatic transection to maintain CVP as low as possible, since this minimises blood loss. Due to the effect of the pneumoperitoneum on the CVP transducer, readings are not completely reliable, and so it is best to take these measurements before achieving pneumoperitoneum.

Several different devices can be used for the transection of normal hepatic parenchyma. The superficial part of the hepatic parenchyma has little vascularisation, facilitating the use of a monopolar scalpel down to a depth of 0.5–1 cm. From this point, the underlying vessels are dissected using a laparoscopic ultrasonic dissector, with coagulation achieved using 5 mm Ligasure Atlas[®] or Ultracision[®] combined with laparoscopic Tissuelink[®] (Salient Surgical Technologies). Tissuelink[®] provides superficial coagulation of the hepatic parenchyma and small vessels, while the Ligasure Atlas[®] and Ultracision[®] coagulate and divide the major vessels and parenchyma.

The transection is performed under continuous occlusion of the hepatic pedicle (Pringle manoeuvre). During the transection, the round ligament of the liver is under traction to the right using a clamp through the epigastric trocar, a line is marked for transection on the surface of the hepatic parenchyma by means of Tissuelink[®] surface coagulation, and then the transection of the superficial parenchyma performed using the two previously described devices until reaching the area of the main portal pedicles of the segment. Generally, one staple is needed for the segment III portal pedicle, another for the segment II portal pedicle (Fig. 3), and another for the sectioning of the left suprahepatic vein (Fig. 4).

With the goal of ensuring haemostasis and bile stasis, we applied a sponge made of a haemostatic-sealant material of




Fig. 3 – Division of the portal branches of segments II and III with an endostapler.




Fig. 4 – Division of the left suprahepatic vein with an endostapler.

collagen with fibrin and thrombin (Tachosil[®], Nycomed) over the surface of the hepatic transection.

The whole surgical specimen is extracted through an accessory incision, within a bag. If the tumour was small, usually one of the 10–12 mm trocar holes must be widened by 3–4 cm. In patients with larger tumours, we used a Pfannenstiel incision. A suction drain is placed beneath the resection bed through the 5 mm trocar in the right flank. The accessory incision and 10–12 mm trocars are then closed using absorbable sutures.

Results

Between May 2005 and July 2010, 21 patients were operated upon using the procedure described here (Table 1). Nine of these patients were men and 12 were women, with ages ranging between 35 and 89 years.

The surgery was for a malignant pathology in 17 patients (9 colorectal cancer metastases, 4 HCC, 2 breast cancer metastases, 1 melanoma metastasis, and 1 primary lung

metastasis), and for benign pathologies in 4 cases (3 adenomas and 1 case of sclerosing cholangitis) (Table 1).

The mean number of lesions found per patient was 1.4 (range: 1–4), with a mean size of 3.8 cm (range: 1–5 cm). The mean duration of the procedure was 142 min (range: 90–120 min). In most cases (76%), we used a Pringle manoeuvre during the hepatic transection, with a mean duration of 26 min (range: 10–60 min).

We had to switch to a laparotomy in one case in order to complete the resection of a lesion that went undiagnosed prior intraoperative ultrasound examinations. Postoperative complications were produced in 3 patients (14%): one infection of the surgical wound (grade I on the Dindo-Clavien scale), one pulmonary thromboembolism in a patient operated on for a metastasized (M1) melanoma, with little clinical repercussions (grade II), and one intra-abdominal collection in the resection bed that required a percutaneous drain (grade III). There were no second operations for any patients. We provided intraoperative blood transfusions in one patient with M1 breast cancer that had intraoperative bleeding from a lesion on an epiploic vessel that was controlled using laparoscopy.

The patients started oral tolerance at a mean 30 h after the operation (range: 16–72 h). The duration of analgesic requirements was 2.6 days (range: 2–4 days). The mean hospital stay was 4.3 days (range: 2–7 days). The histological analysis of the resection margins was negative in all cases (Table 2).

Table 1 – Aetiology of the Laparoscopic LLS Performed at the Hospital Mútua de Terrassa (n=21). Time period: 2005–2010.

	No.	%
CRCHM	9	43%
HCC	4	19%
Non-CRCHM	4	19%
Breast cancer	2	50%
Melanoma	1	25%
Lung cancer	1	25%
Adenoma	3	14%
Sclerosing cholangitis	1	4%

HCC: hepatocellular carcinoma; CRCHM: colorectal cancer hepatic metastasis; non-CRCHM: non-colorectal cancer hepatic metastasis.

Discussion

Hepatic resection of segments II and III is a feasible and safe technique. The current results reproduce and even improve upon those from open surgery in selected cases, with a mortality rate of <5% and morbidity of <20%.^{20–24}

LLS should be considered as the technique of choice in benign or malignant lesions located on segments II and III in patients that do not have severe comorbidities that would contraindicate laparoscopy. A patient history involving

Table 2 – Results from Laparoscopic LLS Performed at the Hospital Mútua de Terrassa (n=21). Time period: 2005–2010.

	No., %	Mean ± Standard Deviation
Sex (M/F)	9/12	
Age, years		56±16.90
Number of lesions		1.4±0.75
Lesion size, cm		3.8±2.30
Operational time, min		142±53.88
Clamping time, min		25.3±8.07
Preoperative haemoglobin, mg/dl		13.36±1.77
Postoperative haemoglobin, mg/dl		11.79±1.81
Transfusion	1/21 (4)	
Complications	3/21 (14)	
Oral food tolerance, h		30.8±16.61
Duration of analgesia, days		2.6±2.25
Duration of hospital stay, days		4.3±3.07

previous open surgical procedures is not uncommon in our field (laparotomy for colon resection), but this does not contraindicate the laparoscopic approach, although it may increase the technical difficulty of the procedure due to adhesences.

A few groups support the use of hand-assisted laparoscopic surgery in the case of large lesions,^{25–27} but since the transection line to the left of the round and falciform ligaments of the liver is always the same, we believe that the size of the tumour alone is not an exclusion criteria for performing a completely laparoscopic procedure, if it is possible to achieve adequate mobilisation of the liver.

The trocars used in this procedure should be placed following a concave line with respect to the lesion, placing one 5 mm trocar on the right flank in the most posterior and superior position possible, in order to facilitate the Pringle manoeuvre with a better angle for access to the foramen of Winslow.

A complete mobilisation of the left hepatic lobe facilitates transection and allows for proper access to the suprahepatic veins. The dissection of the anterior face of the suprahepatic veins allows for directly clamping these vessels in the case of haemorrhage, and avoids damaging the left diaphragmatic vein when separating the liver from the diaphragm. Better control of the intrahepatic left suprahepatic vein is best when the left liver is mobilised, since the vein is then in a perpendicular position with respect to the endostapler.

Laparoscopic intraoperative ultrasound must always be used in order to rule out lesions that may have gone unnoticed during the preoperative analysis. This point is of vital importance when dealing with a malignancies, especially colorectal cancer hepatic metastases (CRCHM), since in 20% of cases, IOUS diagnoses hepatic lesions that were not identified using other diagnostic techniques.^{28–30} This occurred in one of the patients from our study, in which the IOUS showed a deep metastatic lesion in segment V that had gone undiagnosed, requiring conversion to a laparotomy in order to safely complete the oncological resection.

Clamping the hepatic hilum during transection is a highly debated issue. Several different possible postoperative

complications are associated with prolonged clamping (longer than 1 h).^{2,3} In our experience, we have not registered any postoperative complications attributable to the Pringle manoeuvre. Additionally, continuous clamping does not tend to last for more than 25 minutes, and is usually well tolerated by most patients both during the procedure and postoperatively.

Using ischaemic preconditioning of the area with a 10-min clamping and 10-min unclamping could provide a beneficial effect on liver cells through various mechanisms, including the inhibition of apoptosis and the local release of proinflammatory cytokine antagonists, protecting the liver from later ischaemia-reperfusion damage.^{31–33}

In spite of the fact that several groups do not support a systematic clamping of the hilum in laparoscopic LLS,³⁴ we used hilar clamping with ischaemic preconditioning as a standard practice because: (a) no secondary side effects were observed in our study after continuous clamping, both in laparoscopic and open surgeries; (b) this provides as a blood-free operative field as possible (the presence of blood in the surgical area absorbs light, darkening the image and making it difficult to identify key structures during the hepatic transection); and (c) blood losses and consequent transfusion in patients that undergo hepatic resection due to metastatic colon cancer can be a prognostic factor against long-term survival.^{35–37} In our study, one patient required a blood transfusion due to damage to a gastroepiploic vessel during the mobilisation manoeuvre, causing severe haemorrhage due to the increased splenic pressure caused by the hilar clamping, although this was controlled without the need for conversion to open surgery.

We find it best to work with a low CO₂ pressure in the pneumoperitoneum, around 10–12 mm Hg, in order to minimise the possibility of gas entering the veins in the case of damage to a suprahepatic vein. Also, in contrast to the standard procedures used for open surgery, i.e. a central venous pressure close to 0 mm Hg in order to minimise haemorrhage, it is preferable to maintain this pressure close to 6 mm Hg in order to decrease the gradient between the intra-abdominal pressure and CVP, thus reducing the risk of gas embolism.^{3,38–40}

For the treatment of hepatic malignancies, LLS is indicated in selected cases as long as the principles of oncological resection are respected, with adequate tumour-free margins. Doubts exist regarding the risk of facilitating intra-abdominal dissemination of the tumour and metastases in the trocar access ports during laparoscopic resections for malignancies.^{41–44} The results obtained in studies focusing on colon cancer appear to contradict this theory. In our cohort, all cases of malignancies were resected with tumour-free margins greater than 1 cm, and to this date we have not registered any cases of recurrence in the abdominal wall in any patients.

Laparoscopic resections of HCC located in the left lateral sector of the liver offer several advantages over open surgery. It is a less invasive approach that preserves collateral circulation of the abdominal wall and minimises the possibility of ascitic decompensation in the postoperative period. For this reason, the surgical indications for this procedure could possibly be expanded to include selected patients with hepatic cirrhosis and portal hypertension.^{45–48}

In conclusion, laparoscopic LLS is a feasible and reproducible technique. The anatomical positioning of segments II and III and their portal and suprahepatic pedicles facilitates a laparoscopic approach. Additionally, this technique has been proven effective and safe for the treatment of both benign and malignant pathologies in selected patients.⁴⁹ Finally, standardising this procedure could facilitate widespread acceptance of this method for the majority of liver surgeons with laparoscopic experience.

Conflicts of Interest

The authors have no conflicts of interest to declare.

REFERENCES

- Descottes B, Lachachi F, Sodji M, Valleix D, Durand-Fontanier S, Pech de Laclause B, et al. Early experience with laparoscopic approach for solid liver tumors: initial 16 cases. *Ann Surg.* 2000;232:641-5.
- Cherqui D. Laparoscopic liver resection. *Br J Surg.* 2003;90:644-6.
- Cherqui D, Husson E, Hammoud R, Malassagne B, Stephan F, Bensaid S, et al. Laparoscopic liver resections: a feasibility study in 30 patients. *Ann Surg.* 2000;232:753-62.
- Buell JF, Koffron AJ, Thomas MJ, Rudich S, Abecassis M, Woodle ES. Laparoscopic liver resection. *J Am Coll Surg.* 2005;200:472-80.
- Katkhouda N, Hurwitz M, Gugenheim J, Mavor E, Mason RJ, Waldrep DJ, et al. Laparoscopic management of benign solid and cystic lesions of the liver. *Ann Surg.* 1999;229:460-6.
- Cugat E, Perez N, Rotellar F, Suarez M, Gastaca M, Artigas V, et al. Laparoscopic liver surgery: 8 years of multicenter Spanish register. *J Hepatobiliary Pancreat Surg.* 2010;17:262-8.
- Gagner M, Rheault M, Duluq J. Laparoscopic partial hepatectomy for liver tumor. *Surg Endosc.* 1992;6:99.
- Azagra JS, Gowergen M, Gilbert E, Jacobs D. Laparoscopic anatomical left lateral Segmentectomy—Technical aspect. *Surg Endosc.* 1996;10:758-61.
- Cugat E, García Domingo MI, Bretcha P, Rodríguez A, Marco C. Laparoscopia y cirugía hepática: técnica e indicaciones. *Cir Esp.* 2004;75:23-8.
- Lesurtel M, Cherqui D, Laurent A, Tayar C, Fagniez PL. Laparoscopic versus open left lateral hepatic lobectomy: a case-control study. *J Am Coll Surg.* 2003;196:236-42.
- Chang S, Laurent A, Tayar C, Karoui M, Cherqui D. Laparoscopy as a routine approach for left lateral sectionectomy. *Br J Surg.* 2007;94:58-63.
- Robles R, Marín C, López A, Abellán B, Pastor P, Parrilla P. La resección laparoscópica de los segmentos del lóbulo izquierdo debe ser realizada por laparoscopia en centros con experiencia. *Cir Esp.* 2009;85:214-21.
- Simillis C, Constantinides VA, Tekkis PP, Darzi A, Lovegrove R, Jiao L, et al. Laparoscopic versus open hepatic resections for benign and malignant neoplasms—a meta-analysis. *Surgery.* 2007;141:203-11.
- Morino M, Morra I, Rosso E, Miglietta C, Garrone C. Laparoscopic vs open hepatic resection: a comparative study. *Surg Endosc.* 2003;17:1914-8.
- Rau HG, Buttler E, Meyer G, Schardey HM, Schildberg FW. Laparoscopic liver resection compared with conventional partial hepatectomy—a prospective analysis. *Hepatogastroenterology.* 1998;45:2333-8.
- Mala T, Edwin B, Gladhaug I, Fosse E, Soreide O, Bergan A, et al. A comparative study of the short-term outcome following open and laparoscopic liver resection of colorectal metastases. *Surg Endosc.* 2002;16:1059-63.
- Farges O, Jagot P, Kirstetter P, Marty J, Belghiti J. Prospective assessment of the safety and benefit of laparoscopic liver resections. *J Hepatobiliary Pancreat Surg.* 2002;9:242-8.
- Koffron AJ, Auffenberg G, Kung R, Abecassis M. Evaluation of 300 minimally invasive liver resections at a single institution: less is more [discussion 392-94]. *Ann Surg.* 2007;246:385-92.
- Gigot JF, Glineur D, Santiago Azagra J, Goergen M, Ceuterick M, Morino M, et al. Laparoscopic liver resection for malignant liver tumors: preliminary results of a multicenter European study. *Ann Surg.* 2002;236:90-7.
- Felsher J, Brody F. Laparoscopic resection of nodular hyperplasia. *Surg Laparosc Endosc Percutan Tech.* 2003;13:276-9.
- Mala T, Edwin B, Rosseland AR, Gladhaug I, Fosse E, Mathisen O. Laparoscopic liver resection: experience of 53 procedures at a single center. *J Hepatobiliary Pancreat Surg.* 2005;12:298-303.
- Farges O, Jagot P, Kirstetter P, Marty J, Belghiti J. Prospective assessment of the safety and benefit of laparoscopic liver resections. *Hepatobiliary Pancreat Surg.* 2002;9:242-8.
- Dulucq JL, Wintringer P, Stabilini C, Berticelli J, Mahajna A. Laparoscopic liver resections: a single center experience. *Surg Endosc.* 2005;19:886-9.
- Tri Nguyen K, Clark Gamblin C, Geller DA. World review of laparoscopic liver resection—2804 patients. *Ann Surg.* 2009;250.
- Robles R, Marín C, Abellán B, López A, Pastor P, Parrilla P. A new approach to hand-assisted laparoscopic liver surgery. *Surg Endosc.* 2008;22:2357-64.
- Fong Y, Jarnagin W, Conlon KC, DeMatteo R, Dougherty E, Blumgart LH. Hand-assisted laparoscopic liver resection: lessons from an initial experience. *Arch Surg.* 2000;135:854-9.
- Huang M, Lee W, Wang W, Wei P, Chen R. Hand-assisted laparoscopic hepatectomy for solid tumor in the posterior portion of the right lobe: initial experience. *Ann Surg.* 2003;238:674-9.
- Vogl TJ, Schwarz W, Blume S, Pietsch M, Shamsi K, Franz M, et al. Preoperative evaluation of malignant liver tumors: comparison of unenhanced and SPIO (Resovist)-enhanced MR imaging with biphasic CTAP and intraoperative US. *Eur Radiol.* 2003;13:262-72.
- Conlon R, Jacobs M, Dasgupta D, Lodge JP. The value of intraoperative ultrasound during hepatic resection compared with improved preoperative magnetic resonance imaging. *Eur J Ultrasound.* 2003;16:211-6.
- Gaitini D, Kopelman D, Soudak M, Epelman M, Assalia A, Hashmonai M, et al. Impact of intraoperative sonography on resection and cryoablation of liver tumors. *J Clin Ultrasound.* 2001;29:256-72.
- Petrowsky H, McCormack L, Trujillo M, Selzner M, Jochum W, Clavien PA. A prospective, randomized, controlled trial comparing intermittent portal triad clamping versus ischemic preconditioning with continuous clamping for major liver resection. *Ann Surg.* 2006;244:921-30.
- Rudiger HA, Kang KJ, Sindram D, Riehle H, Clavien P. Comparison of ischemic preconditioning and intermittent and continuous inflow occlusion in the murine liver. *Ann Surg.* 2002;235:400-7.
- Clavien PA, Selzner M, Rudiger HA, Graf R, Kadry Z, Rausson V, et al. A prospective randomized study in 100 consecutive patients undergoing major liver resection with versus

- without ischemic preconditioning [discussion 851-2]. *Ann Surg.* 2003;238:843-50.
34. Troisi R, Van Huysse J, Berrevoet F, Vandenbossche B, Sainz-Barriga M, Vinci A. Evolution of laparoscopic left lateral sectionectomy without the Pringle maneuver: trough resection of benign and malignant tumors to living liver donation. *Surg Endosc.* 2011;25:79-87.
35. Kooby DA, Stockman J, Ben-Porat L, Gonen M, Jarnagin W, Dematteo R, et al. Influence of transfusions on perioperative and long-term outcome in patients following hepatic resection for colorectal metastases [discusión 869-70]. *Ann Surg.* 2003;237:860-9.
36. Tsao JI, Loftus JP, Nagorney DM, Adson MA, Ilstrup DM. Trends in morbidity and mortality of hepatic resection for malignancy: a matched comparative analysis. *Ann Surg.* 1994;220:199-205.
37. Yamamoto J, Kosuge T, Takayama T, Shimada K, Yamasaki S, Ozaki H, et al. Perioperative blood transfusion promotes recurrence of hepatocellular carcinoma after hepatectomy. *Surgery.* 1994;115:303-9.
38. Chang S, Laurent A, Tayar C, Karoui M, Cherqui D. Laparoscopy as a routine approach for left lateral sectionectomy. *Br J Surg.* 2006;94:58-63.
39. Melendez JA, Arslan V, Fischer ME, Wuest D, Jarnagin W, Fong Y. Perioperative outcomes of major hepatic resections under low central venous pressure anesthesia: blood loss, blood transfusion, and the risk of postoperative renal dysfunction. *J Am Coll Surg.* 1998;187:620-5.
40. Bhattacharya S, Jackson DJ, Beard CI, Davidson BR. Central venous pressure and its effects on blood loss during liver resection. *Br J Surg.* 1999;86:282-3.
41. Vittimberga FJ, Foley DP, Meyers WC, Callery MP. Laparoscopic surgery and the systemic immune response. *Ann Surg.* 1998;227:226-34.
42. Cook T, Dehn T. Port-site metastases in patients undergoing laparoscopy for gastrointestinal malignancy. *Br J Surg.* 1996;83:1419-20.
43. Neuhaus S, Texler M, Hewett P, Watson D. Port-site metastases following laparoscopic surgery. *Br J Surg.* 1998;85:735-41.
44. Watson D, Mathew G, Ellis T, Baigrie CF, Rofe A, Jamieson G. Gasless laparoscopy may reduce the risk of port-site metastases following laparoscopic tumor surgery. *Arch Surg.* 1997;132:166-8.
45. Laurent A, Cherqui D, Lesurtel M, Brunetti F, Tayar C, Fagniez PL. Laparoscopic liver resection for subcapsular hepatocellular carcinoma complicating chronic liver disease. *Arch Surg.* 2003;138:763-9.
46. Daniel C, Alexis L, Claude T, Stephen C, Jeanne T, Jerome L, et al. Laparoscopic liver resection for peripheral hepatocellular carcinoma in patients with chronic liver disease: midterm results and perspectives. *Ann Surg.* 2006;243:499-506.
47. Lang BH, Poon RT, Fan ST, Wong J. Perioperative and long-term outcome of major hepatic resection for solitary hepatocellular carcinoma in patients with cirrhosis. *Arch Surg.* 2003;138:1207-13.
48. Bryant R, Laurent A, Tayar C, Cherqui D. Laparoscopic liver resection-understanding its role in current practice. *Ann Surg.* 2009;250:103-11.
49. Vigiano L, Laurent A, Tayar C, Tomatis M, Ponti A, Cherqui D. The Learning Curve in Laparoscopic Liver Resection. Improved feasibility and reproducibility. *Ann Surg.* 2009;250.