

**Image of the Month****Phlegmasia Cerulea Dolens[☆]****Flegmasía cerúlea dolens**

**Ainhoa García Familiar,* Ioar Larrañaga Oyarzabal, Víctor Rodríguez Sáenz De Buruaga,
Jose María Egaña Barrenechea**

Servicio de Angiología y Cirugía Vascular, Hospital Donostia, San Sebastián, Guipúzcoa, Spain

**Fig. 1****Fig. 2**

A 50 year old man, with a prior history of stage IV rectal cancer in palliative treatment, came to the Emergency Department with sudden severe pain of 12 h duration in the left lower extremity. On physical examination edema with fovea in the left lower extremity with cyanosis below the knee (Figs. 1 and 2, arrows) was found. No popliteal or distal pulse could be found, with anesthesia and paralysis of the left foot. A Doppler ultrasound showed deep venous thrombosis of the iliac, femoral and popliteal veins. Anticoagulation was started with low molecular weight heparin. The patient progressed to venous gangrene and died 4 days later.

* Please cite this article as: García Familiar A, Larrañaga Oyarzabal I, Rodríguez Sáenz De Buruaga V, Egaña Barrenechea JM. Flegmasía cerúlea dolens. Cir Esp. 2013;91:e21.

* Corresponding author.

E-mail address: ainhoa.garciafamiliar@osakidetza.net (A. García Familiar).
2173-5077/\$ – see front matter © 2012 AEC. Published by Elsevier España, S.L. All rights reserved.