

## Image of the month

## Enterocoele Containing Caecum and Terminal Ileum at Labium Majus Level ☆

### Enterocoele conteniendo ciego e íleon terminal a nivel de labio mayor

Leire Zarain Obrador, \* Marta Cuadrado Ayuso, Cristina Rey Valcárcel, M. Dolores Pérez Díaz

Servicio de Cirugía General II, Hospital General Universitario Gregorio Marañón, Madrid, Spain



**Fig. 1** – Photograph of the patient prior to surgery. The existing tumour at labium majus level can be observed.



**Fig. 2** – CT scan coronal cut showing caecum and terminal ileum at labium majus level.

An 87-year-old woman came to the Emergency Department with a 48-hour history of intestinal obstruction and a soft and palpable 15 cm mass on the left labia majora. Relevant clinical history was a hysterectomy 44 years ago and four subsequent

surgical interventions for colpocele and cystocele; the last surgery was performed 15 years ago. The CT scan showed the presence of caecum and terminal ileum at labium majus level, between the vagina and rectum. The patient underwent

\* Please cite this article as: Zarain Obrador L, Cuadrado Ayuso M, Rey Valcárcel C, Pérez Díaz M.D. Enterocoele conteniendo ciego e íleon terminal a nivel de labio mayor. *Cir Esp.* 2013;91:676.

\* Corresponding author.

E-mail address: [leirezarainobrador@gmail.com](mailto:leirezarainobrador@gmail.com) (L. Zarain Obrador).

2173-5077/\$ – see front matter © 2013 AEC. Published by Elsevier España, S.L. All rights reserved.