

## REFERENCES

- San Norberto EM, Montes JM, Romero A, Núñez E, Vaquero C. Síndrome del ligamento arcuato medio: a propósito de tres casos y revisión de la literatura. *Angiología*. 2012;64:167-72.
- Grotemeyer D, Duran M, Iskandar F, Blondin D, Nguyen K, Sandmann W. Median arcuate ligament syndrome: vascular surgical therapy and follow-up of 18 patients. *Langenbecks Arch Surg*. 2009;394:1085-92.
- Horton KM, Talamini MA, Fishman EK. Median arcuate ligament syndrome: evaluation with CT angiography. *Radiographics*. 2005;25:1177-82.
- Matsumura Y, Nakada TA, Kobe Y, Hattori N, Oda S. Median arcuate ligament syndrome presenting as hemorrhagic shock. *Am J Emerg Med*. 2013;31:1152-4.
- Storm J, Kerr E, Kennedy P. Rare complications of a low lying median arcuate coeliac ligament. *Ulster Med J*. 2015;84:107-9.
- Karavelioğlu Y, Kalçık M, Sarak T. Dunbar syndrome as an unusual cause of exercise-induced retrosternal pain. *Turk Kardiyol Dern Ars*. 2015;43:465-7.
- Berard X, Cau J, Déglise S, Trombert D, Saint-Lebes B, Midy D, et al. Laparoscopic surgery for coeliac artery compression syndrome: current management and technical aspects. *Eur J Vasc Endovasc Surg*. 2012;43:38-42.
- Hill E, Sultan M, Chalhoub W, Jackson P, Mattar M. Median arcuate ligament syndrome: a cause of postprandial abdominal pain in a patient with ulcerative colitis. *J Med Cases*. 2014;5:344-6.

Gloria María Novo Martínez<sup>a\*</sup>, Alejandro Rodríguez Morata<sup>b</sup>, Gonzalo Alonso Argüeso<sup>b</sup>, Juan Pedro Reyes Ortega<sup>b</sup>, Rafael Gómez Medialdea<sup>b</sup>

<sup>a</sup>Hospital Universitario de León, León, Spain

<sup>b</sup>Hospital Clínico Universitario Virgen de la Victoria, Málaga, Spain

\*Corresponding author.

E-mail address: [glori\\_mry@hotmail.com](mailto:glori_mry@hotmail.com) (G.M. Novo Martínez).

2173-5077/

© 2016 Published by Elsevier España, S.L.U. on behalf of AEC.

## Treatment of Presacral Bleeding After Colorectal Surgery With Bakri Balloon<sup>☆</sup>



## Tratamiento de la hemorragia presacra tras cirugía colorrectal mediante el uso del balón de Bakri

Presacral bleeding (PB) is a non-pulsatile haemorrhage due to the disruption of the presacral venous plexus (PVP). These haemorrhages are difficult to manage and are sometimes non-controllable with conventional surgical manoeuvres.<sup>1</sup> The use of tamponade balloons is suggested as a feasible and effective option.

The Bakri balloon is a silicone balloon designed for the treatment of uterine bleeding in postpartum haemorrhage<sup>2</sup> (Fig. 1). In our series, the balloons were inserted vaginally, in perineal wounds after rectal resection or anally through the open rectal stump and filled with sterile saline solution (maximum capacity 500 mL).


The first patient underwent surgery for disease recurrence in the rectovaginal septum after resection for sigmoid adenocarcinoma. The operation involved hysterectomy and bilateral adnexectomy, resection of the posterior vaginal wall and resection of the rectum with a colorectal mechanical anastomosis and loop ileostomy. On the 5th day post-op, the patient was re-operated on due to anastomotic leak and the remaining colonic segment was resected up to the efferent

loop of the ileostomy, presenting PB that required packing. Seven days after this reoperation, the packing material was removed but, 7 days later, the patient once again underwent surgery for suspicion of another PVP haemorrhage. The approach was vaginal, and 3 balloons were inserted vaginally.

The second patient was treated surgically for disease recurrence in the right hemipelvis after abdominoperineal resection due to rectal adenocarcinoma. Surgery involved midline laparotomy with tumour resection. During surgery, PB began that was not controllable by electrocoagulation, suture or haemostatic biological materials, so a balloon was inserted through the perineum.

The third patient, with a history of subtotal colectomy due to familial adenomatous polyposis and an ileal pouch done at another hospital (without proctectomy), presented a tubular adenoma with high-grade dysplasia in the rectal stump 5 years later. Surgery involved midline laparotomy with division of the ileum above the pouch plus resection of the rectum up to the pectinate line and associated ileostomy in the RIF. During the postoperative period, the patient

<sup>☆</sup> Please cite this article as: Lopez-Lopez V, Abrisqueta J, Lujan J, Ferreras D, Parrilla P. Tratamiento de la hemorragia presacra tras cirugía colorrectal mediante el uso del balón de Bakri. *Cir Esp*. 2016;94:303-305.



**Fig. 1 - (A) Bakri balloon; (B) Dimensional perspective of Bakri balloon placement in the presacral venous plexus space, within the abdominal cavity; (C) Anal placement of the Bakri balloon in the region of the presacral venous haemorrhage.**

**Table 1 - Characteristics of Patients With Presacral Venous Bleeding.**

|           | Sex    | Age (years) | Technique used to control PB  | Blood transfusions (RBC) | No. of Bakri balloons | Placement | No. of days for control of the PB |
|-----------|--------|-------------|---|--------------------------|-----------------------|-----------|-----------------------------------|
| Patient 1 | Female | 50          | Packing with gauze<br>Monopolar electrocoagulation<br>Simple suture<br>Biological haemostatic agents      | 20                       | 3                     | Vaginal   | 4                                 |
| Patient 2 | Male   | 43          | Monopolar electrocoagulation<br>Simple suture<br>Biological haemostatic agents                            | 10                       | 1                     | Perineal  | 2                                 |
| Patient 3 | Male   | 29          | Monopolar electrocoagulation<br>Simple suture<br>Biological haemostatic agents                            | 7                        | 1                     | Anal      | 2                                 |
| Patient 4 | Female | 28          | Vascular embolisation<br>Monopolar electrocoagulation<br>Circular suture<br>Biological haemostatic agents | 15                       | 2                     | Anal      | 3                                 |

RBC: packed red blood cells; PB: presacral bleeding.

presented several episodes of anal bleeding; relaparotomy was therefore performed, which identified bleeding of the PVP that did not cease with conventional surgical manoeuvres and required the placement of 2 balloons through the non-functioning anus.

The fourth patient was diagnosed with a vascular malformation in the right pelvis who reported metrorrhagia, rectal bleeding and haematuria, which required vascular embolisation. Endoscopy revealed an anal ulcer measuring 20 cm, suggesting ischaemic colitis. The patient underwent hysterectomy, bilateral adnexectomy and division of the distal rectum at the pelvic floor and proximal sigmoid colon, closure of the rectal stump and terminal colostomy. On the 2nd day post-op, the patient presented haemoperitoneum and a haemorrhage was found at the PVP. Haemostasis techniques used included electrocoagulation and biologic haemostatic agents, but, given the persistent bleeding, 2 balloons were inserted anally (Table 1).

PB is a serious complication of pelvic tumour surgery (rectal, advanced ovarian and presacral tumours), with an incidence of 3%–9% and a mortality rate of 1%–4%.<sup>3</sup>

Monopolar electrocoagulation or suture can even aggravate PVP lesions and bleeding.<sup>4</sup> The use of surgical clips can be effective, although disadvantages include chronic pain and displacement or difficult placement due to the anatomic curvature of the sacrum.<sup>3</sup> Compression with a fragment of anterior rectus muscle and indirect electrocoagulation with monopolar forceps can achieve haemostasis, although this is not recommended for large haemorrhages or if there is contraction of the vertebral vein endings.<sup>5</sup> In these cases, bipolar coagulation seems more effective. Electrocauterisation through the epiploic appendices is useful when the origin is due to involvement of the basivertebral veins in a sacral nerve foramen.<sup>1</sup> Jiang et al.<sup>6</sup> described the circular suture of the PVP in the intact presacral fascia that surrounds the haemorrhage, although this is difficult in a narrow pelvis, previous rectal surgery or atypical anatomic pattern of the PVP. Other alternatives, such as biological haemostatic agents, are widely used but are ineffective in a massive PB.<sup>7</sup> Traditionally, packing has been used, but the need for re-operation to remove it is a disadvantage.<sup>8</sup>

McCourteny et al.<sup>9</sup> used a Sengstaken–Blakemore balloon to control a massive PB, with resolution of the haemorrhage 4 days later. Recently, Charoenkwan described the usefulness of the Bakri balloon for untreatable pelvic floor haemorrhage after hysterectomy or caesarean section when other techniques have failed.<sup>10</sup>

In our experience, the Bakri balloon has been shown to be a useful method in the management of PB after colorectal surgery when conventional treatment has failed. Its advantage is its easy placement and good adaptation to the presacral space while exerting greater intraluminal pressure than the pressure of the venous and capillary system. For its removal, surgical re-operation is not necessary. In fact, when poor haemorrhage control is still suspected, this technique enables

us to reduce the pressure gradually and even re-fill the balloon in cases of persistent bleeding.

## REFERENCES

- Lou Z, Zhang W, Meng RG, Fu CG. Massive presacral bleeding during rectal surgery: from anatomy to clinical practice. *World J Gastroenterol.* 2013;7:4039–44.
- Bakri YN. Uterine tamponade-drain for hemorrhage secondary to placenta previa-accreta. *Int J Gynaecol Obstet.* 1992;37:302–3.
- Van der Vurst TJ, Bodegom ME, Rakic S. Tamponade of presacral hemorrhage with hemostatic sponges fixed to the sacrum with endoscopic helical tackers: report of two cases. *Dis Colon Rectum.* 2004;47:1550–3.
- Filippakis GM, Leandros M, Albanopoulos K, Genetzakis M, Lagoudianakis E, Pararas N, et al. The use of spray electrocautery to control presacral bleeding: a report of four cases. *Am Surg.* 2007;73:410–3.
- Casal Núñez JE, Martínez MT, Poblador AR. Electrocoagulation on a fragment of anterior abdominal rectal muscle for the control of presacral bleeding during rectal resection. *Cir Esp.* 2012;90:176–9 [article in Spanish].
- Jiang J, Li X, Wang Y, Qu H, Jin Z, Dai Y. Circular suture ligation of presacral venous plexus to control presacral venous bleeding during rectal mobilization. *J Gastrointest Surg.* 2013;17:416–20.
- Losanoff JE, Richman BW, Jones JW. Cyanoacrylate adhesive in management of severe presacral bleeding. *Dis Colon Rectum.* 2002;45:1118–9.
- Zama N, Fazio VW, Jagelman DG, Lavery IC, Weakley FL, Church JM. Efficacy of pelvic packing in maintaining hemostasis after rectal excision for cancer. *Dis Colon Rectum.* 1988;31:923–8.
- McCourteny JS, Hussain N, Mackenzie I. Ballon tamponade for control of massive presacral haemorrhage. *Br J Surg.* 1996;83:222.
- Charoenkwan K. Use of the Bakri postpartum balloon in a patient with intractable pelvic floor hemorrhage: when other methods failed to stop postcesarean bleeding, physicians tried something new. *Am J Obstet Gynecol.* 2013;209:277.

Victor Lopez-Lopez\*, Jesús Abrisqueta, Juan Lujan, David Ferreras, Pascual Parrilla

Departamento de Cirugía General del Aparato Digestivo, Hospital Clínico Universitario Virgen de la Arrixaca, Universidad de Murcia, IMIB, Murcia, Spain

\*Corresponding author.

E-mail address: [victorlopez@gmail.com](mailto:victorlopez@gmail.com) (V. Lopez-Lopez).

2173-5077/

© 2015 AEC. Published by Elsevier España, S.L.U. All rights reserved.