

REFERENCES

1. Abcarian H, Alexander-Williams J, Christiansen J, Johanson J, Killingback M, Nelson RL, et al. Bening anorectal disease: definition, characterization and analysis of treatment. *Am J Gastroenterol.* 1994;89 Suppl. 8:S182-93.
2. Nivatvongs S. Hemorrhoids. In: Philip G, editor. *Principles and practice of surgery.* 3rd ed. Nueva York: Informa Healthcare; 2007; p. 143-66.
3. Madoff RD, Fleshman JW, Clinical Practice Committee, American Gastroenterological Association. American Gastroenterological Association technical review on the diagnosis and treatment of hemorrhoids. *Gastroenterology.* 2004;126:1463-73.
4. Lewis RT, Maron DJ. Anorectal Crohn's disease. *Surg Clin N Am.* 2010;90:83-97.
5. Radcliffe AG, Ritchie JK, Hawley PR, Lennard-Jones JE, Northover JM. Anovaginal and rectovaginal fistulas in Crohn's disease. *Dis Colon Rectum.* 1988;31:94-9.
6. Morrison JG, Gathright JB Jr, Ray JE, Ferrari BT, Hicks YC, Timmcke AE. Results of operation for rectovaginal fistula in Crohn's disease. *Dis Colon Rectum.* 1989;32:497-9.
7. Cracco N, Zinicola R. Is haemorrhoidectomy in inflammatory bowel disease harmful? An old dogma re-examined. *Colorectal Dis.* 2014;16:516-9.
8. D'Ugo S, Franceschilli L, Cadeddu F, Leccesi L, Blanco G del V, Calabrese E, et al. Medical and surgical treatment of haemorrhoids and anal fissure in Crohn's disease: a critical appraisal. *BMC Gastroenterol.* 2013;13:47.
9. Vidal V, Sapoval M, Sielezneck Y, de Parades V, Tradi F, Louis G, et al. Emborrhoid: a new concept for the treatment of hemorrhoids with arterial embolization: the first 14 cases. *Cardiovasc Intervent Radiol.* 2015;38:72-8.
10. Vidal V, Louis G, Bartoli JM, Sielezneck I. Embolization of the hemorrhoidal arteries (the emborrhoid technique): a new concept and challenge for interventional radiology. *Diagn Interv Imaging.* 2014;95:307-15.

Vicent Primo Romaguera^{a*}, Abel Gregorio Hernández^b,
Luís Andreo Hernández^b, Eduardo de la Morena Valenzuela^a

^aServicio de Cirugía General y Digestiva, Hospital de Dénia Marina-Salud, Dénia, Alicante, Spain

^bServicio de Radiología, Hospital de Dénia Marina-Salud, Dénia, Alicante, Spain

*Corresponding author.

E-mail address: sentopr@gmail.com (V. Primo Romaguera).

2173-5077/

© 2017 AEC. Published by Elsevier España, S.L.U. All rights reserved.

Surgical Treatment of Supraclavicular and Internal Mammary Recurrence of Breast Cancer[☆]



Tratamiento quirúrgico de la recidiva supraclavicular y a nivel de la mamaria interna por un carcinoma de mama

Brito et al.¹ treated patients with supraclavicular lymph node involvement using multimodal therapy with curative intent (chemotherapy [CTx], surgery and radiotherapy). With this method, they obtained better results than in patients who presented distant disease that was not supraclavicular. These results led to the review and modifications of the American Joint Committee on Cancer (AJCC-TNM) classification. Since its sixth edition, metastases in the ipsilateral supraclavicular lymph nodes have ceased to be considered distant metastases (M1) and have been classified as locoregional disease (N3c).^{2,3} To this day, the debate continues and no consensus has been reached for the treatment of these patients.

Likewise, contralateral axillary involvement should be considered stage IV and therefore the treatment of these patients should be palliative.⁴ However, there is a hypothesis that dissemination to the contralateral axilla could occur by the lymphatic pathway, therefore locoregional curative treatment would be justified.⁵ Therefore, the management

of this situation is not standardized and must be individualized for each patient.

We present a controversial clinical case using multimodal treatment, associating surgery of a metastatic breast carcinoma with contralateral and ipsilateral supraclavicular lymph nodes.

A 57-year-old woman presented with a 10 cm ulcerated tumor infiltrating the skin and chest wall of the right breast. Mammography, magnetic resonance imaging with gadolinium contrast and core-needle biopsy diagnosed the mass as invasive ductal carcinoma (IDC) of the right breast with ipsilateral and contralateral axillary lymph node involvement (cT4c cN1c M1). Immunohistochemistry study reported: estrogen receptors (ER) +++, progesterone receptors (PR) +, cerbB2 ++ (positive FISH), Ki 67 20%.

After 8 cycles of CTx with docetaxel, trastuzumab and pertuzumab with good radiological response, radical mastectomy was performed with right and left axillary lymph node dissection. The pathology study of the piece reported an

[☆] Please cite this article as: de Andrés Gómez A, Villalba Ferrer F, Navarro Moratalla C, Guijarro Jorge R, Fuster Diana C. Tratamiento quirúrgico de la recidiva supraclavicular y a nivel de la mamaria interna por un carcinoma de mama. *Cir Esp.* 2018;96:241-244.




Fig. 1 – Total-body PET/CT: (a) coronal PET image showing hyperuptake of the internal breast (arrow); (b) axial PET/CT showing the same lesion (circle); (c) coronal PET/CT image with the lesion in white (arrow); (d) coronal PET view with hyperuptake in the right supraclavicular region; (e) axial PET/CT slice demonstrating the same involvement (circle); (f) coronal PET/CT showing right cervical hyperuptake in white (arrow).

IDC lesion measuring 3.5×3 cm (RE –, RP –, cerb B2 +++ and Ki 67 30%), grade 1, N + (1/14) and an invasive lobular carcinoma measuring 2×1.5 cm (RE –, RP –, cerbB2 –, Ki 67 5%–10%). The axillary lymphadenectomy dissection revealed bilateral lymph node involvement: 1/14 in the right axilla and 4/24 in the left axilla.

The patient did not receive radiotherapy due to limited mobility of the right glenohumeral joint and followed adjuvant treatment with trastuzumab-pertuzumab every 21 days.

Positron emission tomography/computed tomography (PET/CT) one year after surgery (Fig. 1) detected right supraclavicular lymph node and internal mammary involvement, confirmed by fine-needle aspiration. The internal mammary ultrasound detected nodal involvement with a hypoechoic lesion measuring 4×3×4 cm.

Three cycles of CTx were administered with TDM1 (trastuzumab emtansine) and subsequent PET/CT showed a decrease in size and right supraclavicular lymph node metabolism (SUV 2.1, previously 3.3), but increased internal mammary metabolism (SUV 7.6, previously 7) compared to the previous study. Surgical intervention was performed one month after the end of CTx, involving functional right lateral cervical dissection, excision of the internal mammary chain lesion and 5th, 6th and 7th anterior costal arches due to macroscopic involvement of the intercostal muscles. The wall defect was covered with a double-sided mesh (Fig. 2). The

patient progressed favorably and was discharged on the 5th postoperative day.

The pathological anatomy of the internal mammary chain lesion revealed a metastasis measuring 5×1.6 cm, with involvement of soft tissues (striated muscle and fibrous-fatty tissue) by lobular carcinoma, with no bone involvement, free resection margins and immunohistochemistry: RE –, RP –, cerbB2 –, Ki 67 5%–10%. From the cervical lymph node dissection, 52 lymphadenopathies were isolated with no tumor involvement. The patient currently continues with adjuvant CTx with TDM1 and is disease-free 9 months after the intervention.

There are numerous publications that evaluate the results of radiotherapy in the involvement of the internal and supraclavicular mammary arteries^{6,7}; however, the role of surgery has not yet been studied.⁸ In the review of the literature, there is a study that included patients with recurrence of the disease at the supraclavicular level where surgery was part of the treatment and concluded that surgical excision of these supraclavicular lymph nodes would be a prognostic factor for overall survival.⁹

The negative pathology results of the lymph nodes removed in the cervical lymphadenectomy could be explained by a complete response to CTx that was already evident in the re-evaluation PET after the end of the last cycle, where there was a decrease in metabolic activity.




Fig. 2 – Surgery image: (a) after lateral cervical lymphadenectomy; (b) right supraclavicular lymphadenopathy; (c) mesh on chest wall; (d) metastasis of the internal breast.

Although locoregional therapy is not standard in the treatment of *de novo* stage IV cancers, it is usually used as a palliative treatment.¹⁰ In this group of patients, this would be the case by including surgery in multimodal treatment for better control of the disease. There are authors who defend that patients with advanced tumor disease and certain cancer subtypes, such as luminal-like or Her2, can be offered local treatment with surgery or radiotherapy when this means better control of the primary tumor or the regional involvement in order to improve overall survival.¹⁰ In cases in which radiotherapy cannot be applied, as in our patient, surgery would be the option for better locoregional control of the disease.

REFERENCES

1. Brito RA, Valero V, Buzdar AU, Booser DJ, Ames F, Strom E, et al. Long-term results of combined-modality therapy for locally advanced breast cancer with ipsilateral supraclavicular metastases: the University of Texas M.D. Anderson Cancer Center experience. *J Clin Oncol.* 2001;19:628-33.
2. Greene FLPD, Fleming ID, Fritz AG, Balch CM, Haller DG, Morrow M. *AJCC cancer staging manual*, 6th ed. New York: Springer Verlag; 2002.
3. Singletary SE, Allred C, Ashley P, Bassett LW, Berry D, Bland KI, et al. Revision of the American Joint Committee on Cancer staging system for breast cancer. *J Clin Oncol.* 2002;20:3628-36.
4. Sakurai K, Enomoto K, Kitajima A, Tani M, Amano S. A case of advanced breast cancer successfully treated with multi-disciplinary therapy and S-1 administration. *Gan To Kagaku Ryoho.* 2007;34:2065-7.
5. Castillo G, Sabadell D, Carreras R, Vernet MM. Metástasis ganglionares axilares contalaterales en cáncer de mama: controversia sobre su vía de diseminación, manejo y pronóstico. *Rev Senol Patol Mamar.* 2016;29:86-9.
6. Poortmans PM, Collette S, Kirkove C, Van Limbergen E, Budach V, Struikmans H, et al. EORTC Radiation Oncology and Breast Cancer Groups. Internal mammary and medial supraclavicular irradiation in breast cancer. *N Engl J Med.* 2015;373:317-27.
7. Poortmans PM, Collette S, Kirkove C, van Limbergen E, Budach V, Struikmans H, et al. EORTC Radiation Oncology and Breast Cancer Groups. Internal mammary and medial supraclavicular irradiation in breast cancer. *N Engl J Med.* 2015;373:317-27.

8. Rotenhuis BA, Klem TM, Vrijland WW. Treatment outcome in breast cancer patients with ipsilateral supraclavicular lymph node metastasis at time of diagnosis: a review of the literature. *Eur J Surg Oncol*. 2013;39:207-12.
9. Chatterjee S, Lee D, Kent N, Wintle T, Mott JH, Kelly CG, et al. Managing supraclavicular disease from breast cancer with brachial plexus-sparing techniques using helical tomotherapy. *Clin Oncol (R Coll Radiol)*. 2011;23:101-7.
10. Chen P, Cheng S, Hung C, Yu B, Chen C. Locoregional therapy in luminal-like and HER2-enriched patients with de novo stage IV breast cancer. *Springer Plus*. 2013;2:1-6.

Alejandra de Andrés Gómez^{a*}, Francisco Villalba Ferrer^a, Carla Navarro Moratalla^a, Ricardo Guijarro Jorge^b, Carlos Fuster Diana^a

^aServicio de Cirugía General y del Aparato Digestivo, Hospital General Universitario de Valencia, Valencia, Spain

^bServicio de Cirugía Torácica, Hospital General Universitario de Valencia, Valencia, Spain

*Corresponding author.

E-mail address: adeangomez@gmail.com

(A. de Andrés Gómez).

2173-5077/

© 2017 AEC. Published by Elsevier España, S.L.U. All rights reserved.

Ileocolic Intussusception by Appendicular Mucocele[☆]



Intususcepción ileocólica por mucocele apendicular

Appendiceal intussusception is a rare pathology. It represents only 0.01% of indications for appendectomy¹ and is a diagnostic challenge.²

We present the case of a 20-year-old female patient who came to the Emergency Room with colicky abdominal pain in the right iliac fossa that had lasted for several days, with no other associated symptoms. Blood work-up showed no alterations. Due to the persistence of the pain, imaging tests were indicated. Ultrasound revealed an image in the right iliac fossa. A CT scan showed a tubular structure with hypodense content at the hepatic angle that ended in a cecal sac (Fig. 1), as well as the invagination of the terminal ileum and the ileocecal valve in the right colon (Fig. 2). The findings were compatible with the diagnosis of invagination of an appendiceal mucocele, therefore urgent laparoscopic ileocecectomy was performed. The postoperative evolution was favorable, and the patient was discharged from hospital on the third postoperative day. The pathology study reported an appendi-

ceal mucocele with no malignant histological features whose base protruded into the lumen of the cecum, causing intussusception.

Appendiceal intussusception is defined as the invagination of a portion of the appendix into its own lumen or that of the cecum. It is secondary to abnormal peristaltic movements caused by inflammation of the appendix and favored by lax mesoappendices with a broad appendicular lumen or by very mobile and thin appendices.³ This pathology appears more frequently in adults (76%) than in children, and in women than in men, with a 2:1 ratio. The most frequent age at onset is the fourth decade of life.² In pediatric patients, the most frequent cause is inflammation of the appendix, while in adults the most common etiology is endometriosis (33%) followed by appendiceal mucocele (19%) and appendiceal inflammation (19%). In the remaining cases, the etiology is tumorous and there have been descriptions of carcinoid tumors, metastasis, hamartomas or lymphomas.^{2,4,5}




Fig. 1 – Appendix with abundant hypodense content in the lumen of the colon in the region of the hepatic angle.



Fig. 2 – Ileocecal valve and ileum invaginated in the lumen of the cecum and right colon.

[☆] Please cite this article as: Sagarra Cebolla E, Burgos de la Obra Ferrari E, Velasco Sánchez EA, Gómez Lanz L. Intususcepción ileocólica por mucocele apendicular. *Cir Esp*. 2018;96:244-245.