



## Original article

# Risk Factors for Bad Outcomes in Incisional Hernia Repair: Lessons Learned From the National Registry of Incisional Hernia (EVEREG)<sup>☆,☆☆</sup>



José Antonio Pereira,<sup>a,b,\*</sup> Blanca Montcusí,<sup>a</sup> Manuel López-Cano,<sup>c,d</sup>  
Pilar Hernández-Granados,<sup>e</sup> Laura Fresno de Prado,<sup>b</sup> Members of the EVEREG Registry<sup>◇</sup>

<sup>a</sup>Servicio de Cirugía General, Hospital Universitari del Mar, Barcelona, Spain

<sup>b</sup>Departament de Ciències Experimentals i de la Salut, Universitat Pompeu Fabra, Barcelona, Spain

<sup>c</sup>Servicio de Cirugía General, Hospital Vall d'Hebron, Barcelona, Spain

<sup>d</sup>Departament de Cirurgia, Universitat Autònoma de Barcelona, Barcelona, Spain

<sup>e</sup>Hospital Universitario Fundación Alcorcón, Madrid, Spain

## ARTICLE INFO

## Article history:

Received 27 October 2017

Accepted 24 March 2018

Available online 8 August 2018

## Keywords:

Incisional hernia

Registries

Risk factors

## A B S T R A C T

**Introduction:** Registries are powerful tools for identifying factors predicting bad results. Our objective was to analyze data from the Spanish Registry of Incisional Hernia (EVEREG) to detect risk situations for the development of complications and recurrences.

**Methods:** We have analyzed data of the cohort of hernias registered during the period from July 2012 to June 2014. We have compared the data between complicated and non-complicated patients in the short and long term follow-up. Data compared were: patient demographics, comorbid condition, hernia defect characteristics and surgical technique to determine which of them may be predictors of poor outcomes.

**Results:** During the period of study, we collected data from 1336 hernias (43.7% males; 56.3% females) with a mean age of 63.6 years (SD 12.4) and BMI of 30.4 (SD 5.4). In the multivariate analysis, factors associated with complications were: age >70 years, previous neoplasm, diameter greater than 10 cm, previous repair and bowel resection. Factors related with recurrences were: parastomal hernia, previous repair, emergency repair, postoperative complications and reoperation. A separation of components was the only protective factor for this type of analysis (OR 0.438; CI 0.27–0.71; P = .0001).

<sup>\*</sup> Please cite this article as: Pereira JA, Montcusí B, López-Cano M, Hernández-Granados P, Fresno de Prado L, Miembros del registro EVEREG. Factores de riesgo de mala evolución en la reparación de hernias incisionales: Lecciones aprendidas del Registro Nacional de Hernia Incisional (EVEREG). Cir Esp. 2018;96:436–442.

<sup>☆☆</sup> This work was presented at the Congress of the European Hernia Society (May 2017, Vienna) with the title: “Registries as tool to identify risk factors for bad outcomes”.

<sup>\*</sup> Corresponding author.

E-mail address: 86664@parcdesalutmar.cat (J.A. Pereira).

<sup>◇</sup> Members of the EVEREG registry are listed in [Appendix A](#).

*Conclusions:* Risk factors for the development of complications and recurrences must be considered for promoting preoperative patient prehabilitation, planning the surgical technique and referring patients to specialized abdominal wall units.

© 2018 AEC. Published by Elsevier España, S.L.U. All rights reserved.

## Factores de riesgo de mala evolución en la reparación de hernias incisionales: Lecciones aprendidas del Registro Nacional de Hernia Incisional (EVEREG)

### R E S U M E N

#### Palabras clave:

Hernia incisional  
Registros  
Factores de riesgo

*Introducción:* Los registros son herramientas potentes para identificar situaciones de riesgo de mala evolución. Nuestro objetivo ha sido analizar los datos del Registro Nacional de Hernia Incisional (EVEREG) para detectar situaciones de riesgo para el desarrollo de complicaciones y recidivas.

*Métodos:* Se analizan los datos de la cohorte de hernias registradas en el período desde julio de 2012 hasta junio de 2014. Se realiza una comparación estadística entre las hernias que presentaron complicaciones y recidivas, y su relación con los pacientes, las características de las hernias y la técnica quirúrgica, para determinar cuáles de ellos pueden predecir malos resultados.

*Resultados:* En el período de estudio disponemos de datos de 1.336 hernias (43,7% varones; 56,3% mujeres), con una edad media de 63,6 años (DE: 12,4) e IMC de 30,4 (DE: 5,4). En el estudio multivariante las variables asociadas a la presencia de complicaciones fueron: edad superior a 70 años, neoplasia, longitud del defecto mayor de 10 cm, reparación previa y resección intestinal. Las variables asociadas a recidiva fueron: las hernias paraestomales, la reparación previa, la cirugía urgente, la aparición de complicaciones postoperatorias y la reintervención quirúrgica. El uso de una separación de componentes fue el único factor protector en este tipo de análisis (OR: 0,438; IC: 0,27-0,71;  $p = 0,001$ ).

*Conclusiones:* La presencia de factores de riesgo para la aparición de complicaciones y recurrencias debe ser tenida en cuenta con el fin de proyectar la prehabilitación del paciente para la cirugía, planificar la técnica quirúrgica y su derivación a unidades especializadas.

© 2018 AEC. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Introduction

National registries are very powerful tools that provide a large volume of data. Their analysis enables us to reach conclusions in order to: improve treatment, identify in advance which patients are at higher risk of complications, and identify subgroups requiring specific measures to avoid the appearance of postoperative complications.<sup>1-3</sup>

The use of registries for abdominal wall surgery has extended worldwide in recent years, and there are numerous contributions related to the analysis of the registered data to achieve a better understanding of this highly prevalent pathology and obtain better results in their treatment.<sup>4-7</sup>

The Spanish National Registry of Incisional Hernias (EVEREG) was initiated in July 2012. By July 2017, it included data from 160 Spanish hospitals and a total of 7505 cases of hernias related to previous laparotomy (no primary ventral hernias were registered). An initial analysis of the first years of data collection has been previously published, which demonstrated the need to introduce improvements in treatment, especially to reduce the number of complications and recurrences.<sup>8</sup>

The aim of this study is to analyze, in a cohort of registered cases, variables for patient-related data, surgical situation, hernia characteristics and the surgical technique used in order to predict the appearance of postoperative complications and recurrences, and, based on this knowledge, suggest the implementation of preoperative, technical and postoperative measures to improve results

## Methods

The registry is promoted by the Abdominal Wall and Sutures Division of the Spanish Association of Surgeons. Data are stored in an external server, whose global data are only accessible by the surgeons who head this nation-wide initiative.<sup>8</sup> The data were collected prospectively from the Spanish National Registry of Incisional Hernias between July 2011 and June 2014. During this period, a total of 2181 hernias had been registered, and 1336 cases (61.25%) were considered valid for the analysis. The causes for exclusion of the remaining registries were: date of intervention prior to the start of the registry ( $n = 108$ ), errors in data collection that impede analysis ( $n = 237$ ) and absence of follow-up data during the first 30 days after surgery ( $n = 530$ ). The data

collection process has been described previously.<sup>8</sup> In short, it is an online database maintained by surgeons from each hospital that outlines the parameters for patients, hernias, operations, complications, and follow-up after one month, 6 months, one year and 2 years, for each of the incisional hernias treated. Complications were recorded specifically and classified according to the affected organ, situation (intraoperative and postoperative) and the time of onset (early or late).

The hernias were categorized especially for this study in terms of complexity (low, moderate and high) according to the criteria described by Slater et al.,<sup>9</sup> which considered patient/hernia characteristics and the circumstances of the surgery.

### Statistical Analysis

The data were exported to the SPSS v.22 statistical analysis software (IBM Inc., Rochester, MN, USA). Quantitative variables were expressed as mean  $\pm$  standard deviation and qualitative variables as proportions. To analyze the association between qualitative variables, we used the Chi-squared test ( $\chi^2$ ), or Fisher's test when necessary, as well as the Student's t test or the Mann-Whitney test for quantitative variables. The normality of the distribution of the quantitative variables was verified using the Kolmogorov-Smirnov test. Statistical significance was established with a  $P < .05$ . In the multivariate analysis, the predictive capacity of each variable and its independence from the other predictor variables were analyzed using a binomial logistic regression model by sequentially introducing the variables with an input  $F$  of 0.5.

## Results

The cohort of patients analyzed is shown in [Table 1](#). During the postoperative period, complications occurred in 429 patients (32.1%), 22 of whom (1.7%) died. After a median follow-up of 15.43 months (CI: 12.3-22.5), there were 265 recurrences (20.7%).

In the analysis of the patient-related factors, age over 70, male gender, history of neoplasm and anesthetic risk correlated with a statistically significant occurrence of postoperative complications ([Table 2](#)).

The situation in which the surgery was performed (elective vs urgent) correlated with the appearance of complications (elective 30.5% vs urgent 52%,  $P < .0001$ ), mortality (elective 1.4% vs urgent 10%,  $P < .0001$ ) and recurrences (elective 20% vs urgent 32%;  $P = .006$ ).

The characteristics of the hernias that were associated with a greater number of complications were ([Table 3](#)): previous incisional hernia repair, location in the midline, hernias associated with an ostomy, and the diameter of the defect (transverse and longitudinal diameters greater than 10 cm were both significant). In general, more complex hernias were associated with a greater number of complications (45.2% vs 29.7%;  $P < .0001$ ), mortality (11% vs 0.4%;  $P < .0001$ ) and recurrences (27.8% vs 19.7%;  $P = .011$ ). In contrast, trocar hernias presented a lower frequency of complications ( $P < .0001$ ) and recurrences ( $P = .011$ ).

**Table 1 – Characteristics of the Cohort Patients.**

	n = 1336
Age (SD)	63.9 (12.5)
Age >70 yrs, n (%)	500 (37.4)
Sex M/F, n (%)	583/753 (43.6/56.4)
BMI (SD)	30.45 (5.5)
Obesity BMI >30, n (%)	639 (47.8)
Overweight BMI >25, n (%)	1140 (85.3)
Smoker, n (%)	237 (17.7)
COPD, n (%)	206 (15.4)
Diabetes, n (%)	295 (22.1)
Immunosuppression, n (%)	85 (6.4)
Cancer, n (%)	373 (27.9)
ASA III/IV, n (%)	414 (31.0)
Complexity, n (%)	
Low	284 (21.7)
Moderate	836 (64.0)
High	187 (14.3)

Regarding the surgical technique ([Table 4](#)), the approach (open or laparoscopic) showed no significant differences. The use of mesh correlated with a lower number of recurrences (20.6% vs 38.1%;  $P = .05$ ). The position of the mesh showed no influence on the development of complications or recurrence. On the other hand, significant differences were detected when the mesh fixation systems were compared ([Table 4](#)). In order to better examine these technical data, we considered the approach separately. In open surgery, the use of mesh was associated with fewer recurrences (20.3% vs 38.1%;  $P = .05$ ). Upon comparing the prostheses according to their configuration (reticular, laminar and composite), no differences were found. The use of staples as a fixation system in open surgery presented a greater number of complications (39.5% vs 31.1%;  $P = .02$ ), but with no influence on mortality and recurrences. The monofilament material produced the fewest complications (30.3% vs 38.3%;  $P = .03$ ) and recurrences (17.3% vs 30.0%;  $P < .0001$ ). The use of absorbable suture for mesh fixation was associated with more complications (36.9% vs 28.8%;  $P = .008$ ) and recurrences (26.9% vs 15.7%;  $P < .0001$ ). There were no differences between the use of continuous or simple sutures.

In open surgery, component separation presented complications more frequently (45.8% vs 30.5%;  $P < .0001$ ) as well as mortality (6.3% vs 1.4%;  $P < .0001$ ), but a lower incidence of recurrences (13.9% vs 21.9%;  $P = .015$ ). The association of another surgical procedure had a negative influence on complications (44.5% vs 29.5%;  $P < .0001$ ) and mortality (7.7% vs 0.6%;  $P < .0001$ ), but not in recurrences (18.7% vs 21.2%;  $P = .40$ ).

In the laparoscopic repair group, composite mesh was associated with a greater number of recurrences, although without reaching statistical significance (25.6% vs 10.8%;  $P = .06$ ). Physiomesh (Ethicon, NJ, USA) was not associated with a greater number of complications, but instead with a significantly higher frequency of recurrences (29.1% vs 14.5%;  $P = .03$ ). No other parameter was associated with a higher frequency of complications and recurrences.

In the multivariate logistic regression analysis ([Table 5](#)), a significant relationship was demonstrated between the

**Table 2 – Comparison of Patient-Related Parameters and the Appearance of Postoperative Complications and Recurrences.**

	Complications			Recurrences		
	Yes	No	P	Yes	No	P
Age (SD)	65.6 (12.5)	63.1 (12.4)	.001	63.6 (12.5)	63.6 (12.4)	.90
Age >70 yrs, n (%)	187 (37.4)	242 (28.9)	.001	103 (21.5)	170 (20.5)	.68
Sex M/F, n (%)	206/223 (35.3/29.6)	377/530 (64.7/70.4)	.03	125/148 (21.9/20.1)	447/589 (78.1/79.9)	.43
BMI (SD)	30.6 (5.4)	30.4 (5.5)	.63	30.8 (5.4)	30.3 (5.4)	.27
Obesity BMI >30, n (%)	215 (33.6)	213 (30.9)	.30	135 (21.6)	136 (20.1)	.51
Overweight BMI >25, n (%)	357 (31.3)	71 (37.8)	.08	237 (21.2)	34 (18.6)	.42
Smoker, n (%)	69 (29.1)	360 (32.8)	.28	53 (22.5)	220 (20.5)	.50
COPD, n (%)	78 (37.9)	351 (31.1)	.054	42 (21.8)	231 (20.7)	.74
Diabetes, n (%)	102 (34.6)	327 (31.4)	.30	56 (19.9)	217 (21.1)	.66
Immunosuppression, n (%)	28 (32.9)	401 (32.1)	.86	18 (22.2)	255 (20.8)	.75
Cancer, n (%)	156 (41.8)	273 (28.3)	<.0001	80 (22.0)	193 (20.4)	.51
ASA III/IV, n (%)	161 (38.9)	265 (29)	<.0001	90 (23.0)	183 (20.1)	.23

**Table 3 – Comparison of Hernia Characteristics With the Appearance of Postoperative Complications and Recurrences.**

	Complications			Recurrences		
	Yes	No	p	Yes	No	p
Previous repair, n (%)	126 (41.2)	303 (29.4)	<.0001	90 (30.4)	183 (18.1)	<.0001
Complexity – low, n (%)	66 (23.2)	218 (76.8)	.002	57 (20.1)	227 (79.9)	.72
Complexity – moderate, n (%)	264 (31.6)	572 (68.4)	.41	163 (19.5)	673 (80.5)	1.00
Complexity – high, n (%)	95 (45.2)	334 (29.7)	<.0001	52 (27.8)	221 (19.7)	.011
Midline, n (%)	292 (35.2)	137 (27.1)	.002	172 (21.3)	101 (20.1)	.58
Trocar, n (%)	43 (19.0)	386 (34.8)	<.0001	33 (14.6)	240 (22.2)	.011
Ostomy, n (%)	35 (51.5)	394 (31.1)	<.0001	25 (38.5)	248 (19.9)	<.0001
Length >10 cm, n (%)	172 (41.7)	205 (26.8)	<.0001	88 (22.4)	160 (21.1)	.60
Width >10 cm, n (%)	99 (41.3)	280 (29.9)	.001	51 (22.6)	197 (21.3)	.67

**Table 4 – Comparison of the Technical Circumstances and the Appearance of Postoperative Complications and Recurrences.**

	Complications			Recurrences		
	Yes	No	P	Yes	No	P
Laparoscopy, n (%)	42 (25.8)	387 (33.0)	.064	36 (22.2)	237 (20.7)	.65
Mesh, n (%)	417 (31.8)	12 (46.2)	.12	265 (20.6)	8 (38.1)	.05
Two mesh, n (%)	62 (37.8)	362 (31.0)	.08	32 (20.1)	241 (21.0)	.81
Type of mesh, n (%)						
Reticular	266 (31.6)	151(32.3)	.80	157 (18.9)	108 (23.5)	.051
Laminar	14 (31.8)	403 (31.8)	.99	11 (26.2)	254 (20.4)	.36
Composite	139 (32.6)	278 (31.4)	.67	97 (23.3)	168 (19.3)	.09
Mesh position, n (%)						
Suprafascial	194 (32.1)	221 (31.5)	.90	117 (19.7)	147 (21.3)	.49
Bridge	13 (28.6)	402 (31.2)	.65	8 (16.7)	256 (20.7)	.59
Retromuscular	128 (31.8)	287 (30.7)	.85	82 (20.7)	182 (20.5)	.90
Intraperitoneal	79 (31.5)	336 (31.0)	.93	57 (23.3)	207 (19.9)	.25
Fixation with staples, n (%)	126 (33.8)	291 (31.1)	.34	77 (20.9)	188 (20.5)	.87
Fixation with absorbable sutures, n (%)	134 (37.0)	176 (28.8)	.007	95 (26.8)	94 (15.7)	<.0001
Fixation with monofilament, n (%)	236 (30.2)	75 (38.7)	.02	132 (17.3)	59 (27.8)	<.0001
Defect closure, n (%)	333 (33.6)	96 (27.8)	.05	192 (19.8)	81 (24.0)	.09
Associated procedures, n (%)	129 (42.2)	300 (29.1)	<.0001	51 (18.0)	222 (21.7)	.17
Component separation, n (%)	89 (45.9)	340 (29.8)	<.0001	25 (13.7)	248 (22.0)	.01

**Table 5 – Multivariate Analysis of Variables Associated With Postoperative Complications.**

	OR	CI	P
Age >70 yrs	1.370	1.48-1.79	.021
Cancer	1.568	1.18-2.08	.002
Length >10 cm	1.663	1.27-2.17	<.0001
Previous repair	1.718	1.28-2.30	<.0001
Bowel resection	2.124	1.20-3.75	.01
Urgent surgery	1.835	1.09-3.07	.021
Separation comp.	1.543	1.08-2.20	.017

**Table 6 – Multivariate Analysis of Variables Associated With Recurrences.**

	OR	CI	P
Parastomal	2.379	1.26-4.50	.008
Previous repair	2.031	1.47-2.80	<.0001
Urgent surgery	1.830	1.06-3.15	.029
Post-op complications	1.590	1.15-2.20	.005
Reoperation	2.599	1.34-5.03	.005
Separation comp. <sup>a</sup>	0.438	0.27-0.71	.001

<sup>a</sup> Associated with a lower rate of recurrences.

appearance of complications and age over 70 years, history of neoplasm, longitudinal diameter >10 cm, previous repair, urgent surgery, bowel resection and component separation. The combination of these factors was able to predict 70.4% of complications.

In the multivariate study, factors associated with a higher frequency of recurrences included: parastomal hernias, previous repair, urgent surgery, appearance of postoperative complications and reoperation (Table 6). The use of a separation of components was the only protective factor in this analysis (OR: 0.438; CI: 0.27-0.71; P = .001). In view of the fact that urgent surgery was one of the main factors both for complications and recurrences, we performed an analysis only of patients undergoing elective surgery. After logistic regression, the parameters that maintained a significant relationship with complications were: age over 70 years (OR: 1.329; CI: 1.00-1.75; P = .04), history of neoplasia (OR: 1.572; CI: 1.17-2.11; P = .003), longitudinal diameter >10 cm (OR: 1.364; CI: 1.20-1.54; P < .0001), previous repair (OR: 1.720; CI: 1.27-2.33; P < .0001) and associated dermolipsectomy (OR: 2.220; CI: 1.04-4.75; P < .04). For recurrences in elective surgery, the model yielded: parastomal hernias (OR: 2.538; CI: 1.31-4.91; P = .006), transverse diameter >10 cm (OR: 1.316; CI: 1.09-1.58; P = .003) and previous repair (OR: 2.062; CI: 1.47-2.89; P < .0001); meanwhile, component separation was a protective factor (OR: 0.397; CI: 0.23-0.67; P = .001). Both models were able to predict 85% of complications and recurrences in elective surgery.

## Discussion

The analysis of our study revealed very interesting data that can be applied to the treatment of incisional hernias. First of all, the registry itself, as previously shown by registries from

other countries,<sup>4-7</sup> confirms its usefulness for the collection and analysis of data. The main limitations of our study are the inability to study all the patient data recorded in the registry, as many cases from the database were not useful due to the lack of basic elements for analysis or errors in their compilation. Another limitation is that they are just a sample of reality, because Spain does not have a universal registry like other national registries,<sup>5</sup> which may lead to biases related to the type of patients and the hospitals that treat them. In our study, the high frequency of complications, affecting 32.1% of surgeries, is striking. This demonstrates that we are facing a pathology with a considerable surgical risk. Likewise, the number of recurrences (20.7%) is high, demonstrating that improvements are necessary for treatment.

Patient-related risk factors affect the appearance of postoperative complications and not recurrence. Age over 70 years, male sex, history of cancer and anesthetic risk were significantly associated with complications, while other factors such as COPD, DM and immunosuppression did not reach significant differences. These factors must be taken into account in order to prioritize the programming of elective repair, as has been suggested by other authors.<sup>10,11</sup>

The lack of correlation between the BMI and complications/recurrences is noteworthy, since other authors<sup>9,12-14</sup> have found an association of this parameter with both. This is probably due to the characteristics of our population, in which a considerable number of patients were obese with a high average BMI (30.45 kg/m<sup>2</sup>).

We found this specific datum very interesting as, like other authors,<sup>12</sup> we believe that reducing the BMI of patients should be a priority objective to reduce postoperative complications and recurrences. This is also one of the few patient-related factors that can be modified before surgery.

The circumstances of the repair correlated with a greater number of complications and recurrences. Therefore, any repair in an emergency context could have better results if treated by an expert team. Other studies on urgent incisional hernia surgery<sup>10</sup> have shown a low frequency of intestinal resections due to strangulation and even lower mortality than other types of hernias, so that initially considering a conservative approach can be safe in well-selected cases.

Hernia characteristics are crucial to be able to predict the appearance of both postoperative complications and recurrences. As a protective variable, our data show that trocar herniation had the lowest recurrence rate, although this was not confirmed in the multivariate analysis. In this case, it is likely that these data are related to the size of the defect.

In studying complexity,<sup>9</sup> we have found that this classification system is useful for predicting complications and recurrences. In fact, all the statistically significant factors in our study are taken into account in this classification system. So, in our opinion, our study validates this method as useful for the decision-making process of incisional hernia repair, and even encourages us to again recommend that moderately to highly complex hernias be treated by expert surgeons whenever possible.

As for the technical aspects, the approach, the type of mesh used and its position do not stand out as factors related with prognosis. Other details, however, are relevant. Mainly, the association of component separation correlated with a greater number of complications, probably because it was used in

more complex hernias. However, patients treated with this technical resource presented fewer recurrences. These same data were maintained among the factors revealed in the multivariate study, both in the group of patients as a whole and in only those undergoing elective surgery. Again, we found another detail that seems to indicate that treatment conducted by specialized teams could be associated with better prognosis, this time in the case of recurrences.

When the repair was associated with other procedures, there was a higher frequency of complications in the univariate analysis. Specifically, bowel resection was the only significant procedure in the multivariate analysis of all the patients. Surely this factor is unmodifiable and related to urgent surgery, therefore entailing a poorer prognosis.

Another datum of the technique that has not been highlighted by the literature on this subject is that the use of absorbable sutures for mesh fixation is related to a greater frequency of complications, and especially of recurrences. In contrast, monofilament sutures are associated with fewer complications and recurrences. Therefore, based on our data, we recommend using non-absorbable monofilament sutures for mesh fixation in these patients.

Finally, regarding the separate analysis of the different types of prosthetic mesh, our study confirms what has been demonstrated by other research<sup>15</sup> in reference to an increased frequency of recurrences with Physiomesh (Ethicon, NJ, USA).

In summary, several measures for improvement should be implemented in the treatment of patients with incisional hernia. Surgeons should insist on reducing patient BMI, referring patients to or creating specialized weight loss units for the global treatment of obesity. Patients should be very carefully evaluated in the emergency room to improve their status and ensure that they are referred to specialized units, if required. In view of the results, the creation of specific abdominal wall surgery units seems recommendable; in this manner, the negative effects of complex hernias can be attenuated and the use of technical resources for their treatment can be streamlined. This suggestion has also been included in other studies about abdominal wall surgery<sup>16</sup> and intestinal anastomoses.<sup>17</sup> In order to provide data on surgeons, it seems clear that we should incorporate parameters related to surgical training in the registry.

---

## Funding

B. Braun.

---

## Conflict of Interests

The authors have no conflict of interests to declare.

---

## Acknowledgements

Silvia Martínez and Xavier Masramón; SAIL (Servicio de Asesoría a la Investigación y Logística)

---

## Addendum. Members of the EVEREG Group

Antonio Utrera González, Hospital Juan Ramón Jiménez, Huelva

José M. Capitán Vallvey, Complejo Hospitalario de Jaén, Jaén

Matías Pradas, Hospital Comarcal de Ronda, Ronda, Málaga  
Daniel Pérez Gomar, Hospital Universitario Puerta del Mar, Cádiz

Manuel Bustos, Hospital Virgen del Rocío, Sevilla  
Francisco Antonio Herrera Fernández, Hospital Comarcal Santa Ana, Motril

Juan A. Blanco, Hospital Civil de Málaga  
Joaquim Robres Puig, Consorci Sanitari Integral de l'Hospitalet, Barcelona

Rafael Villalobos, Hospital Arnau de Vilanova, Lleida  
Dolors Güell Puigcercos, Hospital de Terrassa, Terrassa  
José María Puigcercos Fusté, Hospital Dos de Maig, Barcelona

Vicente Ayuso Osuna, Hospital Espíritu Santo, Santa Coloma Gramanet

Marta Piñol, Hospital Espíritu Santo, Santa Coloma Gramanet

Carles Olona Casas, Hospital Joan XXIII, Tarragona  
Francisco Martínez Ródenas, Hospital Municipal de Badalona, Badalona

Jordi Comajuncosas, Hospital Parc Sanitari Sant Joan de Déu, Sant Boi

Enric Sebastián Valverde, Hospital Parc Sanitari Sant Joan de Déu, Sant Boi

Sara Amador, Hospital de Granollers  
Jaume Gelonch, Hospital de Palamós  
Helena Vallverdú, Hospital de Vic  
Magda Abelló Sala, Hospital Pius, Valls

Antonio Veres Gómez, Hospital de la Cerdaña  
Rosa Gamón, Hospital General de Castellón  
Pedro García Peche, Hospital General Universitario Reina Sofía, Elche

Miguel González Valverde, Hospital General Universitario Reina Sofía, Elche

Miguel Ángel Martín Blanco, Hospital de Vinaroz  
Ramón J Ferri, Hospital Lluís Alcanyís, Valencia  
Elena Martí Cifrat, Hospital Malvarrosa, Valencia  
Providencia García Pastor, Hospital Universitario La Fe, Valencia

Carlos Montero, Hospital Virgen de los Lirios, Alcoy  
Miguel Ángel Lorenzo Liñán, Hospital General de Onteniente

Vicente Barbero, Complejo Hospitalario Virgen de la Salud, Toledo

Miguel Ángel Morlan, Complejo Hospitalario Virgen de la Salud, Toledo

José María Jover, Hospital de Getafe  
Isabel Delgado Lillo, Hospital Clínico San Carlos, Madrid  
María Mercedes Sanz Sánchez, Hospital General Universitario Gregorio Marañón, Madrid

Leire Zarain Obrador, Hospital General Universitario Gregorio Marañón, Madrid

Alejandro Lusilla, Hospital General Universitario Gregorio Marañón, Madrid

Juan Jesús Cabeza Gómez, Hospital Clínico San Carlos, Madrid

Antonio López, Hospital Nuestra Señora del Prado, Talavera de la Reina

Teresa Butrón, Hospital Universitario 12 de Octubre, Madrid

Guillermo Supelano Eslait, Hospital Universitario 12 de Octubre, Madrid

José Luis Álvarez Conde, Complejo Hospitalario de Palencia

Baltasar Pérez Saborido, Hospital Río Ortega, Valladolid

Mario Rodríguez López, Hospital Río Ortega, Valladolid

Santiago Revuelta Álvarez, Hospital Universitario Marqués de Valdecilla, Santander

Pedro Trillo Parejo, Complejo Hospitalario Universitario, Orense

José Luis Rodicio, Hospital Universitario Central de Asturias, Oviedo

Antonio Blanco, Hospital Universitario Central de Asturias, Oviedo

#### REFERENCES

1. Nilsson E, Haapaniemi S. Hernia registers and specialization. *Surg Clin North Am.* 1998;78:1141-55.
2. Nilsson E, Haapaniemi S, Gruber G, Sandblom G. Methods of repair and risk for reoperation in Swedish hernia surgery from 1992 to 1996. *Br J Surg.* 1998;85:1686-91.
3. Bay-Nielsen M, Kehlet H, Strand L, Malmstrøm J, Andersen FH, Wara P, et al., Danish Hernia Database Collaboration. Quality assessment of 26,304 herniorrhaphies in Denmark: a prospective nationwide study. *Lancet.* 2001;358:1124-8.
4. Israelsson LA, Smedberg S, Montgomery A, Nordin P, Spangen L. Incisional hernia repair in Sweden 2002. *Hernia.* 2006;10:258-61.
5. Helgstrand F, Rosenberg J, Bay-Nielsen M, Friis-Andersen H, Wara P, Jorgensen LN, et al. Establishment and initial experiences from the Danish Ventral Hernia Database. *Hernia.* 2010;14:131-5.
6. Bisgaard T, Kehlet H, Bay-Nielsen MB, Iversen MG, Wara P, Rosenberg J, et al. Nationwide study of early outcomes after incisional hernia repair. *Br J Surg.* 2009;96:1452-7.
7. Kokotovic D, Bisgaard T, Helgstrand F. Long-term recurrence and complications associated with elective incisional hernia repair. *JAMA.* 2016;316:1575-82.
8. Pereira JA, López-Cano M, Hernández-Granados P, Feliu X, en representación del grupo EVEREG. Resultados iniciales del Registro Nacional de Hernia Incisional. *Cir Esp.* 2016;94:595-602.
9. Slater NJ, Montgomery A, Berrevoet F, Carbonell AM, Chang A, Franklin M, et al. Criteria for definition of a complex abdominal wall hernia. *Hernia.* 2014;18:7-17.
10. Martínez-Serrano MA, Pereira JA, Sancho JJ, López-Cano M, Bombuy E, Hidalgo J, Study Group of Abdominal Hernia Surgery of the Catalan Society of Surgery. Risk of death after emergency surgery of abdominal wall hernias. Still waiting for improvement. *Langenbecks Arch Surg.* 2010;395:551-6.
11. Helgstrand F, Rosenberg J, Kehlet H, Jorgensen LN, Bisgaard T. Nationwide prospective study of outcomes after elective incisional hernia repair. *J Am Coll Surg.* 2013;216:217-28.
12. Liang Mk, Holihan JL, Itani K, Alawadi ZM, Gonzalez JR, Askenasy EP, et al. Ventral hernia management: expert consensus guided by systematic review. *Ann Surg.* 2017;265:80-9.
13. Kaoutzanis C, Leichtle SW, Mouawad NJ, Welch KB, Lampan RM, Wahl WL, et al. Risk factors for postoperative wound infections and prolonged hospitalization after ventral/incisional hernia repair. *Hernia.* 2015;19:113-23.
14. Owei L, Swendiman RA, Kelz RR, Dempsey DT, Dumon KR. Impact of body mass index on open ventral hernia repair: a retrospective review. *Surgery.* 2017;162:1320-9.
15. Köckerling F, Simon T, Hukauf M, Hellinger A, Fortelny R, Reinpold W, et al. The importance of registries in the postmarketing surveillance of surgical meshes. *Ann Surg.* 2017. <http://dx.doi.org/10.1097/SLA.0000000000002326> [in press].
16. Aquina CT, Fleming FJ, Becerra AZ, Xu Z, Hensley BJ, Noyes K, et al. Explaining variation in ventral and inguinal hernia repair outcomes: a population-based analysis. *Surgery.* 2017;162:628-39.
17. García-Granero E, Navarro F, Cerdán Santacruz C, Frasson M, García-Granero A, Marinello F, et al. Individual surgeon is an independent risk factor for leak after double-stapled colorectal anastomosis: an institutional analysis of 800 patients. *Surgery.* 2017;162:1006-16.