



## Image of the month

## Image-guided Excision With I<sup>125</sup> Seed in Lymph Node Metastasis of Medullary Thyroid Carcinoma <sup>☆</sup>



### Exéresis radioguiada con semilla de I<sup>125</sup> en recidiva metastásica ganglionar en carcinoma medular de tiroides

María Pérez Reyes,<sup>\*</sup> Carolina Jiménez Mazure, Isabel Pulido Roa, Cristina Rodríguez Silva

Servicio de Cirugía General, Aparato Digestivo y Trasplantes, Unidad de Cirugía Endocrina, Hospital Regional Universitario de Málaga, Málaga, Spain



Fig. 1.



Fig. 2.

A 60-year-old male underwent total thyroidectomy with central and right lateral lymph node dissection in 2006 for medullary carcinoma. In 2013, he had surgery for cervical recurrence. During follow-up in 2019, CT scan revealed a metastatic cervical lymphadenopathy in sector II (Fig. 1).

In the oncology committee, we decided to perform radio-guided surgery by implanting an I<sup>125</sup> capsule to confirm its position. After excision, the specimen was studied by x-ray (Fig. 2), and pathology results confirmed the metastasis.

The use of I<sup>125</sup> capsules in recurrent lesions is effective because of its simple placement and greater precision in intraoperative localization. Its use in recurrent medullary thyroid cancer had not been described to date.

<sup>☆</sup> Please cite this article as: Pérez Reyes M, Jiménez Mazure C, Pulido Roa I, Rodríguez Silva C. Exéresis radioguiada con semilla de I<sup>125</sup> en recidiva metastásica ganglionar en carcinoma medular de tiroides. Cir Esp. 2020;98:357.

<sup>\*</sup> Corresponding author.

E-mail address: [maria.perezreyes1991@gmail.com](mailto:maria.perezreyes1991@gmail.com) (M. Pérez Reyes).