

ORIGINAL ARTICLE

Postprandial lipemia induces endothelial dysfunction and higher insulin resistance in healthy subjects[☆]

Robinson Ramírez-Vélez^{a,b}^a School of Medicine, University of Valle, Cali, Colombia^b School of Medicine, University ICESI, Cali, Colombia

Received 16 May 2011; accepted 21 July 2011

Available online 13 January 2012

KEYWORDS

Endothelium;
Insulin resistance;
Lipids;
Adults

Abstract

Objective: To assess the effect of postprandial lipemia on endothelial function, insulin resistance, and lipid profile in healthy subjects.

Patients and methods: A prospective, interventional study in 14 healthy young men aged 18–25 years who were given a high-fat meal. Endothelial function was measured using flow-mediated dilation (FMD) in the brachial artery, flow velocity, mean arterial pressure and serum nitrite/nitrate levels (NO₂/NO₃). Glucose, insulin, total cholesterol, and triglyceride levels were also tested. Insulin resistance was determined by calculating the HOMA-IR index (Homeostatic Model Assessment-Insulin Resistance).

Results: Baseline FMD was 5.9 ± 1.1%. Postprandial lipemia reduced endothelial function by approximately 50% in the first (3.3 ± 0.5%, *p* = 0.03) and second (3.3 ± 0.4%, *p* = 0.04) moment, respectively. This finding was associated to an increased flow rate in the brachial artery and lower NO₂/NO₃ levels (*p* < 0.05). Higher cholesterol and triglyceride levels were found 1 h and 2 h postprandial (*p* < 0.05). HOMA-IR was significantly increased 1 h and 2 h postprandial (*p* < 0.05). **Conclusions:** Postprandial lipemia causes changes in circulating lipid profile and induces endothelial dysfunction and higher insulin resistance.

© 2011 SEEN. Published by Elsevier España, S.L. All rights reserved.

PALABRAS CLAVE

Endotelio;
Resistencia a la
insulina;
Lípidos;
Adultos

La lipemia pos-prandial induce disfunción endotelial y mayor grado de resistencia a la insulina en sujetos sanos

Resumen

Objetivo: Evaluar el efecto de la lipemia pos-prandial sobre la función endotelial, la resistencia a la insulina y el perfil de lípidos en sujetos sanos.

Materiales y métodos: Estudio de intervención prospectivo en 14 hombres jóvenes sanos entre los 18 y 25 años de edad a los que se administró una comida rica en grasas. La función endotelial se midió a través de vasodilatación mediada por flujo (VMF) en arteria braquial, la velocidad de flujo y los niveles séricos de nitritos/nitratos (NO₂/NO₃). Se evaluaron los niveles de glucosa,

[☆] Please cite this article as: Ramírez-Vélez R. La lipemia pos-prandial induce disfunción endotelial y mayor grado de resistencia a la insulina en sujetos sanos. Endocrinol Nutr. 2011;58:529–35.

E-mail address: robin640@hotmail.com

insulina, colesterol total y triglicéridos. La resistencia a la insulina se determinó mediante el cálculo del índice *homeostatic model assessment-insulin resistance* (HOMA-IR).

Resultados: El valor basal de la VMF fue de $5,9 \pm 1,1\%$. Se identificó que la lipemia pos-prandial reducía la función endotelial aproximadamente en 50%, en la primera ($3,3 \pm 0,5\%$, $p=0,03$) y segunda hora ($3,3 \pm 0,4\%$, $p=0,04$), respectivamente. Este hallazgo se acompañó de un incremento en la velocidad del flujo braquial, la presión arterial media y menores niveles de NO_2/NO_3 ($p < 0,05$). Se encontraron mayores niveles de colesterol total y triglicéridos a la 1 h y 2 h pos-prandial ($p < 0,05$). El HOMA-IR, se encontró significativamente elevado comparado a la 1 h y 2 h pos-prandial, ($p < 0,05$).

Conclusiones: La lipemia pos-prandial induce cambios en el perfil de lípidos circulantes e induce disfunción endotelial y mayor grado de resistencia a la insulina.

© 2011 SEEN. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

Epidemiological,^{1,2} clinical,^{3,4} and experimental^{5,6} trials have reported an association between postprandial metabolism of triglyceride rich lipoproteins and cardiovascular and metabolic diseases. Postprandial lipemia is a metabolic state characterized by an exaggerated increase in plasma triglyceride concentrations, due to an increase in their clearance after a high-fat intake.⁷ An increase in remnant chylomicrons (RCs) and very low density lipoprotein (VLDL) has also been observed during this period, stimulating the formation of low density lipoprotein (LDL), as well as a reduction in concentrations of high density lipoprotein (HDL).⁸

Over the long term, this metabolic state causes impaired vascular function and increased oxidative stress.^{9,10} Oxidative stress and endothelial dysfunction are considered significant risk factors for the development and progression of atherosclerosis¹¹ and both conditions have been associated with an increased risk of all-cause morbidity and mortality.¹² However, other authors have added that genetic¹³ and environmental¹⁴ factors, including decreased insulin sensitivity secondary to obesity and carbohydrate rich diets, could lead to increased baseline triglyceride levels, thereby also facilitating the development of cardiometabolic diseases.¹⁵

Another relevant aspect in the postprandial state is the effect of insulin and insulin sensitivity, considered important biomarkers of cardiovascular risk.^{16,17} Insulin facilitates the entry of free fatty acids, as well as their esterification and storage in the form of triglycerides in adipose tissue. This does not occur in situations of insulin resistance (IR), so that an inadequate quantity of free fatty acids circulates in the postprandial state to reduce insulin sensitivity and accentuate postprandial lipemia, while also reducing endothelial dependent vasodilation with resulting endothelial dysfunction.¹⁸ Taking into account that a large part of the human life cycle is in the postprandial state, this may be considered as a important metabolic factor in the study of cardiovascular risk. The aim of this study was to assess the effect of postprandial lipemia on endothelial function, insulin resistance and lipid profile in healthy subjects.

Methods

This was a prospective interventional study of 14 apparently healthy young men aged 18–25 years. The subjects included in the study belonged to a higher education institution in the city of Cali (Colombia) and were selected by consecutive sampling during the first quarter of 2011. Subjects with a history of smoking, surgery, or recent major trauma, endocrine, autoimmune, respiratory or heart disease (<1 month at study entry) and body mass index (BMI) greater than 28 kg/m^2 were excluded. In addition, a survey on nutritional habits, sociodemographic data and informed consent, following approval by the ethics committee (UV 10-09), was carried out. Anthropometric variables (weight, height, BMI, % fat) were measured according to the protocol described by Estrada¹⁹ and Ramírez-Vélez et al.²⁰ for the Colombian population.

After approximately a 12 h fast, 10 ml of blood was collected via puncture of the antecubital vein into tubes without additive for measurement of glucose, total cholesterol and triglycerides (TG) by direct colorimetric method in an automated spectrophotometer (Biosystems, Spain).²¹ Insulin levels were measured by a chemiluminescence assay (Immulite 1000 kit, San Jose, CA. USA).²² With these results, the index of insulin resistance was calculated (HOMA-IR: Homeostatic Model Assessment-Insulin Resistance) using the formula: $\text{HOMA-IR} = \text{fasting glucose (mmol/L)} \times \text{fasting insulin } (\mu\text{U/mL}) / 22.5$.²³ Measurement of serum levels of nitric oxide (NO) metabolites nitrite and nitrate (NO_2/NO_3) was performed by colorimetric assay (Cayman Chemical, USA),²⁴ as an indicator of in vivo NO production and bioavailability.

Endothelial function was evaluated using the flow-mediated vasodilation (FMD) test, a technique that includes visualization of the brachial artery, measurement of its diameter and blood flow velocity using vascular plethysmography Images.^{25,26} After 5 min of rest, the patient was placed in a supine position; in this position, the diameter of the brachial artery was measured both 1 min before and after it had been occluded with the sphygmomanometer cuff for 5 min (applying a pressure 60 mmHg above baseline systolic blood pressure). The 7.5 MHz transducer was placed at 60° with respect to the longitudinal axis of the

vessel and arterial flow velocity (cm s^{-2}) was measured. The image was frozen when it coincided with the QRS interval of the electrocardiogram and the intima–lumen interface was identified. The diameter and area of the artery (mm) were measured, and the skin on the arm was marked for subsequent placement of the transducer (in the same position) in subsequent measurements. FMV was expressed as the percent change in arterial diameter calculated with the formula $\text{FMV} = (\text{peak hyperemic diameter} - \text{baseline arterial diameter}) / \text{baseline arterial diameter} \times 100$. Endothelial shear stress (ESS) of the brachial artery (dynes cm^{-2}) was calculated using the formula: $\text{ESS} = (\text{Vm} D^{-1})$, where Vm = mean or peak blood velocity (cm s^{-1}) and D^{-1} = mean baseline brachial arterial diameter (cm).²⁷ During this measurement, blood pressure was recorded according to the recommendations of the American Heart Association²⁸ with an Omron® automatic sphygmomanometer. Mean arterial pressure was calculated using the formula: $(2 \times \text{diastolic blood pressure} + \text{systolic blood pressure}) / 3$. The effect of a high fat intake on vascular and cardiovascular markers was measured after fasting for 8–12 h (baseline), and 1 and 2 h post-intake, and all subjects received a standard diet low in nitrates 24 h before the tests.

To induce the postprandial lipemia state, a menu containing 1049 calories was prepared, containing 31 g protein, 79 g fat (31 g saturated fat), 666 mg cholesterol and 69 g carbohydrates (Ceres software version 1.02, 1997, FAO). A pilot study confirmed that this dietary intake led to a transient change in endothelial function.

The data were presented as mean \pm standard error of the mean (SEM) for metabolic and vascular variables, and mean \pm standard deviation (SD) for the characteristics of the subjects. The Shapiro–Wilk test was used to assess the normality of continuous data. Repeated measure analysis of variance (one-way ANOVA) with Dunnett's T3 post hoc test, a factorial analysis of variance or the Kruskal–Wallis test according to the distribution of the variables, was used to examine the differences between the different times. Statistical significance was assigned to a p -value ≤ 0.05 , and data were processed with the SPSS 15.0 software (SPSS Inc., Chicago, IL, USA).

Results

Table 1 summarizes the baseline characteristics of the participants. All parameters evaluated were within the range considered healthy for this age range. In the subjects evaluated, the value of FMV was $5.9 \pm 1.1\%$ at baseline measurement. It was determined that postprandial lipemia reduced endothelial function by approximately 50% in the first ($3.3 \pm 0.5\%$, $p = 0.03$) and second hours ($3.3 \pm 0.4\%$, $p = 0.04$), respectively (Fig. 1A). This finding was associated with a significant increase in brachial artery flow velocity (Fig. 1B), increasing from $22 \pm 1.2 \text{ cm s}^{-2}$ at baseline to $25.6 \pm 1.5 \text{ cm s}^{-2}$ at 1 h and $26.0 \pm 1.7 \text{ cm s}^{-2}$ at 2 h postprandial, respectively ($p < 0.05$). When testing the metabolism of NO, lower NO_2/NO_3 levels were observed 1 h ($74.4 \pm 4.4 \text{ mmol/L}$) and 2 h ($70.9 \pm 3.8 \text{ mmol/L}$, $p = 0.05$) postprandial vs baseline ($79.1 \pm 6.0 \text{ mmol/L}$) (Fig. 1C). A significant increase was also noted in mean, systolic

Table 1 Baseline characteristics of study subjects.

Characteristics	Mean \pm SD
Age (years)	21.0 \pm 2.8
Height (cm)	176.2 \pm 15.7
Weight (kg)	73.8 \pm 7.5
BMI (kg/m^2)	23.7 \pm 4.3
Body fat (%)	13.7 \pm 8.5
HR at rest (beats/min)	64.7 \pm 8.1
SBP (mmHg)	127.6 \pm 9.2
DBP (mmHg)	71.1 \pm 9.4
MBP (mmHg)	60.4 \pm 11.9
FMV (%)	5.9 \pm 1.1
Brachial diameter (mm)	3.5 \pm 0.2
Flow velocity (cm s^{-2})	23.3 \pm 5.5
Endothelial shear stress	0.09 \pm 0.04
Glucose (mmol/L)	4.6 \pm 0.4
Cholesterol (mmol/L)	4.6 \pm 1.0
Triglycerides (mmol/L)	1.6 \pm 1.5
Insulin ($\mu\text{U/mL}$)	9.5 \pm 4.7
HOMA-IR ($\text{mmol/L} \times \mu\text{U/mL}$)	1.9 \pm 0.9
NO_2/NO_3 (mmol/L)	69.3 \pm 15.6

HR, heart rate; HOMA-IR, Homeostatic Model Assessment-Insulin Resistance; BMI, body mass index; NO_2/NO_3 , nitrites/nitrates; DBP, diastolic blood pressure; SBP, systolic blood pressure; MBP, mean blood pressure; FMV, Flow mediated vasodilation.

and diastolic blood pressure 1 h postprandial ($p < 0.05$) (Fig. 2A–C).

No significant differences were found in glucose concentrations (Fig. 3A). As expected, higher cholesterol and triglyceride levels were found 1 h and 2 h postprandial ($p = 0.05$) (Fig. 3B and C). HOMA-IR was significantly increased 1 h and 2 h postprandial ($p = 0.05$) (Fig. 3D). Plasma insulin concentrations were elevated 1 h postprandial from baseline status ($7.6 \pm 1.2 \mu\text{U/mL}$ vs $28.5 \pm 5.7 \mu\text{U/mL}$, $p = 0.002$) (Fig. 3E).

Discussion

The main finding of this study was that postprandial lipemia causes changes in the circulating lipid profile and induces endothelial dysfunction and higher insulin resistance (IR), measured by the HOMA index. These results agree with prior studies suggesting an association between higher degrees of IR and an increased risk of cardiovascular events in subjects with characteristics similar to those in this study.^{29,30} Although the mechanism responsible for the reduction in vascular function and IR caused by postprandial lipemia is not known exactly, it is thought that elevated levels of triglyceride rich lipoproteins such as chylomicrons and VLDL, and their remnants, together with a state of oxidative stress are the main mechanisms that could explain this metabolic state.³¹

The marked reduction (50% reduction) in FMV in the first postprandial determination ($p = 0.03$) in this study was consistent with the results of De Koning and Rabelink,²⁹ Tsai et al.,³² and Kovacs et al.³³ However, the differences with other studies that did not find these same changes may be explained by the participants' age, the lipid composition of




Figure 1 Effect of postprandial lipemia on endothelial function. FMV, flow mediated vasodilation; NO₂/NO₃, nitrites/nitrates.

the diet administered, the time of vascular measurements and/or sex of the study subjects.^{10,13} Likewise, a similar finding was reported in diabetic and nondiabetic subjects after a high fat intake. For example, Tsai et al.³² observed that after a high fat meal—similar to that given in this study—the concentration of 8-PFG-2 α triglycerides (a marker of oxidative stress) was increased and GSH-Px activity (glutathione peroxidase) was decreased; these findings are consistent with the significant reduction in FMV in this study. In addition, it has been shown that vascular function improves after treatment with ciprofibrate³³ or the intake of supplements containing vitamin C and vitamin E.⁶

There is evidence that increased clearance of triglycerides in plasma can cause endothelial dysfunction and IR, a finding that was confirmed in this study.³⁴ Free fatty acids may lead to endothelial dysfunction due to increased superoxide anion (O₂⁻) production and a subsequent decrease in NO bioavailability,³⁵ as observed in the first and




Figure 2 Effect of postprandial lipemia on blood pressure.

second hour postprandial with NO₂/NO₃ plasma levels. Oxygen free radicals, such as O₂⁻, can react directly with NO forming peroxynitrites (ONOO⁻) and neutralizing the biological function of the latter, causing a loss in endothelium dependent vasodilation.³⁶ The accumulation of ONOO⁻ leads to an inhibition of eNOS (nitric oxide synthase; the enzyme responsible for NO production), generating changes in the vascular system that are reflected even in blood pressure levels, as was found in this study.

Multiple markers of IR have been studied as predictors of the risk of cardiovascular events, and their results are heterogeneous and discordant.^{23,37} The hyperinsulinemic euglycemic clamp is considered the reference test for assessing insulin sensitivity.³⁸ However, due to its technical difficulty, invasiveness and cost, it has not been considered as a tool applicable to large populations, which leads to the use of more practical alternatives for evaluation. HOMA-IR has been used as a clinical and epidemiological tool that can replace the clamp.³⁷ Thus, this model is a




Figure 3 Changes in lipid profile and insulin resistance induced by postprandial lipemia. HOMA-IR, *Homeostatic Model Assessment-Insulin Resistance*.

good tool for establishing IR. This study found significant increases in this evaluation parameter of IR. Although HOMA-IR, by definition, is used in the fasting state, its use in the postprandial state has been shown to account for some of the mechanisms by which IR attenuates postprandial hypertriglyceridemia. For example, decreased lipoprotein lipase (LPL) activity inhibits the clearance of postprandial triglycerides in individuals with IR.³⁹ On the other hand, IR causes less postprandial suppression of the release from the liver of triglyceride rich particles, such as VLDL that interfere with the insulin signaling pathways.⁴⁰

As mentioned above, the etiology of cardiometabolic diseases has not been fully established and it has been suggested that the postprandial state would partly explain the process leading to their development.¹⁻⁷ Recently, it has been shown that subjects who eat high fat diets have impaired endothelial function and higher levels of insulin, tumor necrosis factor- α and C-reactive protein,⁴ markers that have also been reported in patients with cardiovascular diseases associated with insulin resistance.⁴¹ All these mechanisms are the basis for the association between postprandial lipemia and IR.

In general, our results should be interpreted with caution due to the study limitations. First, the small number of subjects and the limited age range should be considered. We

did not consider the eating patterns and physical activity of the subjects that may have modulated their metabolic and vascular response. In conclusion, our results suggest that a high fat meal causes changes in the circulating lipid profile and induces endothelial dysfunction and a higher IR in healthy subjects. While not all studies have shown attenuation of FMV and a higher IR in the postprandial state, our findings may serve as a starting point for the preparation of further studies to investigate and confirm this metabolic phenomenon.

Funding

This study received financial and administrative support from the Program of Medicine of the ICESI University, Cali, Colombia, and the Laboratory of Proteins and Enzymes of Valle University, Cali, Colombia.

Conflicts of interest

During the preparation of this article, none of the authors had ties to activities that could create conflicts of interest.

References

- Miller M. The epidemiology of triglyceride as a coronary artery disease risk factor. *Clin Cardiol.* 1999;22:111–6.
- Castro Cabezas M, De Brain TWA, Jansen H, Kock LAW, Kortland W, Erkelens DW. Impaired chylomicron remnants clearance in familial combined hyperlipidemia. *Arterioscler Thromb.* 1993;13:804–14.
- Williams MJ, Sutherland WH, McCormick MP, De Jong SA, Walker RJ, Wilkins GT. Impaired endothelial function following a meal rich in used cooking fat. *J Am Coll Cardiol.* 1999;33:1050–5.
- Pereira MA, Kartashov AI, Ebbeling CB. Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet.* 2005;365:36–42.
- Cicha I, Suzuki Y, Tateishi N, Maeda N. Enhancement of red blood cell aggregation by plasma triglycerides. *Clin Hemorheol Microcirc.* 2001;24:247–55.
- Huang HY, Appel LJ, Croft KD, Miller ER, Mori TA, Puddey IB. Effects of vitamin C and vitamin E on in vivo lipid peroxidation: results of a randomized controlled trial. *Am J Clin Nutr.* 2002;76:549–55.
- Alipour A, Elte JW, van Zaanen HC, Rietveld AP, Castro Cabezas M. Novel aspects of postprandial lipemia in relation to atherosclerosis. *Atheroscler Suppl.* 2008;9:39–44.
- Bell DS, O'Keefe JH, Jellinger P. Postprandial dysmetabolism: the missing link between diabetes and cardiovascular events? *Endocr Pract.* 2008;14:112–24.
- Ogita H, Liao J. Endothelial function and oxidative stress. *Endothelium.* 2004;11:123–32.
- Plotnick GD, Corretti MC, Vogel RA. Effect of antioxidant vitamins on the transient impairment of endothelium-dependent brachial artery vasoactivity following a single high-fat meal. *JAMA.* 1997;278:1682–6.
- Stocker R, Kearney JR. Role of oxidative modifications in atherosclerosis. *Physiol Rev.* 2004;84:1381–478.
- Heitzer T, Schlinzig T, Krohn K, Meinertz T, Munzel T. Endothelial dysfunction, oxidative stress, and risk of cardiovascular events in patients with coronary artery disease. *Circulation.* 2001;104:2673–8.
- López-Miranda J, Pérez-Martínez P, Marín C, Moreno JA, Gómez P, Pérez-Jiménez F. Postprandial lipoprotein metabolism, genes and risk of cardiovascular disease. *Curr Opin Lipidol.* 2006;17:132–8.
- Lopez-Miranda J, Williams C, Lairon D. Dietary, physiological, genetic and pathological influences on postprandial lipid metabolism. *Br J Nutr.* 2007;98:458–73.
- O'Keefe JH, Bell DS. Postprandial hyperglycemia/hyperlipidemia (postprandial dysmetabolism) is a cardiovascular risk factor. *Am J Cardiol.* 2007;100:899–904.
- Matsuda M. Measuring and estimating insulin resistance in clinical and research settings. *Nutr Metab Cardiovasc Dis.* 2010;20:79–86.
- Singh B, Saxena A. Surrogate markers of insulin resistance: a review. *World J Diabetes.* 2010;1:36–47.
- Wats GF, Mamo JCL. Postprandial dyslipidaemia: new opportunities for prevention of coronary disease? *Br J Cardiol.* 1998;5:260–4.
- Estrada J. Parámetros antropométricos de la población laboral colombiana 1995 (acopl95). *Rev Fac Nac Salud Pública.* 1998;15:112–39.
- Ramírez-Vélez, López-Alban CA, Gallardo CE, Marmolejo LC. Características morfofuncionales de individuos físicamente activos. *Iatreia.* 2008;21:121–8.
- Kishi K, Ochiai K, Ohta Y. Highly sensitive cholesterol assay with enzymatic cycling applied to measurement of remnant lipoprotein-cholesterol in serum. *Clin Chem.* 2002;48:737–41.
- Grassi J, Pradelles P. Compounds labelled by the acetylcholinesterase of *Electrophorus Electricus*. Its preparation process and its use as a tracer or marker in enzyme-immunological determinations. United States patent, N° 1,047,330; September 10, 1991.
- Bonora E, Targher G, Alberiche M, Bonadonna RC, Saggiani F, Zenere MB, et al. Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: studies in subjects with various degrees of glucose tolerance and insulin sensitivity. *Diabetes Care.* 2000;23:57–63.
- Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite and (15N) nitrate in biological fluids. *Anal Biochem.* 1982;126:131–8.
- Pyke KE, Tschakovsky ME. The relationship between shear stress and flow-mediated dilatation: implications for the assessment of endothelial function. *J Physiol.* 2005;568:357–69.
- Corretti MC, Anderson TJ, Benjamin EJ. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol.* 2002;39:257–65.
- Padilla J, Johnson BD, Newcomer SC, Wilhite DP, Mickleborough TD, Fly AD, et al. Normalization of flow-mediated dilation to shear stress area under the curve eliminates the impact of variable hyperemic stimulus. *Cardiovasc Ultrasound.* 2008;4:44.
- Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation.* 2005;111:697–716.
- De Koning EJ, Rabelink TJ. Endothelial function in the postprandial state. *Atheroscler Suppl.* 2002;3:11–6.
- Arcaro G, Cretti A, Balzano S, Lechi A, Muggeo M, Bonora E, et al. Insulin causes endothelial dysfunction in humans: sites and mechanisms. *Circulation.* 2002;105:576–82.
- Westphal S, Taneva E, Kästner S, Martens-Lobenhoffer J, Bode-Böger S, Kropf S, et al. Endothelial dysfunction induced by postprandial lipemia is neutralized by addition of proteins to the fatty meal. *Atherosclerosis.* 2006;185:313–9.
- Tsai WC, Li YH, Lin CC, Chao TH, Chen JH. Effects of oxidative stress on endothelial function after a high-fat meal. *Clin Sci (Lond).* 2004;106:315–9.
- Kovacs I, Toldy E, Abel T, Tarjan J, Csaszar A. The effect of ciprofibrate on flow-mediated dilation and inflammatory markers in patients with combined hyperlipidemia. *Endothelium.* 2005;12:179–83.
- Hennig B, Alvarado A. Nutrition and endothelial cell integrity: implications in atherosclerosis. *Prog Food Nutr Sci.* 1993;17:119–57.
- Kusterer K, Pohl T, Fortmeyer HP, Marz W, Scharnagl H, Oldenburg A, et al. Chronic selective hypertriglyceridemia impairs endothelium dependent vasodilatation in rats. *Cardiovasc Res J.* 1999;42:783–93.
- Szabo C, Ischiropoulos H, Radi R. Peroxynitrite: biochemistry, pathophysiology and development of therapeutics. *Nat Rev Drug Discov.* 2007;6:662–80.
- Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care.* 2004;27:1487–95.
- DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol.* 1979;237:E214–23.

39. Panarotto D, Remillard P, Bouffard L, Maheux P. Insulin resistance affects the regulation of lipoprotein lipase in the postprandial period and in an adipose tissue-specific manner. *Eur J Clin Invest.* 2002;32:84–92.
40. Cardona F, Gonzalo-Marín M, Tinahones FJ. Relación de la hipertrigliceridemia posprandial con la resistencia a la insulina en pacientes con síndrome metabólico. *Endocrinol Nutr.* 2006;53:237–41.
41. Ramírez-Vélez R, Ortega JG, Hurtado C, Agredo-Zuñiga RA. Endothelial function is affected by postprandial lipemia: presentation of two cases and literature review. *Med UIS.* 2011;24:87–93.