

ORIGINAL ARTICLE

Relationship between glycated hemoglobin and glucose concentrations in the adult Galician population: selection of optimal glycated hemoglobin cut-off points as a diagnostic tool of diabetes mellitus

Manuel Antonio Botana López^{a,*}, Mónica López Ratón^b, María Ausencia Tomé^c, Alexis Fernández Mariño^d, José Antonio Mato Mato^e, Antonia Rego Iraeta^e, Román Pérez Fernández^f, Carmen Cadarso Suárez^b

^a Endocrine Section, Hospital Lucus Augusti, Lugo, Spain

^b Department of Statistics and Operations Research (Unit of Biostatistics), School of Medicine, University of Santiago de Compostela, Santiago de Compostela, Spain

^c Endocrine Service, University Clinical Hospital, Santiago de Compostela, Spain

^d Coya Specialty Centre, Vigo, Spain

^e Endocrine Service, Hospital Cristal-Piñor, Orense, Spain

^f Department of Physiology, School of Medicine, University of Santiago de Compostela, Santiago de Compostela, Spain

Received 13 January 2012; accepted 4 June 2012

Available online 1 August 2012

KEYWORDS

HbA_{1c};
Glucose;
Optimal HbA_{1c} cut-off points

Abstract

Aims/hypothesis: To analyze the relationship between glucose and glycated hemoglobin (HbA_{1c}) in the adult Galician population, evaluate the use of HbA_{1c} for the screening and diagnosis of diabetes, and calculate the diagnostic threshold required for this purpose.

Methods: We analyzed data on 2848 subjects (aged 18–85 years) drawn from a study undertaken in 2004 to assess the prevalence of diabetes in Galicia. For study purposes, diabetes was defined using the criteria recommended in 2002. Participants were classified into four glucose-based groups. The relationship between glucose and HbA_{1c} was described using linear regression models, generalized additive models and Spearman's correlation. Diagnostic capacity was assessed, and optimal HbA_{1c} cut-off points were calculated as a diabetes marker using the receiver operating characteristic curve.

Abbreviations: AUC, area under the curve; FPG, fasting glucose; 2hOGTT, glycemia at 2 h of oral glucose tolerance test; HbA_{1c}, glycated hemoglobin; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; KDM, known diabetes; NPV, negative predictive value; NGM, normal glucose metabolism; PPV, positive predictive value; ROC, receiver operating characteristic curve; Se, sensitivity; Sp, specificity; UKDM, unknown diabetes.

Please, cite this article as: Botana López MA, et al. Relationship between glycated hemoglobin and glucose concentrations in the adult Galician population: Selection of optimal glycated hemoglobin cut-off points as a diagnostic tool of diabetes mellitus. *Endocrinol Nutr.* 2012;59(8):496–504

* Corresponding author.

E-mail address: manuel.antonio.botana.lopez@sergas.es (M.A. Botana López).

Results: Prevalence of pre-diabetes, unknown diabetes and known diabetes was 20.86, 3.37 and 4.39%, respectively. The correlations between HbA_{1c} and fasting glucose were higher than those obtained for HbA_{1c} and glycemia at 2 h of the oral glucose overload (0.344 and 0.270, respectively). Taking glucose levels as the gold standard, a greater discriminatory capacity was obtained for HbA_{1c} (area under the curve: 0.839, 95% confidence intervals: 0.788–0.890). Based on the study criteria, the optimal minimum and maximum HbA_{1c} values were 5.9% and 6.7%, respectively.

Conclusions/interpretation: HbA_{1c} did not prove superior to glycemia for diagnosis of diabetes in the adult Galician population, and cannot therefore be used to replace the oral glucose tolerance test for screening and diagnosis purposes. Indeed, determination of glucose is essential to verify the diagnosis in the majority of cases.

© 2012 SEEN. Published by Elsevier España, S.L. All rights reserved.

PALABRAS CLAVE

HbA_{1c};
glucosa;
puntos de corte
óptimos de la HbA_{1c}

Relación entre hemoglobina glucosilada y concentraciones de glucosa en la población gallega adulta: selección de los puntos de corte óptimos de la hemoglobina glucosilada como herramienta diagnóstica de la diabetes mellitus

Resumen

Objetivos/hipótesis: Analizar la relación entre la glucosa y la hemoglobina glucosilada (HbA_{1c}) en la población gallega adulta, evaluar el uso de la HbA_{1c} para cribado y diagnóstico de la diabetes y calcular el umbral diagnóstico necesario para este fin.

Métodos: Se analizaron datos de 2.848 sujetos (de 18–85 años de edad) procedentes de un estudio emprendido en 2004 para valorar la prevalencia de diabetes en Galicia. A efectos del estudio, se definió la diabetes de acuerdo con los criterios recomendados en 2002. Se clasificó a los participantes en cuatro grupos en función de los valores de glucosa. Se describió la relación entre glucosa y HbA_{1c} mediante modelos de regresión lineal, modelos aditivos generalizados y la correlación de Spearman. Se valoró la capacidad diagnóstica y se calcularon los puntos de corte óptimos de la HbA_{1c} como marcador de la diabetes empleando la curva de características operativas del receptor.

Resultados: Las tasas de prevalencia de prediabetes, diabetes desconocida y diabetes conocidas eran del 10,86, 3,37 y 4,39%, respectivamente. Las correlaciones entre la HbA_{1c} y la glucemia en ayunas eran mayores que las obtenidas entre la HbA_{1c} y la glucemia en ayunas dos horas después de la sobrecarga oral de glucosa (0,344 y 0,270, respectivamente). Tomando los valores de glucosa como referencia, se obtuvo una mayor capacidad discriminatoria para la HbA_{1c} (área bajo la curva: 0,839, intervalos de confianza del 95%: 0,788–0,890). Basándose en los criterios del estudio, los valores óptimos mínimos de la HbA_{1c} eran del 5,9 y el 6,7%, respectivamente.

Conclusiones/interpretación: La HbA_{1c} no fue superior a la glucemia para el diagnóstico de la diabetes en la población gallega adulta, por lo que no puede utilizarse en lugar de la prueba de tolerancia oral a la glucosa con fines de cribado y diagnóstico. De hecho, la determinación de la glucosa es esencial para confirmar el diagnóstico en la mayoría de los casos.

© 2012 SEEN. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

Ascertaining the prevalence of diabetes is important because it is a disease that is becoming increasingly prevalent.¹ Fasting glucose (FPG) and glycemia at 2 h after an oral glucose overload test (2hOGTT) were classical and now also glycosylated hemoglobin (HbA_{1c}) are used for diagnosis of diabetes. Although considered the “gold standard” for diagnosis, measurement of blood glucose is subject to several limitations as patient must fast at least 8 h, it has a large biological variability, samples are not stable, numerous factors alter glucose concentrations (diurnal variation, sample source, acute illness or stress), and it reflects glucose homeostasis at a single point in time.²

HbA_{1c} has advantages including its familiarity to clinicians, convenience, preanalytic stability, and assay

standardization. It displays none of the variability inherent in the determination of glucose, it gives a better reflection of chronic hyperglycemia, and its concentration predicts the development of microvascular complications of diabetes.^{2,3} HbA_{1c} has a number of limitations: may be altered by factors other than glucose (e.g., change in erythrocyte life span, ethnicity), some conditions interfere with measurement (e.g., selected hemoglobinopathies), it may not be available in some laboratories/areas of the world and its cost is higher than glucose determination.²

HbA_{1c} cut-off point has been set at $\geq 6.5\%$ for diagnosis and at 5.7–6.4% for the diabetes high risk category.⁴ The question arises, however, as to whether HbA_{1c} identifies the same population as does glucose. Accordingly, the aim of this study was to analyze the relationship between glucose and HbA_{1c} in the adult Galician population, and evaluate

the performance of HbA_{1c} for the screening and diagnosis of diabetes.

Methods

The basic methodology of the study has been previously described.^{5,6} This study was carried out in a random sample representative of the Galician adult population (older than 18 years). Study subjects were selected by a two step cluster sampling procedure from the Galician Public Health Service (SERGAS) database, which covers more than 95% of the population. Primary health care centre dependent populations were randomly selected in each province (Galicia has four provinces; the population of each was considered as independent), and individual subjects aged over 18 years were then randomly selected within each population. Health centres were stratified by municipality type (rural or urban; coastal or interior); individuals were stratified by sex and age. Pregnant women were excluded from this study.

We contacted each person by mail in order to arrange the appointment for the study. For each non-responder a substitute was randomly selected. Information was collected through a personal interview at local health centres using a structured questionnaire, followed by a physical examination to measure blood pressure and anthropometric characteristics. Blood and urine samples were collected for subsequent analysis.

The study protocol was approved by the corresponding research ethics committee (Comité Ético de Investigación Clínica de Galicia). All participants signed informed consent forms.

The anthropometric measurements, including weight, height, waist circumference (WC) and hip circumference (HC), were obtained by trained personnel (physicians and nurses) using standardized techniques and equipment.⁷ Blood pressure was measured twice in recumbent position, with an interval of 3 min; the final value was the arithmetic mean of the two figures. Blood samples were drawn after a fasting period of 10–14 h.

Blood analyses were all done in the same central laboratory to which all samples were sent within the first 24 h after immediate centrifuging and freezing.

To evaluate glucose metabolism, 75 g of anhydrous glucose load was given orally in 250 ml of water to all subjects except those with known diabetes. Fasting and 2 h post-glucose load blood glucose were assessed by the glucose hexokinase method. HbA_{1c} was measured by HPLC. All other laboratory determinations were done using standardized procedures.

We used the diagnostic criteria recommended in 2002 by the Expert Committee on the Diagnosis and Classification of Diabetes mellitus⁸ to classify diabetes and lesser degrees of impaired glucose regulation, as follows (a) normal glucose, i.e., fasting plasma glucose (FPG) <100 mg/dL and 2hOGTT <140 mg/dL; (b) impaired fasting glucose (IFG): PG between 100 and 125 mg/dL; (c) impaired glucose tolerance (IGT): 2hOGTT between 140 and 199 mg/dL; and (d) diabetes: PG ≥126 mg/dL or 2hOGTT ≥200 mg/dL. For comparisons, we used also the criteria recommended in 2011.⁴ These included a new category of increased risk of diabetes, when HbA_{1c} is between 5.7% and 6.4%.

Statistical analysis

Based on their oral glucose tolerance test (OGTT) results, participants were classified into four different groups: (1) normal glucose (NGM); (2) prediabetes (subjects presenting with IFG, IGT or both); (3) unknown diabetes (UKDM); (4) known diabetes (KDM) (subjects who reported suffering from diabetes or using insulin or drugs for treatment of diabetes).

The correlations between HbA_{1c} and FPG, and between HbA_{1c} and 2hOGTT were calculated using Spearman correlations, which were not based on the assumption of normality, whether in the total study population or in the respective subgroups defined.

After patients with known diabetes were excluded, HbA_{1c} diagnostic capacity for diagnosis of diabetes was evaluated by using the receiver operating characteristic (ROC) curve and examining HbA_{1c} sensitivity (Se) and specificity (Sp) measures at different cut-off values, accompanied by the corresponding area under ROC curve (AUC) and its 95% confidence intervals.⁹ The AUC assumes values ranging from 0 to 1, such that the nearer its value approaches 1 the higher the discriminatory capacity.

Based on the ROC curve, optimal cut-off points that best discriminated between the diabetic and non-diabetic populations were then calculated on the basis of different criteria, namely: (1) judging the optimal cut-off point to be that at which the Se and Sp measures were similar or practically the same¹⁰; (2) choosing the optimal cut-off point as being that at which Se and Sp were simultaneously maximized,¹¹ thereby assuming that the consequences of false positives and false negatives were practically identical; (3) seeking the cut-off point which ensured that the Se and Sp values were the closest possible to 1 (criterion of the point closest to the point (0, 1) on the ROC curve)¹²; (4) taking the cut-off point that maximized Youden's index,¹³ which was equivalent to maximizing the sum of the Se and Sp measures; (5) selecting the cut-off point at which Sp was higher than or equal to a designated minimum value and, subject to this pre-condition, at which Se was as high as possible (to prevent a high number of false positives, in view of the high prevalence of diabetes); and (6) deeming the optimal cut-off point to be the value that maximized the percentage of correctly classified individuals or, what amounted to the same thing, that minimized the percentage of incorrect classifications of the diagnosis.¹⁴

To describe the effect of the different HbA_{1c} diagnostic thresholds obtained, we calculated the Se and Sp measures and the positive (PPV) and negative predictive values (NPV) at these cut-off values.

All statistical analyses were performed using the R 2.12.0 statistical software package, with a *p*-value of ≤0.05 considered as statistical significant.

Results

Study population characteristics

Of the 2860 patients included in the study, 2850 (99.65%) were aged between 18 and 85 years. As no HbA_{1c} values were available for two of the latter (0.07%), the analysis

initially included a total of 2848 participants. In this sample, 125 (4.4%) subjects had known diabetes and 96 (3.4%) had unknown diabetes, if we used only the blood glucose criteria. Using also the HbA_{1c} criterion, the number of subjects with unknown diabetes were 119 (4.2%). There were 23 subjects (19.3% of all subjects with diabetes) with HbA_{1c} \geq 6.5% and normal basal and 2h OGTT blood glucose. Seventeen subjects (14.3% with diabetes) had only met the criteria of fasting glucose, 34 (28.6%) met only the criterion of glucose at 2 h, and 21 (17.6%) subjects met the three diagnostic criteria. Forty-five (37.8%) subjects met at least two criteria. Five subjects had a basal glucose higher than 126 mg/dl with an HbA_{1c} higher than 6.5% who did not undergo a 2hOGTT.

If we consider IFG and IGT as prediabetes, there were 2033 subjects with normal glucose metabolism and 594 (20.9%) had prediabetes according to the ADA 2002 criteria. If we considered the "at risk" HbA_{1c} (between 5.7% and 6.4%) as another form of prediabetes, the number of subjects with prediabetes amounted 1600 (56.2%). A total of 1025 subjects had HbA_{1c} between 5.7% and 6.4% with normal basal and 2h OGTT glucose. These data are detailed in Table 1.

Table 2 shows the baseline characteristics of subjects classified according to the diagnostic criteria met. Body mass index, triglycerides and glucose concentrations (basal glucose, 2hOGTT and HbA_{1c}) were statistically different among subgroups. BMI was lower in patients diagnosed only by OGTT than in the other subgroups ($p < 0.05$ vs. diagnosed only by fasting glucose and $p < 0.01$ vs. those diagnosed only by HbA_{1c} or more than one criteria). There were no differences in the prevalence of obesity between subgroups. Triglyceride levels were significantly lower in patients diagnosed only after an OGTT or by HbA_{1c} compared with those diagnosed by at least two criteria ($p \leq 0.01$). Basal glucose levels were higher in those diagnosed only by fasting glucose and in those who had at least two criteria ($p < 0.001$ in all cases). The highest 2hOGTT concentrations were found in the group with at least two criteria ($p < 0.001$). This parameter was also higher in the group diagnosed only by this criterion compared with those diagnosed only by fasting glucose or by HbA_{1c} ($p < 0.001$). The highest HbA_{1c} concentration was that of the group with at least two criteria ($p < 0.001$). There were no differences between subgroups in the prevalence of hypertension, cholesterol, HDL cholesterol or microalbuminuria.

HbA_{1c}, FPG and 2hOGTT correlations

For the calculation of correlations and associations, patients who did not undergo a 2hOGTT (102 individuals, 3.58%) were excluded, leaving a total of 2746 patients for this analysis. Fitting a linear regression model yielded R^2 values of 0.46 for the association between FPG and HbA_{1c}, and 0.33 for that between 2hOGTT and HbA_{1c} (both significant at $p < 0.01$). Table 3 shows the Spearman correlations for HbA_{1c}, FPG and 2hOGTT. In the total population, the correlations between HbA_{1c} and FPG were higher than those between HbA_{1c} and 2hOGTT (0.344 and 0.270, respectively). The correlation between FPG and 2hOGTT levels was 0.445, i.e., slightly higher than those between HbA_{1c} and FPG.




Figure 1 Receiver operating characteristic curve for identification of participants with previously undiagnosed diabetes, using glycated hemoglobin and fasting plasma glucose for diagnosis and fasting plasma glucose and glycemia at 2h of oral glucose tolerance test as disease criteria. AUC: area under the curve; FPG: fasting plasma glucose; HbA_{1c}: glycated hemoglobin; ROC: receiver operating characteristic curve; Se: sensitivity; Sp: specificity.

Diagnostic capacity of HbA_{1c}

Fig. 1 depicts the ROC curve of HbA_{1c} as a tool for screening and diagnosis of diabetes. The AUC was 0.839 (95% CI 0.788–0.890). Table 4 shows the diagnostic properties of different HbA_{1c} cut-off values. As the cut-off level rose, Sp increased and Se decreased. The cut-off point at which Se and Sp were approximately equal (0.771 and 0.717, respectively) was 5.9% (95% CI 5.9–6.0), a value close to one standard deviation above mean HbA_{1c} in healthy subjects. Of the individuals with HbA_{1c} \geq 5.9%, only 9% had glucose levels the were diagnostic of diabetes. The optimal HbA_{1c} value which simultaneously maximized Se and Sp (0.729 and 0.824, respectively) and which corresponded to the point on the ROC curve closest to the point (0,1) was 6% (95% CI 5.9–6.0). Only 13% of patients with HbA_{1c} \geq 6% had glucose indicative of diabetes; 33% of pre-diabetic individuals had HbA_{1c} \geq 6%; and only 54% of subjects with UKDM had HbA_{1c} values \geq 6%.

The cut-off point that maximized the sum of Se (66.7%) and Sp (90.4%) was 6.1% (95% CI 6.0–6.2). Among the individuals with HbA_{1c} \geq 6.1%, 20% had glucose concentrations indicative of diabetes. This cut-off point would detect 69.4% of all subjects with UKDM. A total of 29% of participants at high risk of suffering diabetes (those with IFG or IGT) had HbA_{1c} \geq 6.1%.

The HbA_{1c} value at which Sp was greater than 95% and Se was at its highest possible level, was 6.3% (95% CI 6.2–6.3). This yielded a Se of 55.2% and a Sp of 96.8%. In this case, 39% of patients had glucose concentrations that were diagnostic of diabetes.

Table 1 Distribution of sample subjects according to the number of diagnostic criteria given.

	2010 criteria			2002 criteria	
	n	%		n	%
<i>NGM</i>	1004	35.3%	<i>NGM</i>	2033	71.4%
<i>Prediabetics</i>	1600	56.2%	<i>Prediabetics</i>	594	20.9%
Only IFG	112	3.9%	IFG	367	12.9%
Only IGT	47	1.7%	IGT	113	4%
Only HbA _{1c} ≥5.7 and ≤6.4	1025	36%	IFG and IGT	114	4%
IFG, IGT and HbA _{1c}	84	2.9%			
IFG and 2hOGTT (normal HbA _{1c})	19	0.7%			
IFG and HbA _{1c} (normal 2hOGTT)	249	8.7%			
IGT and HbA _{1c} (normal FPG)	64	2.2%			
<i>NKDM</i>	119	4.2%	<i>NKDM</i>	96	3.4%
Only elevated FPG	17	0.6%	Only elevated FPG	20	0.7%
Only elevated 2hOGTT	34	1.2%	Only elevated 2hOGTT	46	1.6%
Only elevated HbA _{1c}	23	0.8%	Only elevated HbA _{1c}		
Elevated FPG and 2hOGTT but not HbA _{1c}	4	0.1%	Elevated FPG and 2hOGTT	25	0.9%
Elevated FPG and HbA _{1c} but not 2hOGTT	3	0.1%			
Elevated 2hOGTT and HbA _{1c} but not FPG	12	0.4%			
Elevated FPG, 2hOGTT and HbA _{1c}	21	0.7%			
<i>KDM</i>	125	4.4%	<i>KDM</i>	125	4.4%
<i>Total</i>	2848		<i>Total</i>	2848	

HbA_{1c}: glycated hemoglobin; NGM: normal glucose metabolism; IFG: impaired fasting glucose; IGT: impaired glucose tolerance; FPG: fasting plasma glucose; 2hOGTT: plasma glucose value at 2 h in oral glucose tolerance test; KDM: previously known diabetics; NKDM: unknown diabetics.

The cut-off point that minimized the percentage of incorrect classifications was 6.7% (95% CI 6.5–7.0; Se 0.312; Sp 0.998; 83% of subjects with glucose concentrations that were diagnostic of diabetes). This cut-off value coincides exactly with that corresponding to 4 standard deviations above mean HbA_{1c} in normal subjects. Of all subjects having HbA_{1c} ≥6.5% and ≥7%, 64% and 92% had glucose concentrations diagnostic of diabetes, respectively. Of the total study population, however, only 2% had HbA_{1c} concentrations >6.5%, and 0.8% had HbA_{1c} concentrations >7%. In the present study, the cut-off value of 6.5% coincided with approximately three deviations above the mean HbA_{1c} of normal individuals (without diabetes and with a low risk of diabetes).

To compare the diagnostic properties of HbA_{1c} against those of FPG, the ROC curve was also calculated for FPG (see Fig. 1), using the criteria proposed in 2002. The AUC for FPG was 0.938 (95% CI 0.908–0.968), which was higher than that of HbA_{1c}.

Discussion

In this adult Galician population, we found a prevalence of diabetes of 7.8% using the ADA 2002 criteria (57% knew their condition but 43% had unknown diabetes) and a prevalence of 8.6% (51.2% KDM and 48.8% NKDM) according to the ADA 2010 criteria. This prevalence is clearly lower than that of 13.8% described recently for the whole of Spain.¹⁵ There are slight methodological differences between studies and we do not believe that they are responsible for this

difference. In the present study, there were almost no resignations for the performance of the OGTT while these occurred in a significant proportion of the study of Soriguer et al. In fact, probably the study in all Spain would have had more cases of diabetes if they had performed OGTT to the entire study population and in this case the difference with our sample would have been even greater. The only reason we find for explaining differences in both prevalences is the time elapsed between the studies. Distribution of known and unknown diabetes is the same in the Galician and Spain studies which could be due to the use of the same screening methods for detecting diabetes.

We had 20.9% of subjects classified as having prediabetes with the ADA 2002 criteria (12.9% only with IFG, 4% only with IGT and 4% with both IFG and IGT). These figures are different from those of the Di@bet.es Study (14.9% prediabetes: 3.4% IFG, 9.2% IGT, 2.2% combined IFG-IGT). It is surprising that having almost twice the prevalence of diabetes in the most recent study however the prevalence of prediabetes is lower and also has a different distribution of the types of abnormalities of glucose metabolism.

On the basis of a HbA_{1c} 5.7–6.4% criterion, 1025 individuals had prediabetes. These were a total of 1600 with this condition if we also included IFG and IGT. It is difficult to accept that more than half of the population has a metabolic disorder; thus, we believe that the HbA_{1c} in our population should not be used as a criterion for classifying a person as having prediabetes.

The correlations between HbA_{1c} and glucose concentrations are low in comparison with those obtained in patients with known diabetes, and a weak linear relationship is

Table 2 Baseline characteristics of subjects according to the way to diagnose diabetes.

	Total population	Total UKDM	BG	2h OGTT	HbA _{1c}	More than 1 criteria	KDM
<i>n</i> (%)	2848	119 (4.18)	17 (14.29)	34 (28.57)	23 (19.33)	45 (37.82)	125 (4.39)
Male sex (%)	1321 (46.38)	73 (61.34)	14 (82.35)	23 (67.65)	11 (47.83)	25 (55.56)	65 (52)
Age (years)	41.43 ± 15.01	56.68 ± 14.99	49.74 ± 11.91	55.45 ± 16.67	61.90 ± 11.20	57.57 ± 15.65	56.52 ± 15.45
Hypertension (%)	420 (14.78)	46 (38.66)	4 (23.53)	10 (29.41)	13 (56.52)	19 (42.22)	65 (52)
BMI (kg/m ²)	26.78 ± 4.93	29.72 ± 5.43	30.02 ± 4.83 [†]	27.62 ± 4.96 ^{*,‡,§}	30.73 ± 4.87 [†]	30.67 ± 5.94 [†]	29.58 ± 5.11 [†]
Obesity (%)	486 (17.06)	41 (34.45)	7 (41.18)	7 (20.59)	12 (52.17)	15 (33.33)	53 (42.40)
Total cholesterol (mg/dl)	195.68 ± 39.55	204.76 ± 40.27	205.76 ± 38.49	196.65 ± 45.22	201.22 ± 33.85	212.31 ± 39.84	199.22 ± 39.01
HDL cholesterol (mg/dl)	61.20 ± 16.78	54.53 ± 16.41	54.71 ± 25.75	58.18 ± 16.88	55.70 ± 13.77	51.11 ± 12.22	54.17 ± 15.74
Triglycerides (mg/dl)	80 (56–118)	113 (78.50–172.50)	111 (76.00–212.00)	96 (69.25–127.25) [§]	93 (75.50–171.50) [§]	159 (96.00–227.00) ^{†,‡}	100 (66–157) [§]
Microalbuminuria	2.5 (1.5–4.3)	3.35 (1.88–8.93)	2.8 (2.00–3.70)	3.2 (1.90–6.90)	3.6 (1.80–8.50)	4.4 (1.88–18.23)	4.20 (2.30–10.03)
FPG (mg/dl)	94.51 ± 21.54	131.19 ± 45.26	132.65 ± 5.69 ^{†,‡}	105.71 ± 11.30 ^{*,§}	106.70 ± 11.90 ^{*,§}	162.42 ± 59.17 ^{†,‡}	143.85 ± 50.00 ^{†,‡}
2-h postload glucose (mg/dl)	101.40 ± 41.47	221.11 ± 90.48	152.00 ± 28.10 ^{†,§}	228.32 ± 26.40 ^{*,‡,§}	144.57 ± 29.39 ^{†,§}	288.35 ± 110.01 ^{*,†,‡}	112.00 ± 51.37 ^{*,†,‡,§}
HbA _{1c} (%)	5.79 ± 0.60	6.72 ± 1.38	5.99 ± 0.28 ^{‡,§}	5.88 ± 0.32 ^{‡,§}	6.64 ± 0.20 ^{*,†,§}	7.67 ± 1.82 ^{*,†,‡}	7.10 ± 1.41 ^{*,†,§}

Values are mean ± SD, or percentage or medians (25th–75th percentiles). BMI: body mass index; HDL: high density lipoprotein; FPG: fasting plasma glucose; KDM: previously known diabetes; UKDM: unknown diabetes; BG: diabetes diagnosed only by basal glucose criterion; 2h OGTT: diabetes diagnosed only by 2h OGTT criterion; HbA_{1c}: diabetes diagnosed only by HbA_{1c} (glycated hemoglobin) criterion. Percentage of UKDM and KDM is from total sample. Other percentages are expressed from UKDM.

* Significantly different from diabetic subjects detected only by basal glucose

† Significantly different from diabetic subjects detected only by 2hOGTT.

‡ Significantly different from diabetic subjects detected only by HbA_{1c}.

§ Significantly different from diabetic subjects detected by at least two criteria.

Table 3 Spearman correlations between glycosylated hemoglobin, fasting plasma glucose and glycemia at 2 h of oral glucose tolerance test in the total population and in each subgroup.

	Total population	Pre-diabetic			
		NGM	IFG + IGT	UKDM	KDM
HbA _{1c} vs. FPG	0.34*	0.23*	0.19*	0.59*	0.65*
HbA _{1c} vs. 2hOGTT	0.27*	0.14*	0.15*	0.45*	0.60*
FPG vs. 2hOGTT	0.44*	0.28*	-0.21*	0.17*	0.84*

HbA_{1c}: glycosylated hemoglobin; FPG: fasting plasma glucose; 2hOGTT: glycemia at 2 h of oral glucose tolerance test; NGM: normal glycemia; IFG: impaired fasting glucose; IGT: impaired glucose tolerance; UKDM: unknown diabetes; KDM: known diabetes.

* Statistically significant values with a 5% significance level ($p < 0.05$).

observed between the them. These low correlations may indicate that HbA_{1c} and glucose reflect different metabolic conditions, essentially in the range of glucose tolerance values indicative of non-diabetic subjects. Furthermore, the degree of glycosylation is known to vary among individuals,¹⁶ an aspect that cannot be analyzed by this study, as glycemia was not monitored throughout the day. This might also indicate that these low HbA_{1c}-glucose correlations obtained could be due to variability of FPG and 2hOGTT in individuals.¹⁷ A wide range of average glucose levels for individuals with the same HbA_{1c} levels has been described, and this range is wider at the lowest HbA_{1c} levels and decreases with increasing HbA_{1c} concentrations.¹⁸ The correlation values observed in the current study were comparable to those reported by similar previous studies.¹⁹ The correlations between HbA_{1c} and glucose were higher in patients with known versus unknown diabetes. One explanation for this may lie in the degree of glycemic control, particularly among patients who are already receiving appropriate treatment with insulin or oral antidiabetics. One study²⁰ reported that, for any given HbA_{1c} level, the glycemic levels of patients in different treatment groups were not the same. The correlations observed by us among patients with known diabetes were similar to those reported

by Nathan et al.,¹⁸ who carried out glycemic monitoring in subjects with known diabetes, and Van't Riet et al.,¹⁹ who also obtained higher correlations among patients with known than among those with unknown diabetes.

Correlations depend on differences in the ranges of the variables studied, and tend to be lower in subgroups with narrower ranges. In our study, however, the correlations between glucose and HbA_{1c} were higher in subjects with diabetes than in the total population, but the ranges of the HbA_{1c} and glucose values were nevertheless wider in the total population. It must therefore be concluded that differences in the ranges of the variables do not constitute the only explanation for differences found in the correlations between the total population and the subgroups with diabetes.

In this study sample, FPG displayed a greater AUC than did HbA_{1c}, indicating that the diagnostic capacity of FPG is greater than that of HbA_{1c}. This could be related to the fact that glucose is used as the reference method to establish the criteria for diagnosing the disease. This result also agrees with those of a Dutch study¹⁹ and other studies which were also population-based and concluded that HbA_{1c} had no additional diagnostic value compared with FPG used in isolation, except in groups with a high risk of suffering diabetes.²¹

Table 4 Sensitivity, specificity, positive predictive value, and negative predictive value for diabetes mellitus using different glycosylated hemoglobin cut-off points. Brackets include 95% bootstrap confidence intervals for the cutoff points.

Criterion	HbA _{1c}	% Total population (without KDM) <i>n</i> = 2723	% Total population <i>n</i> = 2848	% High risk <i>n</i> = 594	Se	Sp	PPV	NPV
Se ≈ Sp	5.9 (5.9–6.0)	28.72	32.37	46.80	0.771	0.717	0.090	0.988
Maximizing Se and Sp simultaneously	6.0 (5.9–6.0)	18.68	22.19	33.33	0.729	0.824	0.132	0.988
Point under ROC closest to point (0,1)	6.0 (6.0–6.1)	18.68	22.19	33.33	0.729	0.824	0.132	0.988
Youden Index (maximizing Se + Sp)	6.1 (6.0–6.2)	11.10	14.40	22.56	0.667	0.904	0.203	0.987
Sp > 0.95 and maximizing Se	6.3 (6.2–6.3)	4.81	7.79	9.93	0.552	0.968	0.387	0.983
Recommended for diagnosis of diabetes	6.5	2.25	4.81	3.20	0.427	0.991	0.641	0.979
Maximizing % correct classifications	6.7 (6.5–7.0)	1.26	3.48	1.01	0.312	0.998	0.833	0.975

Se: sensitivity; Sp: specificity; PPV: positive predictive value; NPV: negative predictive value; HbA_{1c}: glycosylated hemoglobin; ROC: receiver operating characteristic curve; KDM: previously known diabetics.

In these groups HbA_{1c} may help to optimize the use of OGTT.²²

The HbA_{1c} value of 6% which simultaneously maximized Se and Sp in the current sample coincides exactly with the recently proposed cut-off point for screening individuals with a high risk of suffering diabetes. This would identify 40% of patients with intermediate glucose levels. The cut-off point that maximized Se and Sp was 6.1%.

An HbA_{1c} value $\geq 7\%$ yielded an Sp of close on 100% (99.9%), and 91% of subjects had glucose indicative of diabetes. A slightly higher cut-off point would have to be set in order to be able to dispense with any additional test for diagnosis of diabetes. The principal limitation of these diagnostic criteria is their low Se in return for a high Sp, something that is in line with the results of other studies.^{19,23} Accordingly, an OGTT would have to be performed to confirm the diagnosis in most cases.

Based on HbA_{1c}-glucose correlation values for the total population, the Se and Sp attained, and greater AUC for glucose, the use of HbA_{1c} instead of glucose values in the adult Galician population would not be advisable. Indeed, the advantages of using HbA_{1c} instead of the OGTT for the screening and diagnosis of diabetes mellitus are limited, and in most patients measurement of glucose will still be necessary to verify the diagnosis. Even so, despite its limitations, HbA_{1c} may be superior to OGTT in terms of cost-effectiveness and practical utility in the clinical setting. HbA_{1c} is less time-consuming than OGTT, can be measured at any time of day regardless of fasting, and can be analyzed with a small amount of the sample.²⁴ Furthermore, HbA_{1c} is a more complete measure of total glycemic exposure, inasmuch as it is indicative of glucose, not only in a fasting and but also in a postprandial state. The high correlation between HbA_{1c} and the presence of microvascular diabetic complications,²⁵⁻²⁸ and its association with cardiovascular diseases, even in the nondiabetic glucose-tolerance range of values²⁹ might indicate its usefulness as a diagnostic method among patients with a longer-term risk than that predicted by glucose.

The establishment of optimal cut-off points for HbA_{1c} in clinical practice calls for more in-depth research. There may even be different characteristics/covariates that influence the HbA_{1c} discriminatory capacity as a diagnostic marker of diabetes mellitus, and different optimal cut-off points would therefore have to be defined in accordance with the values of such covariates.

Our study could be completed after 5-10 years with a new cross-section on the same population and detect the patients that not having diabetes, but with levels of HbA_{1c} between 5 and 6.5%, develop diabetes after these years. As shown in the study published by Cheng et al.,³⁰ HbA_{1c} levels ≥ 5 increase the risk of diabetes after 4 years monitoring. Also another study³¹ proved that HbA_{1c} is a strong predictor of diabetes when it is within the limits 5 and 6.5%, and HbA_{1c} has also been shown as a good predictor of future diabetes in Spain.³²

In conclusion, the use of glycated hemoglobin as a criterion for diagnosing diabetes mellitus does not identify exactly the same subjects than the glucose criteria and we must have this in mind when we make this diagnosis. We believe that the maximum utility could be its use for initial screening but always confirming the existence of

diabetes using plasma glucose concentration. More studies are needed to establish in the long term the most useful criteria for identifying individuals with increased risk of morbidity and mortality.

Funding

M. López-Ratón and C. Cadarso-Suárez gratefully acknowledge the financial support of the Spanish Ministry of Science & Innovation (grants MTM2008-0163, MTM2010-09213-E and MTM2011-28285-C02-00).

Conflicts of interest

The authors have no conflicts of interest to declare.

References

1. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004;27:1047-53.
2. Sacks DB. A1C versus glucose testing: a comparison. *Diabetes Care*. 2011;34:518-23.
3. Nathan DM, Singer DE, Hurxthal K, Goodson JD. The clinical information value of the glycosylated hemoglobin assay. *N Engl J Med*. 1984;310:341-6.
4. The American Diabetes Association Clinical Practice Recommendations. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2011;34 Suppl. 1:S62-9.
5. Pérez-Fernández R, Mariño AF, Cadarso-Suárez C, Botana MA, Tomé MA, Solache I, et al. Prevalence, awareness, treatment and control of hypertension in Galicia (Spain) and association with related diseases. *J Hum Hypertens*. 2007;21:366-73.
6. Tomé MA, Botana MA, Cadarso-Suárez C, Rego-Iraeta A, Fernández-Mariño A, Mato JA, et al. Prevalence of metabolic syndrome in Galicia (NW Spain) on four alternative definitions and association with insulin resistance. *J Endocrinol Invest*. 2009;32:505-11.
7. World Health Organization. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee 1995. Technical Report Series 854. Geneva, Switzerland: World Health Organization.
8. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*. 2002;25 Suppl. 1: S5-20.
9. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics*. 1998;44:837-45.
10. Hosmer DW, Lemeshow S. *Applied logistic regression*. 2nd ed. Chichester, New York: Wiley; 2000.
11. Riddle DL, Stratford PW. Interpreting validity indexes for diagnostic tests: an illustration using the Berg Balance Test. *Phys Ther*. 1999;79:939-50.
12. Metz CE. Basic principles of ROC analysis. *Semin Nucl Med*. 1978;8:283-98.
13. Youden WJ. Index for rating diagnostic tests. *Cancer*. 1950;3:32-5.
14. Feinstein SH. The accuracy of diver sound localization by pointing. *Undersea Biomed Res*. 1975;2:173-84.
15. Sorriquer F, Goday A, Bosch-Comas A, Bordiú E, Calle-Pascual A, Carmena R, et al. Prevalence of diabetes mellitus and impaired glucose regulation in Spain: the Diabetes Study. *Diabetologia*. 2012;55:88-93.

16. Modan M, Meytes D, Rozeman P, Yosef SB, Sehayek E, Yosef B, et al. Significance of high HbA1 levels in normal glucose tolerance. *Diabetes Care*. 1988;11:422–8.
17. Selvin E, Crainiceanu CM, Brancati FL, Coresh J. Short-term variability in measures of glycemia and implications for the classification of diabetes. *Arch Intern Med*. 2007;167:1545–51.
18. Nathan DM, Kuenen J, Borg R, Zheng H, Schoenfeld D, Heine RJ. Translating the A1c assay into estimated average glucose values. *Diabetes Care*. 2008;31:1473–8.
19. Van't Riet E, Alsema M, Rijkelijhuizen JM, Kostense PJ, Nijpels G, Dekker JM. Relationship between A1C and glucose levels in the general Dutch population: the New Hoorn Study. *Diabetes Care*. 2010;33:61–6.
20. Kilpatrick ES, Rigby AS, Atkin SL. Variability in the relationship between mean plasma glucose and HbA1c: implications for the assessment of glycemic control. *Clin Chem*. 2007;53:897–901.
21. Perry RC, Shankar RR, Fineberg N, McGill J, Baron AD. HbA1c measurement improves the detection of type 2 diabetes in high-risk individuals with non diagnostic levels of fasting plasma glucose: the Early Diabetes Intervention Program (EDIP). *Diabetes Care*. 2001;24:465–71.
22. de la Hera JM, Vegas JM, Hernández E, Lozano I, García-Ruiz JM, Fernández-Cimadevilla OC, et al. Performance of glycated hemoglobin and a risk model for detection of unknown diabetes in coronary patients. *Rev Esp Cardiol*. 2011;64:759–65.
23. Rohlfing CL, Little RR, Wiedmeyer HM, England JD, Madsen R, Harris MI, et al. Use of GHb (HbA1c) in screening for undiagnosed diabetes in the U.S. population. *Diabetes Care*. 2000;23:187–91.
24. Voss EM, Cembrowski GS, Clasen BL, Spencer ML, Ainslie MB, Haig B. Evaluation of capillary collection system for HbA1c specimens. *Diabetes Care*. 1992;15:700–1.
25. McCance DR, Hanson RL, Charles MA, Jacobsson LTH, Pettitt DJ, Bennett PH, et al. Comparison of tests for glycosylated hemoglobin and fasting and two hour plasma glucose concentrations as diagnostic methods for diabetes. *Br Med J*. 1994;308:1323–8.
26. American Diabetes Association. Report of the Expert Committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care*. 1997;20:1183–97.
27. Diabetes Control Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med*. 1993;329:977–86.
28. U.K. Prospective Diabetes Study Group. Intensive blood-glucose control with sulphonyl ureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*. 1998;352:837–53.
29. Khaw KT, Wareham N, Bingham S, Luben R, Welch A, Day N. Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. *Ann Intern Med*. 2004;141:413–20.
30. Cheng P, Neugaard B, Foulis P, Conlin PR. Hemoglobin A1c as a predictor of incident diabetes. *Diabetes Care*. 2011;34:610–5.
31. Bonora E, Kiechl S, Mayr A, Zoppini G, Targher G, Bonadonna RC, et al. High-normal HbA1c is a strong predictor of type 2 diabetes in the general population. *Diabetes Care*. 2011;34:1038–40.
32. Valdés S, Botas P, Delgado E, Alvarez F, Díaz-Cadorniga F. HbA(1c) in the prediction of type 2 diabetes compared with fasting and 2-h post-challenge plasma glucose: the Asturias study (1998–2005). *Diabetes Metab*. 2011;37:27–32.