



ORIGINAL ARTICLE

High prevalence of obesity in a Spanish working population[☆]

Alberto Goday-Arnó^{a,b,c,*}, Eva Calvo-Bonacho^d, Miguel-Ángel Sánchez-Chaparro^{d,e}, José-Antonio Gelpí^d, Juan-Carlos Sainz^d, Sonia Santamaría^d, Rosa-Isabel Navarro^d, Faustino Gutiérrez^d, Carlos Sanz^d, Elena Caveda^f, Jesús Reviriego^f

^a Servicio de Endocrinología y Nutrición, Hospital del Mar, Barcelona, Spain

^b Departamento de Medicina, Universidad Autónoma de Barcelona, Barcelona, Spain

^c Cardiovascular Risk and Nutrition Research Group, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), Barcelona, Spain. CIBER de Fisiopatología de la Obesidad y Nutrición (CIBEROBN)

^d Ibermutuamur, Mutua de Accidentes de Trabajo y Enfermedades Profesionales de la Seguridad Social N.º 274, Madrid, Spain

^e Departamento de Medicina Interna, Hospital Universitario Virgen de la Victoria, Málaga, Spain

^f Departamento de Investigación Clínica, Lilly Spain, Alcobendas, Madrid, Spain

Received 18 September 2012; accepted 13 October 2012

Available online 3 June 2013

KEYWORDS

Prevalence;
Obesity;
Overweight;
Morbid obesity;
Occupation

Abstract

Background and objectives: To report the prevalence of obesity in a Spanish working population and its changes in recent years.

Material and methods: Data were collected from routine medical examinations performed on workers by a national mutual insurance society for occupational accidents and diseases (Ibermutuamur). A structured questionnaire was completed and physical examinations were performed. Overweight was defined as BMI ranging from 25 to 29.9, obesity as BMI of 30–39.9, and morbid obesity as BMI $\geq 40 \text{ kg/m}^2$.

Results: Data from 1,336,055 medical examinations performed from May 2004 to November 2007 were collected. Prevalence rates in the population examined in 2004 ($n=230,684$; 73% males; average age, 36.4 years) were: morbid obesity, 0.5% (0.6% males, 0.5% females); obesity, 14.5% (17.0% males, 7.7% females); overweight, 38.4% (44.8% males, 21.3% females). Prevalence rates of obesity and overweight were higher in blue-collar workers (16.4% and 40.5% respectively) as compared to white-collar workers (10.9% and 34.4% respectively). There was a progressive increase in prevalence of obesity during the 4-year study (2004–2007) in both males (17.0%, 17.6%, 17.9%, 18.2%) and females (7.6%, 8.0%, 8.4%, 8.7%).

Conclusions: Prevalence of obesity and overweight in the Spanish working population is high, especially in male blue-collar workers, and is increasing. There is a need to promote early prevention programs and specific treatments for obesity.

© 2012 SEEN. Published by Elsevier España, S.L. All rights reserved.

[☆] Please cite this article as: Goday-Arnó A, et al. Alta prevalencia de obesidad en una población laboral en España. Endocrinol Nutr. 2013;60:173–8.

* Corresponding author.

E-mail address: Agoday@hospitaldelmar.cat (A. Goday-Arnó).

PALABRAS CLAVE
 Prevalencia;
 Obesidad;
 Sobrepeso;
 Obesidad mórbida;
 Ocupación

Alta prevalencia de obesidad en una población laboral en España

Resumen

Antecedentes y objetivo: Describir la prevalencia de la obesidad y su evolución reciente en una población laboral en España.

Material y método: Se recogieron los datos de los chequeos médicos rutinarios practicados a trabajadores por la Mutua de Accidentes de Trabajo y Enfermedades Profesionales de la Seguridad Social (Ibermutuamur). Se realizó exploración física y se empleó un formulario estructurado para recoger los datos. Se consideró que un individuo tenía sobrepeso si su índice de masa corporal (IMC) estaba entre 25-29,9, obesidad entre 30-39,9 y obesidad mórbida si era de 40 kg/m² o mayor.

Resultados: Se incluyeron datos de 1.336.055 chequeos médicos realizados entre mayo de 2004 y noviembre de 2007. En los trabajadores examinados en 2004 (n = 230.684; 73% hombres; edad promedio, 36,4 años) las prevalencias fueron: obesidad mórbida: 0,5% (0,6% hombres, 0,5% mujeres), obesidad: 14,5% (17,0% hombres, 7,7% mujeres), sobrepeso: 38,4% (44,8% hombres, 21,3% mujeres); las prevalencias de obesidad y sobrepeso fueron mayores entre los trabajadores manuales (16,4 y 40,5%, respectivamente) que entre los trabajadores intelectuales (10,9 y 34,4%, respectivamente). Se observó un incremento progresivo en la prevalencia de obesidad a lo largo de los 4 años evaluados (2004-2007), tanto en hombres (17,0, 17,6, 17,9 y 18,2%) como en mujeres (7,6, 8,0, 8,4 y 8,7%).

Conclusiones: La prevalencia de obesidad y sobrepeso en una población laboral en España es alta, especialmente entre los trabajadores manuales varones, y va en aumento. Es necesario promover programas de prevención temprana y tratamientos específicos para la obesidad.

© 2012 SEEN. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Introduction

Obesity represents an increasing health problem because of the continuous increase in its worldwide incidence during the last decades and its impact on morbidity, mortality, quality of life, and healthcare costs.^{1,2} Lifestyle changes and decreased physical activity are determinant factors of the global impact of obesity.³ Although leisure time physical activity has slightly increased in industrialized countries, current patterns, as well as long-term trends, suggest that an effective decrease of physical activity in general, and specifically of work-related activity is occurring.^{2,4}

Obesity prevalence in Spain is high⁵ and shows an upward trend, as occurs in all other industrialized countries.^{2,6} The available epidemiological data come from local⁷ or regional⁸⁻¹² studies and from surveys with anthropometric data provided by the surveyed subjects themselves,¹³ who often underestimate their weight and overestimate their height. The reported obesity data may therefore be lower than the actual data.¹⁴ Moreover, the working population is often underrepresented in epidemiological studies conducted in the general population because people are less willing to participate in studies that require them to devote time during working hours.

This study analyzes the prevalence of obesity in a Spanish working population in order to collect up-to-date data and to report its prevalence as a function of the different types of occupation. Data collected during routine medical check-ups performed on active workers throughout Spain for 4 consecutive years allowed these objectives to be explored.

Subjects and methods

Study design

This was an observational, cross-sectional, epidemiological study including data from routine medical check-ups given to active workers between 2004 and 2007. The sample consisted of workers whose employers had agreements for the provision of health services with Ibermutuamur, a mutual insurance society for occupational accidents and diseases that complements the social security system in Spain. Ibermutuamur covers more than 1,100,000 workers from all Spanish regions, and its Prevention Society performs more than 400,000 medical check-ups every year. This study is part of the Ibermutuamur CArdiovascular Risk Assessment (ICARIA) plan, which has been applied in all medical check-ups performed at the facilities of the Ibermutuamur Prevention Society since 2004.¹⁵

Study population and measurements

Workers who underwent routine medical check-ups performed by the Ibermutuamur Prevention Society between May 2004 and November 2007 were enrolled into the study. When more than one medical check-up was recorded for the same year, only data from the oldest check-up (the one with the earliest date) were considered. A structured questionnaire was used to transfer data from the clinical records. The physical examination included direct measurements of weight and height (in light clothes and without shoes) to calculate body mass index (BMI), an estimator of body fat mass.

Workers were classified by BMI into the following categories:¹⁶

- Normal weight, BMI < 25 kg/m².
- Overweight, BMI ≥ 25 kg/m² and <30 kg/m².
- Obesity, BMI ≥ 30 kg/m².
- Morbid obesity, BMI ≥ 40 kg/m².

As regards occupation, workers were classified into two main occupational categories based on the Spanish 1994 National Classification of Occupations:¹⁷

- Manual or blue-collar workers, including those in the restaurant industry and in personal services, protection and security services; shop assistants and similar occupations; craftsmen; skilled workers in the farming and fishing industries; skilled workers in the manufacturing, construction, and mining industries; plant and machinery operators; assembly and non-skilled workers.
- Clerical or white-collar workers, including managers from public and private organizations; intellectuals; scientific and technical support staff, and clerks.

This study was conducted in compliance with the principles of the Declaration of Helsinki. The ethics committee of Ibermutuamur approved the study, and each worker signed an informed consent authorizing the use of their data. Data confidentiality was guaranteed in accordance to the applicable Spanish law on personal data protection.

Statistical analysis

Data were analyzed using statistical SAS software (SAS Institute Inc., Cary, NC Carolina, USA). Categorical variables are given as percentages with 85% confidence intervals (CIs). Continuous variables are given as mean and standard deviation (SD).

Results

Data from the 1,395,398 medical check-ups done between 2004 and 2007 were analyzed. Check-ups with incomplete clinical data on sex, age, occupational information, or BMI were excluded. The first cohort included 230,684 workers examined in 2004. The next cohorts included 380,996 workers examined in 2005; 389,681 in 2006, and 394,037 in 2007.

In the 2004 cohort, 72.9% of subjects were men, the mean age was 36.4 years, and approximately two thirds were manual workers. Their sociodemographic and clinical characteristics are shown in Table 1. The proportions of men and subjects in the younger age groups were greater in this sample than in a recent active population survey in Spain, in which 55.7% were men.¹⁸

The prevalence of obesity, including morbid obesity, was 14.9% (17.6% and 8.2% in men and women respectively). The prevalence of overweight was 38.4% (44.8% in men and 21.3% in women). A progressive increase in the prevalence of obesity and overweight was seen with age (Table 2). A continued increase in the prevalence of obesity was also seen during the four-year period analyzed. Such increases occurred both in men and women (Table 3). Only women experienced a

Table 1 Sociodemographic and clinical characteristics of subjects enrolled into the 2004 cohort.

Characteristics	n = 230,684
<i>Age (years), mean [SD]</i>	36.4 (10.9)
<i>Weight (kg), mean (SD)</i>	74.9 (14.8)
<i>Height (m), mean (SD)</i>	1.7 (0.1)
<i>BMI (kg/m²), mean (SD)</i>	25.8 (4.3)
<i>Abdominal circumference (cm), mean (SD)</i>	88.2 (13.0)
<i>Sex</i>	
Male, % (95% CI)	72.9 (72.7-73.1)
Female, % (95% CI)	27.1 (26.9-27.3)
<i>Occupation</i>	
Manual, % (95% CI)	65.2 (65.0-65.4)
Clerical, % (95% CI)	34.8 (34.6-35.0)
<i>Age group</i>	
<30 years, % (95% CI)	31.5 (31.3-31.7)
30-39 years, % (95% CI)	32.5 (32.3-32.7)
40-49 years, % (95% CI)	21.9 (21.7-22.0)
50-59 years, % (95% CI)	11.7 (11.6-11.9)
≥60 years, % (95% CI)	2.4 (2.4-2.5)

SD: standard deviation; CI: confidence interval; BMI: body mass index.

gradual increase in the prevalence of overweight over the four-year period (Table 3).

By type of occupation, the prevalence rates of obesity (including morbid obesity) and overweight were higher in manual workers (17.0% and 40.5% respectively) than in clerical workers (11.3% and 34.4% respectively) (Table 3). These differences were also seen in women from all age groups (Table 2). The prevalence of obesity was higher among men with manual occupations as compared to those in clerical work; however, overweight was less prevalent in men with manual occupations than in those in clerical work (Table 2). During the four years of the study, the prevalence of obesity increased in parallel in both manual and clerical workers (Table 3). The prevalence of overweight remained stable in both occupational groups during the four-year period (Table 3).

The prevalence of morbid obesity in the 2004 cohort was 0.5%. Table 2 shows these results broken down by age and sex. The prevalence of morbid obesity was higher in men as compared to women, with a progressive increase with age up to the range of ≥60 years, when a decrease was seen (Table 2). Morbid obesity was more prevalent in manual than in clerical workers (Table 2). During the four-year study period, the prevalence of morbid obesity increased slightly (Table 3).

Discussion

The prevalence of obesity and overweight in the Spanish working population is high and continues to increase. More than half the study workers were overweight or obese. The prevalence of obesity is lower in women as compared to men, and is also lower in clerical workers than in manual workers.

Table 2 Prevalence of overweight and obesity by age, sex, and type of occupation in the 2004 cohort.

	Males			Females			Total (n = 230,684)
	Clerical (n = 42,740)	Manual (n = 125,418)	Total (n = 168,158)	Clerical (n = 37,459)	Manual (n = 25,067)	Total (n = 62,526)	
<i>Overweight</i>							
<30 years	37.5 (36.5–38.4)	33.1 (32.6–33.5)	33.9 (33.5–34.3)	13.0 (12.4–13.6)	16.5 (15.7–17.3)	14.4 (14.0–14.9)	28.0 (27.7–28.4)
30–39 years	48.2 (47.4–49.0)	46.7 (46.2–47.2)	47.1 (46.7–47.5)	17.4 (16.8–18.0)	23.5 (22.5–24.5)	19.4 (18.9–19.9)	39.0 (38.7–39.4)
40–49 years	53.5 (52.5–54.4)	49.4 (48.8–50.0)	50.5 (50.0–51.0)	25.7 (24.6–26.7)	32.0 (30.8–33.2)	28.6 (27.8–29.3)	44.9 (44.5–45.3)
50–59 years	55.4 (54.1–56.6)	51.2 (50.4–51.9)	52.3 (51.6–52.9)	33.4 (31.5–35.4)	41.3 (39.5–43.2)	37.7 (36.4–39.1)	49.6 (49.0–50.2)
≥60 years	54.9 (52.3–57.6)	52.7 (51.0–54.4)	53.3 (51.9–54.8)	44.6 (39.1–50.0)	43.6 (39.1–48.2)	44.0 (40.5–47.5)	52.0 (50.7–53.3)
Total	48.2 (47.7–48.7)	43.6 (43.3–43.9)	44.8 (44.5–45.0)	18.6 (18.2–19.0)	25.3 (24.8–25.8)	21.3 (21.0–21.6)	38.4 (38.2–38.6)
<i>Obesity</i>							
<30 years	9.1 (8.5–9.7)	10.3 (10.0–10.6)	10.0 (9.8–10.3)	3.9 (3.5–4.2)	6.0 (5.5–6.5)	4.8 (4.5–5.0)	8.4 (8.2–8.6)
30–39 years	13.5 (13.0–14.1)	17.5 (17.1–17.9)	16.3 (16.0–16.7)	5.1 (4.7–5.4)	8.8 (8.2–9.5)	6.3 (6.0–6.6)	13.4 (13.2–13.6)
40–49 years	19.0 (18.3–19.8)	22.1 (21.6–22.6)	21.3 (20.9–21.7)	7.3 (6.7–7.9)	14.1 (13.3–15.0)	10.4 (9.9–10.9)	18.5 (18.1–18.8)
50–59 years	23.2 (22.1–24.3)	26.0 (25.3–26.7)	25.2 (24.7–25.8)	10.7 (9.5–12.0)	22.0 (20.4–23.5)	16.8 (15.8–17.9)	23.7 (23.2–24.2)
≥60 years	25.8 (23.5–28.1)	26.0 (24.5–27.5)	26.0 (24.7–27.2)	13.1 (9.4–16.8)	30.0 (25.8–34.3)	23.0 (20.1–26.0)	25.6 (24.4–26.7)
Total	15.6 (15.2–15.9)	17.5 (17.3–17.7)	17.0 (16.8–17.2)	5.5 (5.3–5.7)	10.9 (10.5–11.3)	7.7 (7.4–7.9)	14.5 (14.3–14.6)
<i>Morbid obesity</i>							
<30 years	0.3 (0.2–0.4)	0.4 (0.3–0.5)	0.4 (0.3–0.4)	0.3 (0.2–0.4)	0.3 (0.2–0.4)	0.3 (0.2–0.4)	0.4 (0.3–0.4)
30–39 years	0.4 (0.3–0.5)	0.7 (0.6–0.7)	0.6 (0.5–0.7)	0.3 (0.2–0.4)	0.6 (0.4–0.8)	0.4 (0.3–0.5)	0.5 (0.5–0.6)
40–49 years	0.6 (0.5–0.8)	0.8 (0.7–0.9)	0.7 (0.7–0.8)	0.5 (0.3–0.6)	0.9 (0.7–1.1)	0.7 (0.5–0.8)	0.7 (0.7–0.8)
50–59 years	0.5 (0.4–0.7)	0.7 (0.6–0.8)	0.7 (0.6–0.8)	0.8 (0.4–1.1)	1.3 (0.9–1.7)	1.0 (0.8–1.3)	0.7 (0.6–0.8)
≥60 years	0.5 (0.1–0.9)	0.4 (0.2–0.6)	0.4 (0.3–0.6)	0.3 (0.0–0.9)	0.9 (0.0–1.7)	0.6 (0.1–1.2)	0.5 (0.3–0.7)
Total	0.5 (0.4–0.5)	0.6 (0.6–0.6)	0.6 (0.5–0.6)	0.3 (0.3–0.4)	0.7 (0.6–0.8)	0.5 (0.4–0.5)	0.5 (0.5–0.6)

Values are given as percentages (95% CI).

Table 3 Prevalence of overweight and obesity by sex and type of occupation during the four years analyzed.

	2004 (n = 230,684)	2005 (n = 380,996)	2006 (n = 389,681)	2007 (n = 394,037)
<i>Overweight</i>				
Males	44.8 (44.5–45.0)	44.9 (44.7–45.0)	44.7 (44.5–44.9)	44.6 (44.4–44.8)
Females	21.3 (21.0–21.6)	22.0 (21.7–22.2)	22.3 (22.1–22.6)	22.9 (22.6–23.1)
White collar	34.4 (34.1–34.7)	34.1 (33.9–34.4)	34.3 (34.0–34.5)	33.9 (33.7–34.2)
Blue collar	40.5 (40.3–40.8)	41.0 (40.8–41.2)	40.9 (40.7–41.1)	40.9 (40.7–41.1)
Total	38.4 (38.2–38.6)	38.6 (38.4–38.7)	38.5 (38.4–38.7)	38.5 (38.3–38.6)
<i>Obesity</i>				
Males	17.0 (16.8–17.2)	17.6 (17.5–17.8)	17.9 (17.8–18.1)	18.2 (18.1–18.4)
Females	7.7 (7.4–7.9)	8.0 (7.8–8.1)	8.4 (8.2–8.6)	8.7 (8.6–8.9)
White collar	10.9 (10.6–11.1)	11.2 (11.1–11.4)	11.5 (11.3–11.6)	11.6 (11.4–11.7)
Blue collar	16.4 (16.2–16.5)	17.0 (16.9–17.2)	17.5 (17.3–17.6)	17.7 (17.5–17.8)
Total	14.5 (14.3–14.6)	15.0 (14.9–15.1)	15.3 (15.2–15.4)	15.5 (15.4–15.7)
<i>Morbid obesity</i>				
Males	0.6 (0.5–0.6)	0.6 (0.6–0.6)	0.7 (0.6–0.7)	0.7 (0.7–0.7)
Females	0.5 (0.4–0.5)	0.6 (0.5–0.6)	0.6 (0.5–0.6)	0.6 (0.6–0.7)
White collar	0.4 (0.4–0.4)	0.5 (0.4–0.5)	0.5 (0.5–0.5)	0.5 (0.5–0.6)
Blue collar	0.6 (0.6–0.6)	0.6 (0.6–0.7)	0.7 (0.7–0.8)	0.8 (0.7–0.8)
Total	0.5 (0.5–0.6)	0.6 (0.6–0.7)	0.6 (0.6–0.7)	0.7 (0.7–0.7)

Values are given as percentages (95% CI).

The upward trend in the prevalence of obesity between 2004 and 2007 is similar to that previously seen in other European studies.⁶ The prevalence of overweight appears to be more constant. However, the four-year period analyzed is short as compared to other studies showing upward trends in the prevalence of overweight in other countries.¹⁹

The successive reports of the Spanish National Health Survey issued from 1987 to 2001 show a constant increase in the prevalence of overweight and obesity in all age groups, including their equivalents in the working population.¹³ Prevalence rates of overweight and obesity have increased in Spain during the past 20 years. To date, updated data show no stabilization or decrease in such rates. Moreover, data on the prevalence of overweight and obesity in Spanish children suggest that this increase has not yet reached its peak.²⁰ Because of the close relationship between obesity and type 2 diabetes mellitus, hypertension, dyslipidemia, and some types of cancer, the prevalence of these latter diseases may also increase in the future. In fact, such an increase has already been shown for type 2 diabetes mellitus.^{21,22} This study provides strong evidence regarding the magnitude of the increase in obesity in the active Spanish population, but the reasons for this increase have still to be determined.

It should be noted that the data from this study included direct measurements of weight and height performed by healthcare professionals, so avoiding any bias from self-reported measurements.^{23,24} In addition, the large sample size allowed for more detailed analysis of the results.

The prevalence of obesity was lower in active working women as compared to men. This finding does not agree with prior studies on the general Spanish population,^{8,9} in which a greater prevalence was found in women, particularly in the groups with low socioeconomic level and advanced age. The active population in our study possibly included women who had a higher socioeconomic level and who were younger on

average than the general population, which could account for the lower prevalence.

Obesity prevalence rates remained approximately 50% higher in manual workers as compared to clerical workers throughout the study period, despite the greater physical activity associated with manual occupations. However, differences in type of diet and type of physical activity performed during leisure time may explain these results. In an Australian study on a male population, the prevalence of obesity was lower in farmers than in professional or managerial workers.²⁵ A more specific analysis of the working population in Spain would allow for preventive interventions adapted to specific occupational groups.

On the other hand, the prevalence of obesity increased in both occupational groups throughout the four-year study period. This means that strategies to prevent obesity should be aimed at both groups.²⁶

Of particular note is the high prevalence of obesity seen in women aged 60 years or older with manual occupations as compared to men or the overall group of clerical workers. There was no such difference regarding the prevalence of overweight.

This study was limited to the active population; its data cannot therefore be extrapolated to the general Spanish population without considering occupational status. Moreover, these data are not representative of the older and younger groups of the general population. The prevalence of obesity among the unemployed working population may be even higher than seen in this study. In that case, the aggregate data would be even more alarming.

These findings demonstrate again the need for promoting specific programs to prevent and treat obesity. Medical check-ups of workers are an excellent means of collecting information about the epidemiology and early prevention of obesity and other prevalent conditions.

Funding

This study was funded by Ibermutuamur. The authors acknowledge Irmita Cheung, from the PRIMO Scientific Corporation, for her assistance in the preparation of this manuscript. The writing of the manuscript was funded by Eli Lilly and Company.

Conflicts of interest

Alberto Goday-Arnó, Eva Calvo-Bonacho, Miguel-Ángel Sánchez-Chaparro, José-Antonio Gelpí, Juan-Carlos Sainz, Sonia Santamaría, Rosa-Isabel Navarro, Faustino Gutiérrez, and Carlos Sanz state that they have no conflicts of interest. Elena Caveda and Jesús Reviriego are full-time employees of Eli Lilly and Company.

References

- Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*. 2009;373:1083–96.
- York DA, Rossner S, Caterson I, Chen CM, James WP, Kumanyika S, et al. Prevention Conference VII: Obesity, a worldwide epidemic related to heart disease and stroke: Group I: worldwide demographics of obesity. *Circulation*. 2004;110:e463–70.
- World Health Organization. Diet, nutrition and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation: WHO Technical Report Series. Geneva: World Health Organization; 2003.
- Brownson RC, Boehmer TK, Luke DA. Declining rates of physical activity in the United States: what are the contributors. *Annu Rev Public Health*. 2005;26:421–43.
- Martinez JA, Moreno B, Martinez-Gonzalez MA. Prevalence of obesity in Spain. *Obes Rev*. 2004;5:171–2.
- Seidell JC. Obesity in Europe: scaling an epidemic. *Int J Obes Relat Metab Disord*. 1995;19 Suppl. 3:S1–4.
- Quiles Izquierdo J, Vioque J. Prevalencia de la obesidad en la Comunidad Valenciana. *Med Clin (Barc)*. 1996;106:529–33.
- Aranceta J, Perez-Rodrigo C, Serra Majem L, Ribas L, Quiles Izquierdo J, Vioque J, et al. Influence of sociodemographic factors in the prevalence of obesity in Spain. The SEEDO'97 Study. *Eur J Clin Nutr*. 2001;55:430–5.
- Aranceta J, Perez Rodrigo C, Serra Majem L, Ribas Barba L, Quiles Izquierdo J, Vioque J, et al. Prevalencia de la obesidad en España: resultados del estudio SEEDO 2000. *Med Clin (Barc)*. 2003;120:608–12.
- Aguilera-Zubizarreta E, Ugarte-Miota T, Munoz Cacho P, Varagonzalez L, Sanz de Castro S. Prevalencia de obesidad y sobrepeso en Cantabria. *Gac Sanit*. 2008;22:461–4.
- Serrano-Aguilar P, Munoz-Navarro SR, Ramallo-Farinha Y, Trujillo-Martin MM. Obesity and health related quality of life in the general adult population of the Canary Islands. *Qual Life Res*. 2009;18:171–7.
- Garcia-Alvarez A, Serra-Majem L, Ribas-Barba L, Castell C, Foz M, Uauy R, et al. Obesity and overweight trends in Catalonia, Spain (1992–2003): gender and socio-economic determinants. *Public Health Nutr*. 2007;10:1368–78.
- Gutierrez-Fisac JL, Banegas Banegas JR, Artalejo FR, Regidor E. Increasing prevalence of overweight and obesity among Spanish adults, 1987–1997. *Int J Obes Relat Metab Disord*. 2000;24:1677–82.
- Quiles Izquierdo J, Vioque J. Validez de los datos antropométricos declarados para la determinación de la prevalencia de obesidad. *Med Clin (Barc)*. 1996;106:725–9.
- Sanchez-Chaparro MA, Roman-Garcia J, Calvo-Bonacho E, Gomez-Larios T, Fernandez-Meseguer A, Sainz-Gutierrez JC, et al. Prevalencia de factores de riesgo vascular en la población laboral española. *Rev Esp Cardiol*. 2006;59:421–30.
- Salas-Salvado J, Rubio MA, Barbany M, Moreno B. Consenso SEEDO 2007 para la evaluación del sobrepeso y la obesidad y el establecimiento de criterios de intervención terapéutica. *Med Clin (Barc)*. 2007;128:184–96.
- Instituto Nacional de Estadística. Clasificación nacional de ocupaciones. 1994 [accessed 23.02.10]. Available from: <http://www.ine.es/clasifi/cnoph.htm>
- Instituto Nacional de Estadística. Encuesta de población activa. 2010. Nota de prensa del 30 de abril de 2010. Tablas y anexos [accessed 29.07.10]. Available from: <http://www.ine.es>
- Jolliffe D. Continuous and robust measures of the overweight epidemic: 1971–2000. *Demography*. 2004;41:303–14.
- Salcedo V, Gutierrez-Fisac JL, Guallar-Castillon P, Rodriguez-Artalejo F. Trends in overweight and misperceived overweight in Spain from 1987 to 2007. *Int J Obes (Lond)*. 2010;34: 1759–65.
- Wild SH, Forouhi NG. What is the scale of the future diabetes epidemic, and how certain are we about it. *Diabetologia*. 2007;50:903–5.
- Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health*. 2009;9:88.
- Niedhammer I, Bugel I, Bonenfant S, Goldberg M, Leclerc A. Validity of self-reported weight and height in the French GAZEL cohort. *Int J Obes Relat Metab Disord*. 2000;24:1111–8.
- Nyholm M, Gullberg B, Merlo J, Lundqvist-Persson C, Rastam L, Lindblad U. The validity of obesity based on self-reported weight and height: implications for population studies. *Obesity (Silver Spring)*. 2007;15:197–208.
- Allman-Farinelli MA, Chey T, Merom D, Bauman AE. Occupational risk of overweight and obesity: an analysis of the Australian Health Survey. *J Occup Med Toxicol*. 2010;5:14.
- Neira M, de Onis M. Preventing obesity: a public health priority in Spain. *Lancet*. 2005;365:1386.