

REVIEW ARTICLE

Electroencephalography as a post-stroke assessment method: An updated review[☆]



E. Monge-Pereira^{a,b,*}, F. Molina-Rueda^{a,b}, F.M. Rivas-Montero^{a,b}, J. Ibáñez^c, J.I. Serrano^d, I.M. Alguacil-Diego^{a,b}, J.C. Miangolarra-Page^{a,b}

^a Departamento de Fisioterapia, Terapia Ocupacional, Rehabilitación y Medicina Física, Facultad de CC de la Salud, Universidad Rey Juan Carlos, Alcorcón, Madrid, Spain

^b Laboratorio de Análisis del Movimiento, Biomecánica, Ergonomía y Control Motor, Universidad Rey Juan Carlos, Alcorcón, Madrid, Spain

^c Instituto Cajal, Consejo Superior de Investigaciones Científicas, Madrid, Spain

^d Grupo de Bioingeniería, Consejo Superior de Investigaciones Científicas, Arganda del Rey, Madrid, Spain

Received 11 April 2014; accepted 30 July 2014

Available online 15 February 2017

KEYWORDS

Stroke;
Electroencephalography;
Event-related
desynchronisation;
Event-related
synchronisation;
Movement-related
cortical potentials;
Plasticity

PALABRAS CLAVE

Ictus;
Electroencefalografía;
Desincronización
relacionada con
eventos;

Abstract Given that stroke is currently a serious problem in the population, employing more reliable and objective techniques for determining diagnosis and prognosis is necessary in order to enable effective clinical decision-making. EEG is a simple, low-cost, non-invasive tool that can provide information about the changes occurring in the cerebral cortex during the recovery process after stroke. EEG provides data on the evolution of cortical activation patterns which can be used to establish a prognosis geared toward harnessing each patient's full potential. This strategy can be used to prevent compensation and maladaptive plasticity, redirect treatments, and develop new interventions that will let stroke patients reach their new maximum motor levels.

© 2014 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Electroencefalografía como método de evaluación tras un ictus. Una revisión actualizada

Resumen Actualmente, el ictus representa un problema grave en la población, por lo que es necesario emplear técnicas de evaluación de diagnóstico y pronóstico más fiables y objetivas que permitan una eficaz toma de decisiones clínicas. La electroencefalografía es una herramienta sencilla, de bajo coste y no invasiva, que puede proporcionar conocimiento acerca

[☆] Please cite this article as: Monge-Pereira E, Molina-Rueda F, Rivas-Montero FM, Ibáñez J, Serrano JI, Alguacil-Diego IM, et al. Electroencefalografía como método de evaluación tras un ictus. Una revisión actualizada. Neurología. 2017;32:40–49.

* Corresponding author.

E-mail addresses: esther.monge@urjc.es, emongepereira@gmail.com (E. Monge-Pereira).

Sincronización
relacionada con
eventos;
Potenciales corticales
relacionados con el
movimiento;
Plasticidad

de los cambios que acontecen en el córtex cerebral en el proceso de recuperación tras un ictus. Proporciona datos de evolución de los patrones de activación corticales, que permiten establecer un pronóstico para aprovechar el máximo potencial de las personas evitando compensaciones y plasticidad maladaptativa, redireccionar los tratamientos y desarrollar nuevas intervenciones con el objetivo de alcanzar el nuevo máximo nivel motor de los pacientes que han sufrido un ictus.

© 2014 Sociedad Española de Neurología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Background

Stroke is the second most frequent cause of death in the Spanish general population, and the leading cause of death in women.¹ In 2002, the cumulative incidence rate per 100 000 population in Catalonia among people older than 24 was 218 new cases in men and 127 new cases in women.² It also represents the main cause of disability and generates a significant healthcare and social expenditure.¹

Eighty per cent of the subjects who suffer a stroke survive the acute phase; 6 months later, 85% of them still present motor disorders as well as neurological deficits, communication disorders, cognitive disorders, and disorders in visuospatial perception.^{3,4}

Rehabilitation is a therapeutic process aimed at maximising the patient's physical, psychological, and social potential.⁵ Although many subjects partially recover lower limb motor function, most of them cannot perform activities of daily living with the affected upper limb (UL) after months of rehabilitation.⁶ Functional recovery of the UL is observed in less than 15% of the subjects.⁷ Impairment of the UL significantly limits the patient's level of activity and both physical and social interactions.⁸

The different sequelae in the UL with an impact on activities of daily living have been clinically assessed using the available standardised scales. Considering the results on these scales, the greatest potential for recovery is reached between the second and fifth months after stroke. Prospects for functional recovery after 6 months are considered limited.^{9,10} However, these scales demonstrate a limited capacity for detecting less sensitive changes that are nonetheless necessary for measuring outcomes of stroke patients. In addition, they are subject to rater subjectivity.¹¹ Therefore, we suggest combining imaging techniques with standard scales to obtain more accurate and objective assessments of recovery.¹²

New evaluation tools

Recovery after stroke constitutes one of the main enigmas of the neurosciences. Motor rehabilitation patterns are heterogeneous, and the only prognostic indicators to be used

until recently were the degree of motor impairment^{13,14} and the cardioembolic stroke subtype. The latter represents 14% to 30% of ischaemic strokes, with cerebral infarct being the most severe due to its high recurrence and in-hospital mortality rates during the acute phase.¹⁵ However, a significant effort has been made to research changes to the central nervous system after stroke in order to establish more accurate prognoses and develop new strategies promoting motor rehabilitation.¹⁶

In this line, some neuroimaging techniques have undergone significant developments: functional magnetic resonance imaging (fMRI), transcranial magnetic stimulation (TMS), and positron emission tomography (PET).

Based on different studies which used these assessment techniques, researchers have managed to determine the changes induced by the application of new and groundbreaking rehabilitation approaches, such as robotics,¹⁷ music therapy,¹⁸ and TMS.¹⁹ Furthermore, neuroplastic changes have been proven to play an important role in the recovery after stroke. According to the scientific literature, changes contributing to motor rehabilitation can be found in both hemispheres. On the one hand, various authors have shown that the presence of motor evoked potentials (MEPs) triggered by TMS is a predictor of motor rehabilitation.^{20,21} On the other hand, some authors suggest that the plasticity associated with compensatory movements may contribute to maladaptive plasticity.^{22,23} The studies by Talelli et al.²⁴ and Stinear et al.²⁵ used TMS to analyse ipsilesional corticospinal tract projections. In both studies, TMS data was helpful in predicting motor rehabilitation and adapting rehabilitation strategies. Chelette et al.²⁶ used TMS to analyse the changes caused by rehabilitation of the affected UL in chronic stroke patients. Between 7 and 20 months after stroke, researchers observed an increased area of activation in the contralesional hemisphere, followed by a decrease in the same after 21 months. However, ipsilesional motor maps remained stable between 7 and 13 months, and their areas started to expand between 13 and 31 months after stroke, especially at 20 to 21 months. At the end of the study, there was a 10% improvement on scores on standardised functional tests (Fugl-Meyer Assessment and Wolf Motor Function Test). This underscores the association between neuroplastic changes and motor rehabilitation. These findings suggest that a lack of MEPs during the earliest stages does not indicate a poor prognosis, that both hemispheres contribute to recovery

during different stages, and that recovery after stroke is more dynamic than was once believed.²⁶ Cramer et al.²⁷ observed that decreased activity in fMRI studies of the ipsilesional cortex during movements of the paretic hand was actually associated with greater functional gains 6 weeks after rehabilitation in chronic-phase stroke patients. Ward et al.²⁸ examined the degree of recovery in chronic stroke patients and the recruitment pattern of brain regions during a handgrip task as measured using fMRI. They observed that patients with poorer outcomes activated more areas in the contralesional hemisphere than did controls. Also, patients with better recovery showed relatively normal patterns. Therefore, there seems to be a negative association between functional outcomes and the degree of activity in the ipsilateral hemisphere during a motor task. Jang et al.²⁹ used diffusion tensor tractography (DTT) and TMS to evaluate the corticospinal tract of a patient who showed motor apraxia at 30 months after stroke. After one month of rehabilitation and dopaminergic treatment, the patient's motor recovery increased by 22% (Motricity Index). However, no modifications were observed in the corticospinal tract fibres according to the motor evoked potential parameters measured using DDT and TMS. In this way, they demonstrated that the motor rehabilitation was mainly linked to improvement of motor apraxia and that the corticospinal tract was not impaired.²⁹

The 2 studies published in 2008 and 2010 by Formaggio et al.^{30,31} combined fMRI and electroencephalography (EEG) to study the blood-oxygenation-level dependent signal and changes in brain oscillatory activity in motor areas associated with physiological and pathological events. These studies reported a significant correlation with both finger movements and motor imagery tasks as they observed a decreased event-related desynchronisation (ERD) in the α and β bands in the contralateral hemisphere during performance of the motor or motor imagery task. This decrease is associated with blood-oxygen-level dependent activation in the same areas. This combination provided useful information helping identify the regions contributing to changes in electrical responses more accurately.

In summary, neuroimaging techniques have important implications in terms of diagnosis and clinical decision-making.²⁹ In general, evidence shows that the analysis of cortical activity using these tools expands our knowledge about motor rehabilitation mechanisms and may help us design new treatment strategies and sensitive outcome measures. However, these tools are not always accessible to everyone, since they are expensive and have contraindications and limited availability.³²

EEG as a tool for diagnosing and assessing stroke patients

Contrary to predictions issued by many authors, the introduction of neuroimaging techniques has not relegated EEG to a second plane; it is still widely used in neurosciences since it provides a model for real-time brain functioning.

EEG is commonly used to diagnose vascular epilepsy secondary to stroke in adults; it lets physicians study

the characteristics and clinical outcomes of patients, as well as analyse the effectiveness of different antiepileptic treatments.³³ Furthermore, EEG is typically used as a monitoring method during carotid endarterectomy to detect cerebral ischaemia, which causes an ipsilateral voltage attenuation.³⁴ EEG is also performed to assess coma and pseudocoma states and differentiate them from other states of consciousness,³⁵ as well as to confirm brain death.³⁶

Characterising and studying electrophysiological signals of the central nervous system with EEG equipment is also well established because of different features which make it an optimal clinical evaluation tool. The technique is non-invasive and equipment is portable and inexpensive.³⁷ Several authors have used it to assess cortical functional reorganisation after stroke.^{38–43} EEG studies have shown different patterns of movement-related electric brain activity and different neuronal mechanisms related to motor control.⁴⁴ Furthermore, EEG has shown that activity-dependent brain plasticity takes place throughout an individual's lifetime. As such, analysing the development of plasticity may be a useful method for determining the functional sequelae of stroke and how they evolve.⁴⁵ Considering this potential, an updated review should be conducted to analyse and compile available evidence on the use of EEG to study changes in the cerebral cortex in stroke patients. Additionally, we will attempt to answer the following unresolved questions: could EEG be a useful tool for increasing knowledge of the patterns of cortical changes after stroke? Could it constitute a diagnostic method in clinical practice? Could it be used as an objective tool for determining stroke patients' outcomes? And lastly, can EEG shed light on the effectiveness of traditional rehabilitation interventions?

Development

Characteristics of movement-related cortical activity measured with EEG: slow potentials and cortical oscillatory activity

Brain electrical activity is the consequence of ion currents generated by cellular biochemical processes; this theory was promulgated by Richard Caton in 1875.^{37,46} These currents manifest as oscillations or fluctuations of potentials in neuronal groups.

A motor event generates various responses within the neuronal structures of the brain. On the one hand, it provokes slow movement-related cortical potentials (MRCP) which represent the electroencephalographic evidence of the participation of motor cortex during movement and conscious preparation for the intentional movement.⁴⁷ On the other hand, it causes changes in the amplitude of cortical oscillatory activity. In general terms, there are 2 types of cortical oscillations related to motor behaviour: the α and β rhythms.^{48–51} Motor-related changes in the α and β frequencies result in ERD or event-related synchronisation (ERS).⁵² The ERS represents an increase in the amplitude of α and β frequencies, whereas ERD represents a decrease (Fig. 1).⁴⁷




Figure 1 Characteristic ERD/ERS pattern in a healthy subject. ERD/ERS pattern in a healthy subject before a voluntary and self-initiated reaching movement of the right arm. Movement onset is at $t=0$. The patterns were obtained using the techniques described by Ibáñez et al.⁷⁴

Movement-related frequency desynchronisation and synchronisation

The α frequency is the main background rhythm of normal EEGs in adults.⁴⁶ It is measured with the patient awake and with eyes closed. This activity, whose maximum amplitude is recorded in the occipital regions, occurs during wakefulness and periods of relaxation and shows reactivity to eye closure, which is attenuated with opening of the eyes.^{37,46} By definition, rhythm frequency is 8 to 13 Hz.³⁷ Frequency amplitude is usually symmetrical, but it may often be higher in the non-dominant hemisphere. If the patient is not relaxed, α waves may not occur, and their absence in one hemisphere is a pathological sign.⁴⁶

According to the cortical region in question, α frequency presents specific characteristics and has a set nomenclature. Mu (μ) rhythm is located around the central sulcus (primary sensorimotor cortices).³⁷ Mu rhythms present a particular reactivity, since they occur at rest and resolve with contralateral limb movements. They are particularly pronounced around the area of the motor cortex associated with the hand and resolve when making a fist, or even during motor imagery tasks.⁵⁰ They are regarded as intrinsic activity of the central region and are thus very useful in the functional evaluation of stroke patients.⁵³

Low amplitude β activity presents frequencies of 14 to 30 Hz and indicates mental activity and attention.³⁷ Maximum β amplitude is observed in the fronto-central regions. The most significant finding in the analysis of β activity is interhemispheric asymmetry. A decrease in amplitude in one of the hemispheres tends to indicate presence of alterations.⁴⁶

Amplitude of oscillatory activity decreases when frequency increases. Therefore, μ rhythm with a frequency of 8 to 13 Hz has a higher amplitude than β activity. Furthermore, oscillation amplitude is proportional to the number of synchronised active neurons, so that lower oscillations involve more neurons than faster oscillations.⁵²

In general, sensory stimulation and the actual movement or motor imagery task will result in ERD or ERS of μ and β rhythms. Movement is preceded by ERD in healthy subjects, and as the movement is carried out, variations occur in the

synchronisation of cortical rhythms (ERS). Movement offset is followed by a β -ERS generated in the precentral region, which reflects a deactivation of motor cortex. This occurs in the first second after termination of voluntary movement, when the μ rhythm still presents a low-amplitude ERD. Bilateral movements cause bilateral activation.⁵⁴

μ -ERD does not depend on movement duration or localisation, and it is the same for movements of the index, the thumb, or even the hand. This suggests that μ -ERD is due to a non-specific preactivation of neurons in the motor areas.⁵² In this line, studies with electrocorticography have shown that β -ERD is more discrete than μ -ERD. Likewise, it is located in the prerolandic cortex, a more anterior area (motor area), whereas μ -ERD is located in the postrolandic area (primary somatosensory cortex).⁵⁵

ERD and ERS can take place simultaneously, both in α activity and β activity. In the first case, during perception of a visual stimulus, occipital α -ERD and μ -ERS of central rhythms can be observed in EEG recordings. However, occipital α -ERS and central μ -ERD occur during self-initiated voluntary movements.⁵⁶ A possible explanation would be the involvement of thalamic structures which activate cortical structures engaged in some movements while deactivating others not involved in the task. At the β activity level, ERD and ERS take place simultaneously during a hand motor imagery task, which activates a contralateral ERD and ipsilateral ERS.⁵²

For ERD/ERS calculations using EEG at movement onset, variations of power in μ and β bands are expressed with reference to a value calculated in a reference time period, always taken at rest, long before the movement is to be performed, in order to remain as neutral as possible (Tables 1 and 2).⁵⁷

Slow MRCP

MRCP are especially relevant in the assessment of cortical activation patterns, since they are associated with movement planning and execution. This being the case, Bereitschaftspotential (BP) will require a special mention as they are considered the initial stage of MRCPs.^{41,47,58} This parameter, introduced by Kornhuber and Deecke in 1964, is an electrical signal corresponding to the supplementary motor area, which activates before the primary motor area.⁵⁹ This is a slow and negative potential which reflects the preparatory activity and whose maximum peak coincides with voluntary movement.⁶⁰ BP has 2 components: an early component initiating 2 seconds before the movement starts, and a late component, which appears 400 ms before the movement itself. Early BP is maximal at the midline of the central parietal area, and widely and symmetrically distributed across both hemispheres. Late BP is much larger over the right central region (contralateral to the movement) (Fig. 2). In general terms, when a movement pattern is present, the BP commonly starts much earlier, compared to movements executed in more natural conditions.⁴⁷

Although the mechanisms generating BP are yet to be understood, these potentials are modulated by the way the task is performed. Multiple factors influence the type of BP: the preparatory state, movement selection

Table 1 Characteristics of cortical motor rhythms.

Cortical oscillatory activity	Frequency	Event	Location	Consequence of the event
Alpha (α) rhythms	8–13 Hz	Levels of consciousness. Reactivity to eye closure.	Occipital area	
Mu (μ) rhythms	8–13 Hz	At rest. They resolve with contralateral limb movements or motor imagery	Somatosensory cortex (central region)	ERD (before movement regardless of movement duration, similar in slow and vigorous movements, independent from location)
Beta (β) rhythms	14–30 Hz	Mental activity and attention	Frontal-central region	ERS (during the event) ERD (before movement) ERS (during the event and at the end of movement, it reflects deactivation of motor cortex)

ERD, event-related desynchronisation; ERS, event-related synchronisation.

Table 2 Characteristics of motor cortex oscillations in healthy subjects during movement planning and execution.

Healthy subjects	ERD	ERS	MRCPs
Preparation	μ -ERD and β -ERD in the hemisphere contralateral to movement		Associated with movement planning and execution
Execution		β -ERS	The highest amplitude coincides with movement onset and then starts to decrease

μ -ERD, β -ERD, event-related desynchronisation in the μ and β frequency bands; β -ERS, event related synchronisation in the β frequency band.




Figure 2 Characteristic BP (Bereitschaftspotential) pattern in a healthy subject. The figure includes BP patterns obtained by 28 EEG channels in a healthy subject who performed a voluntary and self-initiated reaching movement of the right arm. Movement onset is at $t = 0$. The patterns were obtained using the techniques described by Ibáñez et al.⁷⁴

(free or repeated pattern), learning acquisition, movement repetition rhythm, practice, difficulty, speed, and pathological lesion in brain structures, among others.^{41,50} Neurological changes modify the time of appearance and amplitude of MRCPs; as such, they can be useful in diagnosis and follow-up.⁶¹

Characteristics of cortical activity in stroke patients measured with EEG

Studies with EEG by Pfurtscheller^{62–65} in the 1980s revealed that the dynamic amplitude of neural oscillations represents an indicator of neuronal activation during movement preparation and execution. Pfurtscheller et al.⁶⁴ found an attenuation in ERD in the affected hemisphere of stroke patients compared to the unaffected hemisphere during movements of the affected hand.

In the last 10 years, numerous studies have assessed cortical activity in stroke patients with EEG to analyse cortical reorganisation throughout the course of their recovery.

Platz et al.⁴⁴ used EEG to study MRCPs and ERD in α and β frequencies in subjects with acute and subacute hemiparesis (3–20 weeks after stroke). They analysed the preparation and execution stages during self-initiated and repeated movements of the affected hand. The authors observed that in healthy controls, slow potentials extended from the central electrode throughout the cortex during movement preparation, to later become more obvious in the contralateral hemisphere during movement execution. They also verified that ERD started in the contralateral hemisphere and extended to the ipsilateral hemisphere, with bilateral distribution during execution. However, patients showed significant differences with regard to control subjects, especially during movement preparation. The authors observed a more contralateral and frontal distribution of MRCPs, which may suggest more extensive involvement of motor and premotor cortices in motor planning in stroke patients. In terms of cortical oscillatory activity, β -ERD was more localised in the occipital area in patients than in controls during movement preparation. This finding may suggest a higher involvement of visual areas (movement ideation). Furthermore, an increase in the α -ERD in the frontal and ipsilateral area of the cortex was observed during execution. This is associated with an increased need for concentration on the task and compensatory measures in the unaffected hemisphere.⁴⁴

Serrien et al.⁶⁶ described the ipsilateral hemisphere as playing a functional role during movement of the affected hand in chronic stroke patients. For the execution of the same task, somatosensory and motor cortical recruitment originates in the contralateral hemisphere in control subjects and well-recovered patients, and in the ipsilateral hemisphere in patients with a poorer recovery. They established that the more extensive the involvement of contralateral areas, the better the prognosis for functional recovery.

Wiese et al.⁶⁷ compared MRCPs from chronic stroke patients and healthy control subjects as they moved the index finger on the affected hand. In the patient group, the component directly preceding movement onset was significantly reduced in the lesioned hemisphere. The researchers

observed a significant increase in MRCP activity in the contralesional hemisphere during the movement itself. The presence of this early MRCP component, which is generated by the primary motor cortex, suggests that stroke is accompanied by damage to these structures in the affected hemisphere.

Daly et al.⁶⁸ used EEG to study MRCPs in 10 patients with chronic stroke in order to determine motor planning time of a reaching task. They observed a high MRCP amplitude in the sensorimotor cortex as well as in the frontal areas, which reflects a lower degree of automatization of the task and the need for compensatory strategies to perform the movement. Gerloff et al.⁶⁹ showed that α -ERD and β -ERD were reduced in the injured hemisphere of stroke patients in the chronic stage, and increased in the unaffected hemisphere during movements of the affected hand. This suggests a shift in compensatory functional connectivity towards the contralesional hemisphere.

Fattapposta et al.⁷⁰ followed up on 9 acute stroke patients during 1 year by taking a total of 4 measurements at 0, 3, 9, and 12 months. They conducted a study which assessed the pre-programming of movements taking place during execution of a simple task. Patients had a significantly lower BP amplitude than control subjects in the first recording, but amplitudes increased in the subsequent 3 recordings. This result may be associated with early neuronal plasticity and other processes involved in neurorepair, such as angiogenesis and neurogenesis, which occur in the early stages after the lesion.⁷¹ However, regaining automatization of simple tasks would take longer. This provides an explanation for a habituation phenomenon in which the motor task becomes memorised. Mobility scales (Nine Hole Peg Test and Motricity Index Test) showed significant improvements in motor skills throughout the process. The authors stated that patients recovered a relatively good motor skill level, while lacking an adequate recovery of attention and control abilities in the skilled motor act, thereby requiring development of motor compensation strategies. These associative functions are crucial for automating skilled movements, considering that most of daily motor practice is automated.

Stępień et al.⁷² used EEG to assess ERD in stroke patients in the acute stage (less than 8 days of progression). They observed that movements of the affected hand caused a stronger α -ERD in the sensorimotor area (peri-central area) of the ipsilateral hemisphere than in the ipsilesional hemisphere. Furthermore, they reported that α -ERD amplitude was preserved in the affected hemisphere during movements of the unaffected hand; in contrast, activity of the affected hand caused a decrease in the frequency of α oscillations in the affected hemisphere. The ERD is sensitive to the recovery of networks after stroke. A relatively well-preserved ERS in the affected hemisphere during movements of the unaffected hand seems to indicate good potential for recovery, and therefore a positive prognosis. This also suggests that bilateral rehabilitation during at least the first stages of recovery is also important.⁷²

Dean et al.⁷³ studied motor planning in chronic stroke patients and observed reduced MRCP amplitude in the affected hemisphere, especially during preparation of two-handed movements. This shows altered anticipatory

attention activity and motor processing in those patients with poorer functional recovery.

Ibáñez et al.⁷⁴ used EEG to analyse the temporal precision of intention to perform a movement in 3 chronic stroke patients. They used ERD as a parameter to measure oscillatory activity. The authors observed that one subject presented bilateral ERD, while another presented a significant predominance of the central area; both situations were indicative of cortical reorganisation. The third subject presented activation of the regions contralateral to the affected limb that became bilateral once the patient began executing movement. This patient tested at the highest functional level on the UL (Table 3).

There is little evidence on using EEGs for diagnostic purposes. Pfurtscheller et al.⁶² conducted a study with a large sample of patients ($n=50$) to assess cortical oscillatory activity during movements of the affected hand and predict the location of the lesion in the territory of the left middle cerebral artery. Results suggested that an increased rhythm in the ipsilateral hemisphere, together with a symmetric ERD, indicates a deep lesion with a probability of 95% (and a superficial lesion with a probability of 5%). Hemispheric μ amplitude symmetry and asymmetric ERD indicate superficial cortical ischaemia with a probability of 81%; an ipsilaterally attenuated μ rhythm accompanied by an asymmetric or abolished ERD

Table 3 Summary of scientific evidence using EEG to assess motor cortex in stroke patients.

Reference	Stroke phase	MRCP findings	Oscillation findings	Event
Pfurtscheller et al. ^{64,65}	Subacute		ERD attenuated in the affected hemisphere	Movements of the affected hand
Platz et al. ⁴⁴	Acute and subacute	Contralateral and frontal MRCPs during movement preparation (motor and premotor cortices involvement)	Occipital β -ERD during movement preparation (less involvement of visual areas, movement ideation) Frontal and ipsilateral α -ERD during movement execution	Movements of the affected hand
Serrien et al. ⁶⁶	Chronic		Information flows from the ipsilateral hemisphere to the contralateral hemisphere (affected)	Movements of the affected hand
Wiese et al. ⁶⁷	Chronic	Amplitude is significantly reduced in the affected hemisphere during movement preparation. Significant increase in the unaffected hemisphere during execution		Movements of the index finger of the affected hand
Daly et al. ⁶⁸	Chronic	Increased amplitude in the sensorimotor cortices and frontal areas		Reaching task
Gerloff et al. ⁶⁹	Chronic		During execution, α -ERD and β -ERD are decreased in the affected hemisphere and increased in the unaffected hemisphere	Movements of the affected hand
Fattapposta et al. ⁷⁰	Subacute (follow-up of one year)	Higher amplitude at the end of follow-up		Movements of the affected hand
Stępień et al. ⁷²	Acute		μ -ERD of higher amplitude in the ipsilateral hemisphere than in the affected hemisphere during execution.	Movements of the affected hand
Dean et al. ⁷³	Chronic	Amplitude is reduced in the affected hemisphere during movement preparation.		
Ibáñez et al. ⁷⁴	Chronic		Patient with better functionality, ERD in the affected hemisphere during preparation which becomes bilateral during execution	Reaching task

ERD, event-related desynchronisation; ERS, event-related synchronisation; MRCPs, movement-related cortical potentials; α , β , and μ , frequency bands which show the oscillations.

indicates an infarct extending broadly over the entire arterial territory.

Conclusions

EEG-based analysis of movement preparation and execution by comparing stroke patients and healthy controls is useful for determining diagnoses and predicting outcomes. Near-normal cortical activation is associated with better functionality, and progression toward normal characteristics of cortical activation suggests that therapy has been effective. Beyond a certain progression time, patients are excluded from rehabilitation programmes after having been assessed using standardised scales. We need tools that are sensitive to functional changes and thus useful for assessing patient outcomes, effectiveness of interventions, and the real prognosis to facilitate decision-making in clinical practice.

EEG is an inexpensive, non-invasive, simple, portable, and potentially useful tool for diagnosis and prognosis. It provides insight into neuroplastic changes in terms of motor recovery, thereby making it possible to assess the effectiveness of interventions and in turn facilitating clinical decision-making. It also lets us maximise the recovery potential of patients with motor disability by preventing maladaptive plasticity, and developing new interventions intended to modulate cerebral function. An early prediction of the type of motor rehabilitation in question may help to determine the time, duration, and objectives of therapeutic protocols that will be tailored for each patient with neuroplasticity in mind.

A promising new line of research would entail studying motor cortex behaviour using surface EEG recordings, taking into account lesion type and time of progression, in a large sample of patients.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Ministerio de Sanidad, Política Social e Igualdad. Sistema Nacional de Salud. Estrategia del ICTUS del SNS 2008.
- Marrugat J, Arboix A, García-Eroles L, Salas T, Vila J, Castell C, et al. Estimación de la incidencia poblacional y la mortalidad de la enfermedad cerebrovascular establecida isquémica y hemorrágica en 2002. *Rev Esp Cardiol.* 2007;60:573–80.
- Belda-Lois JM, Mena-del Horno S, Bermejo-Bosch I, Moreno JC, Pons JL, Farina D, et al. Rehabilitation of gait after stroke: a review towards a top-down approach. *J Neuroeng Rehabil.* 2011;8:66.
- Dobkin BH. Strategies for stroke rehabilitation. *Lancet Neurol.* 2004;3:528–36.
- Jiménez Muro M, Pedro-Cuesta JJ, Almazán J, Holmqvist W. Stroke recovery in South Madrid. Function and motor recovery, resource utilization, and family support. *Stroke.* 2000;31:1352–9.
- Adams HP, Brott TG, Crowell RM, Burlan AJ, Gomez CR, Grotta J, et al. Guidelines for the management of patient with acute ischemic stroke: a statement for healthcare profession from a special writing group of the stroke council. American Heart Association. *Stroke.* 1994;25:1901–14.
- Hendricks HT, Zwarts MJ, Plat EF, van Limbeek J. Systematic review for the early prediction of motor and functional outcome after stroke by using motor-evoked potentials. *Arch Phys Med Rehabil.* 2002;83:1303–8.
- Kwakkel G, Kollen B. Predicting improvement in the upper paretic limb after stroke: a longitudinal prospective study. *Restor Neurol Neurosci.* 2007;25:453–60.
- Jorgensen HS, Nakayama H, Raaschou HO, Vive-Larsen J, Stoier TS, Olsen TS. Outcome and time course of recovery in stroke. Part I: Outcome. The Copenhagen stroke study. *Arch Phys Med Rehabil.* 1995;76:399–405.
- Jorgensen HS, Nakayama H, Raaschou HO, Vive-Larsen J, Stoier TS, Olsen TS. Outcome and time course of recovery in stroke. Part II: Time course of recovery. The Copenhagen stroke study. *Arch Phys Med Rehabil.* 1995;76:406–12.
- Peña-Pitarch E, Tico-Falguera N. Assessment of damage in affected persons with neurological sequelae. *Trauma.* 2012;23:247–52.
- Stinear C. Prediction of recovery of motor function after stroke. *Lancet Neurol.* 2010;9:1228–32.
- Takeuchi N, Izumi S. Rehabilitation with poststroke motor recovery: a review with a focus on neural plasticity. *Stroke Res Treat.* 2013;2013:128641.
- Montaner J, Álvarez-Sabin J. Escala de ictus del National Institute of Health (NIHSS) y su adaptación al español. *Neurología.* 2006;21:192–202.
- Arboix A, Alio J. Cardioembolic stroke: clinical features, specific cardiac disorders and prognosis. *Curr Cardiol Rev.* 2010;6:150–61.
- Hoyer EH, Celnik PA. Understanding and enhancing motor recovery after stroke using transcranial magnetic stimulation. *Restor Neurol Neurosci.* 2011;29:395–409.
- Turolla A, Daud Albasini OA, Oboe R, Agostini M, Tonin P, Paolucci S, et al. Haptic-based neurorehabilitation in post-stroke patients: a feasibility prospective multicentre trial for robotics hand rehabilitation. *Comput Math Methods Med.* 2013;2013:895492, <http://dx.doi.org/10.1155/2013/895492>.
- Tomaino CM. Creativity and improvisation as therapeutic tools within music therapy. *Ann N Y Acad Sci.* 2013;1303:84–6.
- Volz LJ, Sarfeld AS, Diekhoff S, Rehme AK, Pool EM, Eickhoff SB, et al. Motor cortex excitability and connectivity in chronic stroke: a multimodal model of functional reorganization. *Brain Struct Funct.* 2014, <http://dx.doi.org/10.1007/s00429-013-0702-8>.
- Escudero JV, Sancho J, Bautista D, Escudero M, López-Trigo J. Prognostic value of motor evoked potential obtained by transcranial magnetic brain stimulation in motor function recovery in patients with acute ischemic stroke. *Stroke.* 1998;29:1854–9.
- VanKuijk AA, Pasman JW, Hendricks HT, Zwarts MJ, Geurts AC. Predicting hand motor recovery in severe stroke: the role of motor evoked potentials in relation to early clinical assessment. *Neurorehabil Neural Repair.* 2009;23:45–51.
- Takeuchi N, Izumi S. Maladaptive plasticity for motor recovery after stroke: mechanisms and approaches. *Neural Plast.* 2012;2012:359–728.
- Levin MF, Kleim JA, Wolf SL. What do motor recovery and compensation mean in patients following stroke? *Neurorehabil Neural Repair.* 2009;23:313.
- Talelli P, Greenwood RJ, Rothwell JC. Arm function after stroke: neurophysiological correlates and recovery mechanisms assessed by transcranial magnetic stimulation. *Clin Neurophysiol.* 2006;117:1641–59.

25. Stinear CM, Barber PA, Smale PR, Coxon JP, Fleming MK, Byblow WD. Functional potential in chronic stroke patients depends on corticospinal tract integrity. *Brain*. 2007;130:170–80.
26. Chelette KC, Carrico C, Laurie Nichols L, Sawaki L. Long-term cortical reorganization following stroke in a single subject with severe motor impairment. *NeuroRehabilitation*. 2013;33:385–9.
27. Cramer SC, Parrish TB, Lev RM. Predicting functional gains in a stroke trial. *Stroke*. 2007;38:2108–14.
28. Ward NS, Brown MM, Thompson AJ, Frackowiak RSJ. Neural correlates of outcome after stroke: a cross-sectional fMRI study. *Brain*. 2003;126:1430–48.
29. Jang SH. Motor recovery by improvement of limb-kinetic apraxia in a chronic stroke patient. *NeuroRehabil*. 2013;33:195–200.
30. Formaggio E, Storti SF, Avesani M, Cerini R, Milanese F, Gasparini A, et al. EEG and fMRI coregistration to investigate the cortical oscillatory activities during finger movement. *Brain Topogr*. 2008;21:100–11.
31. Formaggio E, Storti SF, Cerini R, Fiaschi A, Manganotti P. Brain oscillatory activity during motor imagery in EEG-fMRI coregistration. *Magn Reson Imaging*. 2010;28:1403–12.
32. Jiang L, Xu H, Yu C. Brain connectivity plasticity in the motor network after ischemic stroke. *Neural Plast*. 2013;2013:924192.
33. Temprano T, Salas-Puig J, Calleja-Puerta S, Zanabali Al-Sibbai AA, Lahoz CH. Epilepsia postictus. *Rev Neurol*. 2009;48:171–7.
34. Malcharek MJ. Outcomes of combined somatosensory evoked potential. Motor evoked potential, and electroencephalography monitoring during carotid endarterectomy. *Ann Vasc Surg*. 2014;28:1351.
35. Gawryluk JR, D'Arcy RC, Connolly JF, Weaver DF. Improving the clinical assessment of consciousness with advances in electrophysiological and neuroimaging techniques. *BMC Neurol*. 2010;29:10–1.
36. Su Y, Yang Q, Liu G, Zhang Y, Ye H, Gao D, et al. Diagnosis of brain death: confirmatory tests after clinical test. *Chin Med J (Engl)*. 2014;127:1272–7.
37. Gil-Nagel A, Parra J, Iriarte J, Kanner AM. Manual de electroencefalografía. 2.^a ed. Madrid: Mc Graw-Hill Interamericana; 2004.
38. Baird AE, Benfield A, Schlaug G, Siewert B, Lovblad KO, Edelman RR, et al. Enlargement of human cerebral ischemic lesion volumes measured by diffusion-weighted magnetic resonance imaging. *Ann Neurol*. 1997;41:581–91.
39. Chollet F, DiPiero V, Wise RJ, Brooks DJ, Dolan RJ, Frackowiak RS. The functional anatomy of motor recovery after stroke in humans: a study with positron emission tomography. *Ann Neurol*. 1991;29:63–71.
40. Weiller C, Chollet F, Friston KJ, Wise RJ, Frackowiak RS. Functional reorganization of the brain in recovery from striatocapsular infarction in man. *Ann Neurol*. 1992;31:463–72.
41. Cramer SC, Nelles G, Benson RR, Kaplan JD, Parker RA, Kwong K, et al. A functional MRI study of subjects recovered from hemiparetic stroke. *Stroke*. 1997;28:2518–27.
42. Cao Y, D'Olhaberriague L, Vikingstad EM, Levine SR, Welch KM. Pilot study of functional MRI to assess cerebral activation of motor function after poststroke hemiparesis. *Stroke*. 1998;29:112–22.
43. Seitz RJ, Hoflich P, Binkofski F, Tellmann L, Herzog H, Freund HJ. Role of the premotor cortex in recovery from middle cerebral artery infarction. *Arch Neurol*. 1998;55:1081–8.
44. Platz T, Kim IH, Pintschovius H, Winter T, Kieselbach A, Villringer K, et al. Multimodal EEG analysis changes in movement-related electric brain activity after stroke. *Brain*. 2000;123:2475–90.
45. Daly JJ, Wolpaw JR. Brain—computer interfaces in neurological rehabilitation. *Lancet Neurol*. 2008;7:1032–43.
46. Rowan AJ, Tolunsky E. Conceptos básicos sobre EEG. 4.^a ed. Madrid: Elsevier; 2004.
47. Shibasaki H, Hallett M. What is the Bereitschaftspotential? *Clin Neurophysiol*. 2006;117:2341–56.
48. Brown P. Cortical drives to human muscle: the Piper and related rhythms. *Prog Neurobiol*. 2000;60:97–108.
49. Gerloff C, Hadley J, Richard J, Uenisch N, Honda M, Hallett M. Functional coupling and regional activation of human cortical motor areas during simple, self-paced and metronome-paced finger movements. *Brain*. 1998;121:1513–31.
50. Serrien DJ, Brown P. The functional role of interhemispheric synchronization in the control of bimanual timing tasks. *Exp Brain Res*. 2002;147:268–72.
51. Serrien DJ, Brown P. The integration of cortical and behavioral dynamics during initial learning of a motor task. *Eur J Neurosci*. 2003;17:1098–104.
52. Pfurtscheller G, Lopes da Silva FH. Event-related EEG/MEG synchronization and desynchronization: basic principles. *Clin Neurophysiol*. 1999;110:1842–57.
53. Pfurtscheller G. Rolandic mu rhythms and assessment of cerebral functions. *Am J EEG Technol*. 1986;26:19–32.
54. Formaggio E, Storti SF, Boscolo Galazzo I, Gandolfi M, Geroin C, Smania N, et al. Modulation of event-related desynchronization in robot-assisted hand performance: brain oscillatory changes in active, passive and imagined movements. *J Neuroeng Rehabil*. 2013;26:24.
55. Crone NE, Miglioretti DL, Gordon B, Lesser RP. Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis. II. Event-related synchronization in the gamma band. *Brain*. 1998;121:2301–15.
56. Pfurtscheller G, Klimesch W. Functional topography during a visuoverbal judgement task studied with event-related desynchronization mapping. *J Clin Neurophysiol*. 1992;9:120–31.
57. Cassim F, Szurhaj W, Sediri H, Devos D, Bourriez J, Poirot I, et al. Brief and sustained movements: differences in event-related (de)synchronization (ERD/ERS) patterns. *Clin Neurophysiol*. 2000;111:2032–9.
58. Farina D, do Nascimento OF, Lucas MF, Doncarli C. Optimization of wavelets for classification of movement-related cortical potentials generated by variation of force-related parameters. *J Neurosci Methods*. 2007;15(1–2):357–63.
59. Deecke L, Kornhuber HH. An electrical sign of participation of the mesial “supplementary” motor cortex in human voluntary finger movement. *Brain Res*. 1978;159:473–6.
60. Trevena J, Miller J. Brain preparation before a voluntary action: evidence against unconscious movement initiation. *Conscious Cognit*. 2010;19:447–56.
61. Mrachacz-Kresting N, Kristensen SR, Niazi IK, Farina D. Precise temporal association between cortical potentials evoked by motor imagination and afference induces cortical plasticity. *J Physiol*. 2012;590:1669–82.
62. Pfurtscheller G, Sager W, Wege W. Correlations between CT scan and sensorimotor EEG rhythms in patients with cerebrovascular disorders. *Electroencephalogr Clin Neurophysiol*. 1981;52:473–85.
63. Pfurtscheller G, Aranibar A. Evaluation of event-related desynchronization (ERD) preceding and following voluntary self-paced movements. *Electroencephalogr Clin Neurophysiol*. 1979;46:138–46.
64. Pfurtscheller G, Aranibar A, Wege W. Changes in central EEG activity in relation to voluntary movement. II Hemiplegic patients. *Prog Brain Res*. 1980;54:491–5.
65. Pfurtscheller G, Wege W, Sager W. Asymmetrien in der zentralen alpha-Aktivität (mu-Rhythmus) unter ruhe-und aktivitätsbedingungen bei zerebrovaskulären erkrankungen. *EEG EMG. Elektroenzephalogr Elektromyogr Verwandte Geb*. 1980;11:63–71.
66. Serrien DJ, Strensa LH, Cassidy MJ, Thompson AJ, Brown P. Functional significance of the ipsilateral hemisphere during movement of the affected hand after stroke. *Exp Neurol*. 2004;190:425–32.

67. Wiese H, Stude P, Sarge R, Nebel K, Diener HC, Keidel M. Reorganization of motor execution rather than preparation in poststroke hemiparesis. *Stroke*. 2005;36:1474–9.
68. Daly JJ, Fang Y, Perepezko EM, Siemionow V, Yue GH. Prolonged cognitive planning time, elevated cognitive effort, and relationship to coordination and motor control following stroke. *IEEE Trans Neural Syst Rehabil Eng*. 2006;14:168–71.
69. Gerloff C, Bushara K, Sailer A, Wassermann EM, Chen R, Matsumoto T, et al. Multimodal imaging of brain reorganization in motor areas of the contralesional hemisphere of well recovered patients after capsular stroke. *Brain*. 2006;129:791–808.
70. Fattapposta F, D'Agostino VC, My F, Locuratolo N, Vanacore N, Inghilleri M, et al. Psychophysiological aspects of voluntary skilled movement after stroke: a follow-up study. *Arch Ital Biol*. 2008;146(3–4):147–63.
71. Font MA, Arboix A, Krupinski J. Angiogenesis, neurogenesis and neuroplasticity in ischemic stroke. *Curr Cardiol Rev*. 2010;6:238–44.
72. Stępień M, Conradi J, Waterstraat G, Hohlefeld FU, Curio G, Nikulin VV. Event-related desynchronization of sensorimotor EEG rhythms in hemiparetic patients with acute stroke. *Neurosci Lett*. 2011;488:17–21.
73. Dean PJA, Seiss E, Sterr A. Motor planning in chronic upper-limb hemiparesis: evidence from movement-related potentials. *PLoS ONE*. 2012;7.
74. Ibáñez J, del Castillo MD, Serrano JI, Molina Rueda F, Monge Pereira E, Rivas Montero FM, et al. Single-trial detection of the event-related desynchronization to locate with temporal precision the onset of voluntary movements in stroke patients. *IFMBE Proc*. 2014;41:1651–4.