

of cardiovascular risk factors.⁸ Results from the screening tests for opportunistic infections and tumours, and the improvement in visual acuity after antiretroviral treatment support this hypothesis.⁹ HIV infection has also been associated with relapsing-remitting neuritis with inflammation (MS-like neuritis), which responds well to corticosteroid treatment. Some researchers suggest administering penicillin even when serology tests for syphilis are negative due to the high frequency of co-occurrence of these 2 infections.¹

Closely monitoring HIV-positive patients with posterior ischaemic optic neuropathy is essential due to the likelihood of subsequent acute retinal necrosis secondary to herpes zoster or CNS lymphoma. The likelihood of experiencing this complication increases in patients diagnosed with acquired immune deficiency syndrome. Optic neuropathy secondary to HIV infection does not seem to be correlated with either the CD4 lymphocyte level or the presence of opportunistic infections.¹

Conclusion

Unilateral retrobulbar optic neuropathy is an infrequent manifestation of HIV infection. However, it should be included in the differential diagnosis of seropositive patients with atypical symptoms, since it may also constitute the initial manifestation of HIV infection. Diagnosis is made by exclusion, after ruling out presence of opportunistic infections, CNS neoplasm, or drug allergies.

Funding

This study has received no public or private funding.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

1. Stewart MW. Human immunodeficiency virus and its effects on the visual system. *Infect Dis Rep.* 2012;4:e25.
2. Jiménez Caballero PE, Serviá Candela M. Bilateral optic neuropathy in an HIV patient. *Neurología.* 2010;25:267–8.
3. Gautier D, Rabier V, Jallet G, Milea D. Visual loss related to macular subretinal fluid and cystoid macular edema in HIV-related optic neuropathy. *Int Ophthalmol.* 2012;32:405–8.
4. Winward KE, Harmed LM, Glaser JS. The spectrum of optic nerve disease in human immunodeficiency virus infection. *Am J Ophthalmol.* 1989;107:373–80.
5. Babu K, Murthy KR, Rajagopalan N, Satish B. Vision recovery in human immunodeficiency virus-infected patients with optic neuropathy treated with highly active antiretroviral therapy: a case series. *Indian J Ophthalmol.* 2009;57:315–8.
6. Mustak H, Rogers G, Cook C. Ethambutol induced toxic optic neuropathy in HIV positive patients. *Int J Ophthalmol.* 2013;6:542–5.
7. Rebolledo Fernández G, Muñoz Negrete F. In: *Protocolos en Neuro-oftalmología.* Sociedad Española de Oftalmología; Madrid: Macs Line SL; 2010. Cap. 3. p. 81.
8. Newman NJ, Lessell S. Bilateral optic neuropathies with remission in acute human immunodeficiency virus infection. *Acta Ophthalmol Scand.* 1992;76:737–8.
9. Goldsmith P, Jones RE, Ozuzu GE, Richardson J, Ong EL. Optic neuropathy as the presenting featuring of VIH infection: recovery of vision with highly active antirretroviral therapy. *Br J Ophthalmol.* 2000;84:551–3.

F. Pérez-Bartolomé*, E. Santos-Bueso, N. Ávalos-Franco, J. Porta-Etessam

Unidad de Neurooftalmología, Servicio de Oftalmología, Instituto de Investigación Sanitaria, Hospital Clínico San Carlos (IdISSC), Madrid, Spain

* Corresponding author.

E-mail address: Franciscoperezbartolome@gmail.com

(F. Pérez-Bartolomé).

2173-5808/

© 2015 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The effect of manual therapy on tension headache[☆]



Efecto de la terapia manual en la cefalea tensional

Dear Editor:

It was with great interest that we read the article published by Lozano López et al.,¹ which presents a systematic

review of the literature supporting manual therapy as an effective alternative pain relief technique in patients with tension-type headache (TTH). The most frequently used manual therapy techniques in the analysed studies were joint treatment, soft tissue mobilisation techniques, and muscle exercises.² Although these techniques have been shown to have a positive effect on TTH, the heterogeneity of patients' symptoms has not permitted the adoption of a standard protocol with a specific technique. In fact, many of these studies report better results when combining 2 or more manual techniques.³ We agree that the diversity of interventions does not allow us to determine which technique was the most effective for TTH. In our view, however, results are significant in that applying these techniques decreased not only pain intensity but also headache frequency.⁴

Although the true cause of TTH is still to be determined, the frequent co-presence of such symptoms as trigger points

[☆] Please cite this article as: Caballero-Calixto L, Antón-Puescas P, Reyna-Ormeño J. Efecto de la terapia manual en la cefalea tensional. *Neurología.* 2017;32:201–202.

in the head and neck,⁵ cervical rigidity, and inhibition of local stabilising muscles underscores the need for a manual therapy programme addressing each of the many problems associated with TTH.

Manual therapy is a widespread treatment approach in some countries. However, in Peru and other countries in which manual therapy is a new specialty, treatment protocols for TTH are usually limited to pharmacological treatment and do not include physiotherapeutic management of the symptoms associated with TTH. Increasing numbers of manual therapists and further studies of manual therapy will be necessary to generate evidence and draw conclusions about the usefulness of this technique for managing TTH and associated symptoms.

Conflicts of interest

The authors of this letter have no conflicts of interest to declare.

Acknowledgements

The authors would like to thank their families for their support during medical training.

References

1. Lozano López C, Mesa Jiménez J, de la Hoz Aizpurúa JL, Pareja Grande J, Fernández de las Peñas C. Eficacia de la terapia manual en el tratamiento de la cefalea tensional. Una revisión sistemática desde el año 2000 hasta el 2013. *Neurología*. 2014;1:242.
2. Castien RF, van der Windt DA, Dekker J, Mutsaers B, Grooten A. Effectiveness of manual therapy compared to usual care by the general practitioner for chronic tension-type headache: design of a randomised clinical trial. *BMC Musculoskelet Disord*. 2009;10:21.
3. Espi-López GV, Rodríguez-Blanco C, Oliva-Pascual-Vaca A, Benítez-Martínez JC, Lluch E, Falla D. The effect of manual therapy techniques on headache disability in patients with tension-type headache. *Eur J Phys Rehabil Med*. 2014;50:641–7.
4. Gemma V, Antonia GC. Efficacy of manual and manipulative therapy in the perception of pain and cervical motion in patients with tension-type headache: a randomized, controlled clinical trial. *J Chiropr Med*. 2014;13:4–13.
5. Castien RF, van der Windt DA, Blankenstein AH, Heymans MW, Dekker J. Clinical variables associated with recovery in patients with chronic tension-type headache after treatment with manual therapy. *Pain*. 2012;153:893–9.

L. Caballero-Calixto^{a,*}, P. Antón-Puecas^b,
J. Reyna-Ormeño^c

^a *Especialidad de Terapia Física, Facultad de Tecnología Médica, Universidad San Juan Bautista, Lima, Peru*

^b *Departamento de Fisioterapia, Hospital Central de la Fuerza Aérea del Perú, Lima, Peru*

^c *Escuela de Terapia Física, Universidad Peruana de Ciencias Aplicadas, Lima, Peru*

* Corresponding author.

E-mail address: lily.caballero6@gmail.com

(L. Caballero-Calixto).

2173-5808/

© 2015 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Superior vena cava syndrome as a complication of intravenous immunoglobulin treatment[☆]



Síndrome de vena cava superior como complicación del tratamiento con inmunoglobulinas intravenosas

Dear Editor:

Intravenous immunoglobulins are a frequent treatment for a variety of neurological diseases.¹ This treatment has been

associated with thromboembolic complications, with an incidence ranging between 1.2% and 11.3%.² The pathogenic mechanisms of the disease include increased plasma viscosity, increased platelet count and adhesion, and presence of procoagulant antibodies and coagulation factors not eliminated by immunoglobulin fractionation.^{2,3}

Superior vena cava thrombosis in patients receiving intravenous immunoglobulins is an infrequent complication that has rarely been described in the literature.⁴ Management of this complication may be difficult, especially in patients with a central venous catheter, since no specific management guidelines have been established to date.^{4,5}

We present the case of a 57-year-old female smoker diagnosed with chronic inflammatory demyelinating polyneuropathy. She was in treatment with azathioprine, deflazacort 6 mg/24 h, and intravenous immunoglobulins dosed at 0.5 g/kg/day and administered over 12 hours per day for 4 days every month. She had no other relevant history. Immunoglobulin therapy had been started 3 years earlier, by means of a subcutaneous reservoir attached to a catheter in the superior vena cava. The catheter had been implanted following multiple episodes of thrombophlebitis

[☆] Please cite this article as: Crespo-Burillo JA, Alarcia-Alejos R, Capablo-Liesa JL. Síndrome de vena cava superior como complicación del tratamiento con inmunoglobulinas intravenosas. *Neurología*. 2017;32:202–204.