

## Anticipating disaster: mechanical thrombectomy in patients with low NIHSS scores<sup>☆</sup>



### Anticipándonos al desastre: trombectomía mecánica en paciente con NIHSS bajo

Dear Editor,

The effectiveness of endovascular treatment for acute stroke is well established.<sup>1</sup> International treatment guidelines recommend endovascular treatment for patients with proximal intracranial artery occlusions and NIHSS scores  $\geq 6$ .<sup>2</sup> However, patients with proximal occlusions may initially score low on the NIHSS before the collapse of compensatory mechanisms and the onset of the severe symptoms usually associated with large-vessel occlusions.


We present the case of a patient arriving at the emergency department with ischaemic stroke, scoring 1 on the NIHSS. She had a thrombus in the terminal segment of the right internal carotid artery (ICA) and extending up to the M1 segment of the middle cerebral artery (MCA), which was extracted by mechanical thrombectomy.

### Clinical case

The patient was a 71-year-old woman with arterial hypertension and a history of atrial fibrillation managed with acenocoumarol. She was transported to the emergency department due to sudden onset 2 h earlier of difficulty uttering words, a tingling sensation in the left leg, and mild weakness in the left arm and leg. The neurological examination revealed mild paresis of the left leg and an NIHSS score of 1. A blood analysis showed an INR of 2.07. An emergency head CT scan revealed a hyperdense area in the right MCA (Fig. 1A), with no other signs of acute ischaemia or intracranial haemorrhage (ASPECTS score of 10). CT angiography revealed occlusion of the terminal segment of the right ICA.

Symptoms worsened during the patient's transfer to the operating theatre for neurointerventional surgery: she displayed left inferior homonymous quadrantanopia, left supranuclear facial palsy, left arm hypoaesthesia, and mild left-sided hemiparesis (NIHSS 5). Angiography confirmed the results of the CT angiography study (Fig. 1B and C); the thrombus was extracted by stent-retriever mechanical thrombectomy, achieving complete recanalisation (TICI 3) after a single pass.

The patient was asymptomatic (NIHSS 0) the following day and remained so until discharge.



**Figure 1** (A) Non-contrast head CT scan showing the “hyperdense middle cerebral artery sign” (arrow), suggestive of a thrombus at this level. Sagittal (B) and coronal (C) sections from the digital subtraction angiography study, revealing an obstruction to contrast flow in terminal internal carotid artery above the exit of the posterior communicating artery (arrow).

### Discussion

The American Heart Association clinical guidelines establish a minimum NIHSS score of 6 for patients with acute ischaemic stroke to be eligible for endovascular treatment (class I, level of evidence A).<sup>2</sup> However, occlusions of large intracranial arteries may be associated with low NIHSS scores. In the study by Maas et al.,<sup>3</sup> 29% of patients with an initial NIHSS score of 0 had proximal occlusions. This may be due to the fact that symptoms of ischaemic stroke appear when cerebral blood flow drops below 18–22 mL/100 g/min,<sup>4</sup> but such dynamic systems as arteriolar vasodilation and collateral circulation maintain cerebral blood flow above that threshold.

In 62%–71% of cases, these mechanisms eventually fail if blood flow is not restored in the artery involved, leading to symptom worsening and an increase in the size of the infarction.<sup>5–7</sup>

Intravenous fibrinolysis only achieves complete recanalisation of proximal occlusion in up to 30% of cases; the likelihood of spontaneous recanalisation is even lower. Bhogal et al.<sup>8</sup> and Pfaff et al.<sup>9</sup> studied the effectiveness of mechanical thrombectomy in patients with low NIHSS scores and report a median thrombus length of 12 mm (interquartile range, 10–12) and a mean length of 10 mm (range, 4.6–23.1), respectively. There is an extremely low likelihood of achieving recanalisation with intravenous fibrinolysis for thrombus lengths exceeding 8 mm.<sup>10</sup>

Performing mechanical thrombectomy in patients with proximal occlusion and presenting low NIHSS scores would improve outcomes: once collateral circulation collapses, symptoms worsen and infarction size increases at a faster pace than at stroke onset, since compensatory mechanisms are already exhausted.

Our patient's symptoms worsened during transfer to the operating theatre: she had an NIHSS score of 5 in the min-

<sup>☆</sup> Please cite this article as: Tejada Meza H, Martínez García R, Capablo Liesa JL, Marta Moreno J. Anticipándonos al desastre: trombectomía mecánica en paciente con NIHSS bajo. *Neurología*. 2019;34:350–351.

utes before the intervention, probably since compensatory mechanisms were collapsing.

Regarding functional prognosis, in the study by Bhogal et al.<sup>8</sup> (patients with M1 MCA occlusion and NIHSS scores  $\leq 5$  receiving endovascular treatment), around 75% of patients scored  $\leq 2$  on the modified Rankin Scale (mRS) at 3 months. A good functional prognosis is to be expected in our patient, since she was asymptomatic at discharge (mRS 0).

Revascularisation should be considered in all patients with proximal occlusions arriving early at the hospital, regardless of NIHSS score.

## References

- Goyal M, Menon BK, van Zwam WH, Dippel DW, Mitchell PJ, Demchuk AM, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet*. 2016;387:1723–31.
- Powers WJ, Derdeyn CP, Biller J, Coffey CS, Hoh BL, Jauch EC, et al. American Heart Association Stroke Council. 2015 American Heart Association/American Stroke Association Focused update of the 2013 guidelines for the early management of patients with acute ischemic stroke regarding endovascular treatment: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2015;46:3020–35.
- Maas MB, Furie KL, Lev MH, Singhal AB, Ay H, Greer DM, et al. National Institutes of Health Stroke Scale score is poorly predictive of proximal occlusion in acute cerebral ischemia. *Stroke*. 2009;40:2988–93.
- Lin MP, Liebeskind DS. Imaging of ischemic stroke. *Continuum (Minneapolis)*. 2016;22:1399–423.
- Alexandrov AV, Felberg RA, Demchuk AM, Christou I, Burgin WS, Malkoff M, et al. Deterioration following spontaneous improvement: sonographic findings in patients with acutely resolving symptoms of cerebral ischemia. *Stroke*. 2000;31:915–9.
- Campbell BCV, Christensen S, Tress BM, Churilov L, Desmond PM, Parsons MW, et al., EPITHET Investigators. Failure of collateral blood flow is associated with infarct growth in ischemic stroke. *J Cereb Blood Flow Metab*. 2013;33:1168–72.
- Miteff F, Levi CR, Bateman GA, Spratt N, McElduff P, Parsons MW. The independent predictive utility of computed tomography angiographic collateral status in acute ischaemic stroke. *Brain*. 2009;132:2231–8.
- Bhogal P, Bücke P, Ganslandt O, Bänzner H, Henkes H, Aguilar Pérez M. Mechanical thrombectomy in patients with M1 occlusion and NIHSS score  $\leq 5$ : a single-centre experience. *Stroke Vasc Neurol*. 2016:e000052, <http://dx.doi.org/10.1136/svn-2016-000052>.
- Pfaff J, Herweh C, Pham M, Schönenberger S, Nagel S, Ringleb PA, et al. Mechanical thrombectomy in patients with acute ischemic stroke and lower NIHSS scores: recanalization rates, periprocedural complications, and clinical outcome. *Am J Neuroradiol*. 2016;37:2066–71.
- Riedel CH, Zimmermann P, Jensen-Kondering U, Stingele R, Deuschl G, Jansen O. The importance of size: successful recanalization by intravenous thrombolysis in acute anterior stroke depends on thrombus length. *Stroke*. 2011;42:1775–7.

H. Tejada Meza<sup>a,\*</sup>, R. Martínez García<sup>b</sup>,  
J.L. Capablo Liesa<sup>c</sup>, J. Marta Moreno<sup>a</sup>

<sup>a</sup> *Unidad de Ictus, Servicio de Neurología, Hospital Universitario Miguel Servet, Zaragoza, Spain*

<sup>b</sup> *Servicio de Radiodiagnóstico, Hospital Universitario Miguel Servet, Zaragoza, Spain*

<sup>c</sup> *Servicio de Neurología, Hospital Universitario Miguel Servet, Zaragoza, Spain*

\*Corresponding author.

*E-mail address:* [hmeza@gmail.com](mailto:hmeza@gmail.com) (H. Tejada Meza).  
2173-5808/

© 2019 Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Neurología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).