



ORIGINAL ARTICLE

Assessment of the diagnostic accuracy and discriminative validity of the Clock Drawing and Mini-Cog tests in detecting cognitive impairment^{☆,☆☆}

C. Carnero-Pardo^{a,*}, I. Rego-García^b, J.M. Barrios-López^b, S. Blanco-Madera^b, R. Calle-Calle^b, S. López-Alcalde^a, R.M. Vilchez-Carrillo^b

^a FIDYAN Neurocenter, Granada, Spain

^b Servicio de Neurología, Hospital Universitario Virgen de las Nieves, Granada, Spain

Received 24 November 2018; accepted 22 December 2018

Available online 17 September 2021

KEYWORDS

Cognitive impairment;
Detection;
Mini-Cog;
Clock Drawing test;
Screening;
Diagnostic usefulness

Abstract

Introduction and objectives: The Mini-Cog is a very brief, widely used cognitive test that includes a memory task and a simplified assessment of the Clock Drawing Test (CDT). There is not a formal evaluation of the Mini-Cog test in Spanish. This study aims to analyse the diagnostic usefulness of the Mini-Cog and CDT for detecting cognitive impairment (CI).

Methods: We performed a cross-sectional study, systematically including all patients who consulted at our neurology clinic over a 6-month period. We assessed diagnostic usefulness for detecting CI (defined according to the National Institute on Aging-Alzheimer's Association criteria for mild cognitive impairment and dementia) according to the area under the receiver operating characteristic curve (AUC). Sensitivity, specificity, and positive and negative likelihood ratios were calculated for each cut-off point.

Results: The study included 581 individuals (315 with CI); 55.1% were women and 27.7% had not completed primary studies. The Mini-Cog showed greater diagnostic usefulness than the CDT (AUC ± sensitivity: 0.88 ± 0.01 vs 0.84 ± 0.01; $P < .01$). Both instruments were less useful for screening in individuals with a low education level (0.74 ± 0.05 vs 0.75 ± 0.05, respectively). A cut-off point of 2/3 in the Mini-Cog achieved a sensitivity of 0.90 (95% CI, 0.87-0.93) and a specificity of 0.71 (95% CI, 0.65-0.76); a cut-off point of 5/6 in the CDT achieved a sensitivity of 0.77 (95% CI, 0.72-0.81) and a specificity of 0.80 (95% CI, 0.75-0.85).

[☆] Please cite this article as: Carnero-Pardo C, Rego-García I, Barrios-López JM, Blanco-Madera S, Calle-Calle R, López-Alcalde S, et al. Evaluación de la utilidad diagnóstica y validez discriminativa del Test del Reloj y del Mini-Cog en la detección del deterioro cognitivo. *Neurología*. 2022;37:13–20.

^{☆☆} An interim analysis of this study was presented orally at the 70th Annual Meeting of the Spanish Society of Neurology.

* Corresponding author.

E-mail address: ccarnero@neurocenter.es (C. Carnero-Pardo).

Conclusion: In our neurology clinic, the Mini-Cog showed acceptable diagnostic usefulness for detecting CI, greater than that of the CDT; neither test is an appropriate instrument for individuals with a low level of education.

© 2019 Sociedad Española de Neurología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Deterioro cognitivo;
Detección;
Mini-Cog;
Test del Reloj;
Cribado;
Utilidad diagnóstica

Evaluación de la utilidad diagnóstica y validez discriminativa del Test del Reloj y del Mini-Cog en la detección del deterioro cognitivo

Resumen

Introducción y Objetivos: El Mini-Cog es un test cognitivo muy breve de uso extendido que incluye una tarea de memoria y una evaluación simplificada del Test del Reloj (TdR). No existe una evaluación formal del Mini-Cog en español; nuestro objetivo es analizar la utilidad diagnóstica (UD) del Mini-Cog y del TdR para deterioro cognitivo (DC).

Métodos: Estudio transversal en el que se han incluido de forma sistemática todos los sujetos atendidos durante un semestre en una consulta de Neurología. La UD se ha evaluado para DC (incluye sujetos con criterios NIA-AA de *mild cognitive impairment* o demencia) por medio del área bajo la curva ROC (aROC). Se han calculado los parámetros de sensibilidad (S), especificidad (E) y cocientes de probabilidad positivo y negativo (CP+, CP-) para los distintos puntos de corte.

Resultados: Se han incluido 581 sujetos (315 DC), 55.1% mujeres y 27.7% con bajo nivel educativo (< estudios primarios). La UD del Mini-Cog es superior a la del TdR (0.88 ± 0.01 (aROC \pm ee) vs 0.84 ± 0.01 , $P < .01$); para ambos instrumentos, la UD disminuye notablemente en sujetos con bajo nivel educativo (0.74 ± 0.05 y 0.75 ± 0.05 respectivamente). El punto de corte 2/3 del Mini-Cog tiene una S 0.90 (0.87–0.93) y una E 0.71 (0.65–0.76) y el 5/6 del TdR una S 0.77 (0.72–0.81) y E 0.80 (0.75–0.85).

Conclusiones: En consulta de Neurología, el Mini-Cog tiene una UD para DC aceptable, superior a la del TdR; ninguno de ellos es un instrumento adecuado para ser utilizado en sujetos con bajo nivel educativo.

© 2019 Sociedad Española de Neurología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Subjective cognitive complaints and memory loss constitute a frequent reason for consultation with primary care and referral to the neurology department.¹ Initial assessment and follow-up of these patients requires the use of valid cognitive assessment tools that are brief enough to be administered during consultations.²

The Mini-Mental State Examination (MMSE),³ currently the most frequently used cognitive assessment tool, is rapidly losing supporters due to numerous limitations, including its cost,⁴ and was even excluded from the third version of the Alzheimer Disease Centers' Neuropsychological Test Battery.⁵ The Montreal Cognitive Assessment,⁶ in contrast, is increasingly popular. It includes multiple domains, but has an even longer administration time than the MMSE, which limits its use in general consultations and particularly in primary care.

The clock drawing test (CDT), one of the oldest and most widely used brief cognitive tests,⁷ is considered by some authors to be the ideal cognitive screening test due to its simplicity and brevity.⁸ However, it does present some limitations: it does not explicitly evaluate episodic memory; it requires a minimum level of graphomotor skills (which may be problematic for illiterate individuals or those with a low level of education)⁹; it does not have a simple, standardised grading system¹⁰; and its diagnostic accuracy is limited.¹¹

The Mini-Cog,¹² a simple, quick-to-administer instrument, aims to address some of the limitations of the CDT: it includes a 3-item free recall task and a simplified clock-drawing task.⁸ It has been validated in multiple settings (community,¹³ primary care,¹⁴ specialised clinics¹⁵) and in numerous languages (available at <https://mini-cog.com/mini-cog-in-other-languages/>). Furthermore, there are 15 versions of the Mini-Cog, which differ in the set of items the patient has to recall (5 options) and the time to which the clock has to be set (3 options).¹⁶ The Mini-Cog is currently one of the most widely recommended dementia screening tools for primary care and general consultations.^{17–22}

The Mini-Cog has been used in some studies conducted in Spanish-speaking countries,^{23,24} although no formal evaluation of the Spanish-language version has been conducted to date. In Spain, one study did evaluate the test's diagnostic usefulness, calculating results retrospectively by adding together scores for CDT drawings and the MMSE recall task, which had previously been administered independently; however, the Mini-Cog was not actually administered.²⁵

The purpose of this study was to formally evaluate the discriminant validity and diagnostic usefulness of the CDT and the Mini-Cog for identifying individuals with cognitive impairment in a large sample of patients attended at the neurology clinic.

Methods

Design

We conducted a cross-sectional study including all patients attended between March and September 2018 at a neurology clinic specialising in cognitive and behavioural neurology.

For patients attended on more than one occasion during the study period, we only included the data corresponding to the last evaluation, as these were considered to be the most reliable from a diagnostic viewpoint.

Cognitive assessment

All patients underwent a brief cognitive assessment including the Mini-Cog, the CDT, and the Fototest,²⁶ regardless of the reason for consultation.

All patients attended due to subjective or informant-reported memory complaints and/or behavioural alterations, or those scoring below the 10th percentile on the Fototest,²⁷ underwent a thorough cognitive assessment including orientation (temporal, spatial, and personal), attention (forward and backward digit span), memory (learning, free recall, and recognition of the CERAD word list²⁸), language (short version of the Boston Naming Test,²⁹ semantic verbal fluency,³⁰ comprehension of commands), motor apraxia (EULA test³¹), executive function (WAIS similarities, coin test from the Eurotest,³² semantic verbal fluency), and visuospatial function (copy and drawing), in addition to a formal functional evaluation. The researcher performing the assessment (SLA) was blinded to Mini-Cog, CDT, and Fototest results.

Mini-Cog

Version used

We used the Mini-Cog¹⁶ version including the words “capitán,” “jardín,” and “retrato” (captain, garden, picture), the word combination showing the most homogeneous frequency of use in Spanish (58.67, 62.45, and 38.42 occurrences per million words, respectively),³³ and the clock time “10 past 11,” the most frequently used clock time in our setting.

Administration

Patients were given the following instructions: “Please listen carefully. I am going to say 3 words that I want you to repeat back to me; try to remember them as I will ask you to repeat them again.” If the person was unable to repeat all 3 words, the instructions were repeated up to 3 times. Patients were then given a blank sheet and a pencil, and the following instructions were given: “Next, I want you to draw a large, round clock for me. First, put all of the numbers in their places. Then, set the hands to 10 past 11.” After the clock-drawing task was completed, the patient was asked to recall the 3 words from the first task.

Scoring

In the recall task, one point was scored for each word that was correctly remembered. The clock drawing task was scored the maximum score (2 points) if the drawing met all of the following conditions:

- 1 The clock has all the numbers, with no repetitions and no additional numbers.
- 2 The numbers are placed in the correct order.
- 3 The numbers are placed in the correct positions on the face of the clock.
- 4 The hands are unequivocally set to the time asked.

Any other situation, including the patient refusing to draw the clock or being unable to do so due to physical limitations (paresis, tremor, ataxia, etc), was scored 0 points.

Registry

We recorded patients’ scores, the time taken to complete the test, the total number of attempts to repeat the 3 words, and the number of times the rater had to repeat the instructions for the clock drawing task.

Clock drawing test

We used the scoring system of the Spanish-language version of the Seven-Minute Screen³⁴ to score the clock drawing task in the Mini-Cog; this system scores one point for each of the following criteria:

- 1 The clock has all numbers from 1 to 12 (using either Arabic or Roman numerals).
- 2 The numbers are placed in the correct ascending order (even if not all 12 numbers have been placed).
- 3 The numbers are in the correct positions.
- 4 The clock has 2 hands (circles or other marks are not valid).
- 5 The hour hand (or any other mark) is set to 11.
- 6 The minute hand (or any other mark) is set to 10.
- 7 The hour hand is shorter than the minute hand (or the patient indicates so orally).

Diagnosis

Regardless of the reason for consultation and the final aetiological diagnosis, all patients were categorised by cognitive status, as follows:

- Normal cognitive function (NCF): no subjective cognitive complaints and age-, sex-, and education-adjusted Fototest scores above the 10th percentile.²⁷
- Subjective cognitive complaints (SCC): presence of subjective cognitive complaints, with normal results in the formal cognitive assessment.
- Cognitive impairment without associated functional limitations (Clw/o): presence of alterations in at least one cognitive domain according to the formal cognitive assessment, with no significant impact on functional status; these individuals meet the National Institute of Aging–Alzheimer’s Association criteria for mild cognitive impairment.³⁵
- Cognitive impairment with associated functional limitations (Clw): presence of alterations in at least one cognitive domain according to the formal cognitive assessment, with significant impact on functional status; these individuals meet the National Institute of Aging–Alzheimer’s Association criteria for dementia.³⁶ In line with the recommendations of other authors,³⁷ we avoid using the term “dementia” due to its negative connotations and the associated stigma.

Statistical analysis

We conducted a descriptive study of sociodemographic variables and results, stratified by diagnosis; comparisons were performed using ANOVA and the chi-square test, depending on whether variables were continuous or categorical.

The diagnostic usefulness of the tests was evaluated with the area under the ROC curve in patients with (Clw/o + Clw) vs without cognitive impairment (NCF + SCC). We calculated the sensitivity, specificity, and positive and negative likelihood ratios for different cut-off points. We used the Hanley and McNeil method for comparing ROC curves from the same sample³⁸ and the DeLong method for comparing ROC curves from independent samples.³⁹ All parameters

Table 1 Sociodemographic characteristics and cognitive test results.

	NCF	SCC	Clw/o	Clw	Total	No CI	CI
Participants, <i>n</i>	172	94	127	188	581	266	315
Age, years (<i>n</i> = 579)	47.9 (18.5)	63.7 (11.4)	70.5 (12.6)	73.9 (9.4)	63.8 (14.4)	53.4 (18.0)	72.5 (10.9)
Sex, women (<i>n</i> = 568)	102 (60.4%)	53 (57.6%)	63 (50.0%)	95 (52.5%)	313 (55.1%)	155 (59.4%)	158 (51.5%)
Education level (<i>n</i> = 573)							
< Primary education	15 (8.7%)	23 (25.3%)	46 (37.4%)	75 (40.1%)	159 (27.7%)	38 (14.4%)	121 (39.0%)
≥ Primary education	157 (91.3%)	68 (74.8%)	77 (62.6%)	112 (59.9%)	414 (72.3%)	226 (85.6%)	189 (61.0%)
Illiterate (<i>n</i> = 572)	2 (1.2%)	2 (2.2%)	4 (3.2%)	17 (9.1%)	25 (4.4%)	4 (1.5%)	21 (6.8%)
Clock drawing test (<i>n</i> = 578)	6.3 (1.4)	5.7 (1.7)	4.0 (2.2)	2.4 (2.4)	4.4 (2.6)	6.0 (1.5)	3.1 (2.4)
Mini-Cog (<i>n</i> = 581)	3.6 (1.3)	2.4 (0.9)	0.9 (1.1)	0.6 (1.1)	1.9 (1.8)	3.2 (1.5)	0.7 (1.1)
Time, seconds (<i>n</i> = 517)	85.7 (28.6)	106.7 (28.9)	119.0 (33.6)	119.9 (36.2)	107.6 (36.2)	93.2 (30.4)	119.5 (36.3)

CI: cognitive impairment; Clw: cognitive impairment with associated functional limitations; Clw/o: cognitive impairment without associated functional limitations; NCF: normal cognitive function; SCC: subjective cognitive complaints.

Data are expressed either as absolute frequency (%) or mean (standard deviation).




Figure 1 Violin plot showing Mini-Cog scores, stratified by cognitive status. The white dots represent median scores, the ends of the black bars represent the 25th and 75th percentiles, and the length of the line is 1.5 times the interquartile range; the violin shape represents the density distribution of the data, extending to the extreme high and low values. Clw: cognitive impairment with associated functional limitations; Clw/o: cognitive impairment without associated functional limitations; NCF: normal cognitive function; SCC: subjective cognitive complaints.

were calculated with 95% confidence intervals and two-way comparisons with an alpha error of 0.05. Statistical analysis was performed with MedCalc, version 18.9.1.⁴⁰

Formal considerations

The study design and manuscript preparation follow the STARD recommendations for studies evaluating diagnostic tests.⁴¹

Results

We included a total of 581 individuals, 266 of whom (45.8%) did not have cognitive impairment; 94 of these (16.2%) had SCC. The remaining 315 (54.2%) had cognitive impairment: 127 (21.9%) had no functional limitations and 188 (32.1%) presented functional limitations. Table 1 summarises the sociodemographic characteristics and cognitive assessment results of our sample. Participants had a mean (SD) age of 63.8 (14.4) years (range, 15–92) and 55.1% were women. No differences in sex distribution were observed between




Figure 2 Violin plot showing clock drawing test scores, stratified by cognitive status. The white dots represent median scores, the ends of the black bars represent the 25th and 75th percentiles, and the length of the line is 1.5 times the interquartile range; the violin shape represents the density distribution of the data, extending to the extreme high and low values. Clw: cognitive impairment with associated functional limitations; Clw/o: cognitive impairment without associated functional limitations; NCF: normal cognitive function; SCC: subjective cognitive complaints.

groups ($\chi^2 = 3.94$; not significant) but we did observe differences in age, which increased with poorer cognitive status ($F = 121.9$; $P < .001$), and education level, which was lower in individuals with poorer cognitive status ($\chi^2 = 53.68$; $P < .001$). We also observed an inverse correlation between the percentage of illiterate individuals and cognitive status ($\chi^2 = 61.17$; $P < .001$).

Cognitive test results showed a significant negative correlation with cognitive status ($F = 210.7$ for the Mini-Cog and $F = 126.9$ for the CDT; $P < .0001$ for both) (Figs. 1 and 2).

Individuals with cognitive impairment completed the Mini-Cog in a mean of 119.5 (36.3) seconds (range, 55–266), nearly half a minute longer than it took individuals without cognitive impairment to complete the test (93.2 [30.4] seconds [range, 49–251]); the difference is highly significant ($F = 34.2$; $P < .0001$).

The diagnostic usefulness (area under the ROC curve \pm standard error) of the Mini-Cog was significantly higher than that of the CDT (0.88 ± 0.01 vs 0.84 ± 0.01 ; $P < .01$) (Fig. 3). The diagnostic usefulness of the Mini-Cog and the CDT was highly influenced by level of education; performance was clearly poorer among individuals without primary studies (0.74 ± 0.05 and 0.75 ± 0.05 , respectively) than




Figure 3 ROC curves for the Mini-Cog and the clock drawing test (CDT) for cognitive impairment. Data represent the area under the ROC curve ± standard error.

among individuals with at least primary studies (0.90 ± 0.01 and 0.84 ± 0.02 , respectively) (Table 2).

With the optimal cut-off point (2/3), the Mini-Cog has greater sensitivity (0.90) than specificity (0.71), correctly classifying 81.4% of the sample (Table 3); the CDT (cut-off point of 5/6, the score with the best diagnostic performance), in contrast, has greater specificity (0.80) than sensitivity (0.77), correctly classifying only 78.5% of individuals (Table 4).

Discussion

Our study included a large, heterogeneous sample of individuals attended at a neurology clinic, with a high prevalence of cognitive problems. In our sample, the Mini-Cog showed clearly greater

diagnostic usefulness than the CDT; for both tests, the diagnostic usefulness is highly influenced by the patient education level.

The CDT presents limited diagnostic usefulness for cognitive impairment (area under the ROC curve: 0.84 ± 0.02); this is consistent with the results reported in previous studies, regardless of the correction method used, the setting, or the geographical location of the sample. Lee et al.⁴² and Duro et al.⁴³ evaluated the diagnostic usefulness of the CDT for MCI using different correction methods; Ehreke et al.⁴⁴ conducted a systematic review and concluded that the CDT is not suitable for cognitive impairment screening. A previous study conducted by our research group, including patients from primary care consultations and using the same grading system, shows identical results (0.84 ± 0.02); however, the study conducted by Cacho et al.,⁴⁵ including patients attended at a neurology department and using a 10-point grading system,⁴⁶ reports an even lower diagnostic usefulness for cognitive impairment (0.78).

The limited usefulness of the CDT may be due to the fact that the test does not specifically evaluate memory, the cognitive domain most frequently affected in patients with cognitive impairment; this would explain the superiority of the Mini-Cog, whose main difference with respect to the CDT is the inclusion of a free recall task.

The diagnostic usefulness of the Mini-Cog in our setting is acceptable (area under the ROC curve: 0.88 ± 0.02), similarly to the results of a study including a multiethnic sample with high prevalence of cognitive impairment (62.3%),⁴⁷ and was clearly superior to the values obtained in 2 studies conducted in the primary care setting (with areas under the ROC curve of 0.66 and 0.77, respectively).^{20,25} The main difference between this study and our previous study²⁵ is that in the previous study, Mini-Cog scores were calculated retrospectively by combining the results of the CDT and the recall task of the MMSE, using a different word combination (bicycle, horse, apple) and clock time (20 to 8) in part of the sample; this may have led to an underestimation of the test’s usefulness.

Both the Mini-Cog and the CDT also present other limitations. Firstly, they have a narrow score range (0-5 for the Mini-Cog and 0-7 for the CDT), which may favour ceiling or floor effects, as observed in Figs. 1 and 2. Furthermore, and more importantly, both tests require minimum graphomotor skills, and are therefore unsuitable for illiterate individuals or those with a low education level; this may explain why the diagnostic usefulness of both tools is limited in individuals who had not completed primary education. These tools should therefore not be used in this population group. This is consis-

Table 2 Diagnostic usefulness of the Mini-Cog and the clock drawing test.

Test	Total (n = 581)	< Primary education (n = 159)	≥ Primary education (n = 412)
Mini-Cog	0.88 ± 0.01	0.74 ± 0.05	0.90 ± 0.01
CDT	0.84 ± 0.02	0.75 ± 0.05	0.84 ± 0.02

CDT: clock drawing test.
Data represent the area under the ROC curve ± standard error.

Table 3 Discriminant validity of the Mini-Cog in detecting cognitive impairment.

COP	Sen.	Sp.	+LR	-LR
≤ 0	0.64 (0.58-0.69)	0.93 (0.89-0.96)	9.4	0.39
≤ 1	0.75 (0.70-0.80)	0.85 (0.80-0.89)	4.88	0.29
≤ 2	0.90 (0.87-0.93)	0.71 (0.65-0.76)	3.09	0.13
≤ 3	0.97 (0.95-0.99)	0.48 (0.42-0.54)	1.87	0.06
≤ 4	0.99 (0.98-1.00)	0.24 (0.19-0.30)	1.31	0.01
≤ 5	1.00 (0.99-1.00)	0.00 (0.00-0.03)	1.00	

+LR: positive likelihood ratio; -LR: negative likelihood ratio; COP: cut-off point; Sen.: sensitivity; Sp.: specificity.
Numbers shown in parentheses are 95% confidence intervals.

Table 4 Discriminant validity of the clock drawing test in detecting cognitive impairment.

COP	Sen.	Sp.	+LR	−LR
≤ 0	0.25 (0.20-0.30)	0.97 (0.95-0.99)	9.4	0.77
≤ 1	0.33 (0.27-0.38)	0.96 (0.93-0.98)	8.55	0.70
≤ 2	0.45 (0.40-0.51)	0.95 (0.92-0.97)	9.25	0.57
≤ 3	0.56 (0.50-0.62)	0.94 (0.42-0.54)	8.77	0.47
≤ 4	0.69 (0.64-0.74)	0.88 (0.84-0.92)	5.93	0.35
≤ 5	0.77 (0.72-0.81)	0.80 (0.75-0.85)	3.85	0.29
≤ 6	0.89 (0.85-0.92)	0.54 (0.48-0.61)	1.97	0.20
≤ 7	1.00 (0.99-1.00)	0.00 (0.00-0.01)	1.00	

+LR: positive likelihood ratio; −LR: negative likelihood ratio; COP: cut-off point; Sen.: sensitivity; Sp.: specificity. Numbers shown in parentheses are 95% confidence intervals.

tent with the results of other studies evaluating the performance of the Mini-Cog in samples of individuals with low education levels.^{48,49}

The main strength of these tools is their brevity. In our study, the maximum completion time was 266 seconds, and 95% of participants completed the Mini-Cog in less than 3 minutes (95th percentile: 178.2 seconds); in fact, mean (SD) completion time for patients with cognitive impairment was less than 2 minutes (119.4 [36.3]). These results are very similar to those reported in a study including individuals attended at a primary care consultation,⁵⁰ who completed the Mini-Cog in a mean time of 118 (45) seconds, with no participant requiring longer than 5 minutes.

One of the strengths of this study is the large size and naturalistic character of the sample, since we systematically included all individuals attended at a neurology clinic, regardless of the reason for consultation. This recruitment method limits the risk of selection bias and increases the external validity of our results. Its weaknesses include the fact that controls did not undergo a formal cognitive assessment; presence of cognitive impairment was determined based on the clinical history and Fototest results.²⁶ Though valid, this tool cannot rule out the presence of false negatives, making it difficult to discriminate between patients with and without cognitive impairment. The risk of false positives was not a concern since all participants underwent a formal cognitive assessment. Finally, the functions of the neurology clinic where the study was conducted fall between those of a unit specialising in cognitive impairment and those of a general neurology consultation, which limits the applicability of our results to other settings, such as primary care or the general population.

In conclusion, the Mini-Cog has greater diagnostic usefulness than the CDT in screening for cognitive impairment in neurology consultations, but is unsuitable for the assessment of individuals with low levels of education. The test's simplicity and speed of use make it particularly useful for application in combination with other brief cognitive tests.²

Conflicts of interest

C. Carnero Pardo created the Fototest and has received professional fees for academic and consulting activities for Nutricia, Schwabe Farma Ibérica, Biogen, Piramal, Janssen Cilag, Pfizer, Eisai, Esteve, Novartis, Lundbeck, and Grunenthal.

Under the terms of its Creative Commons licence, Fototest may be used and distributed for non-commercial purposes provided that it is not modified and its authorship is explicitly acknowledged.

References

- Lopez-Pousa S, Monserrat-Vila S, Turro-Garriga O, Aguilar-Barbera M, Caja-Lopez C, Vilalta-Franch J, et al. Análisis de la demanda asistencial neurológica generada por la atención primaria en una área geográfica de las comarcas de Girona. *Rev Neurol.* 2009;49:288–94.
- Olazarán J, Hoyos-Alonso MC, del Ser T, Garrido Barral A, Conde-Sala JL, Bermejo-Pareja F, et al. Aplicación práctica de los test cognitivos breves. *Neurologia.* 2016;31:183–94.
- Folstein MF, Folstein SE, McHugh PR. 'Mini-mental state'. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12:189–98.
- Carnero-Pardo C. ¿Es hora de jubilar al Mini-Mental? *Neurologia.* 2014;29:473–81.
- Weintraub S, Besser L, Dodge HH, Teylan M, Ferris S, Goldstein FC, et al. Version 3 of the Alzheimer Disease Centers' Neuropsychological Test Battery in the Uniform Data Set (UDS). *Alzheimer Dis Assoc Disord.* 2018;32:10–7.
- Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc.* 2005;53:695–9.
- Hazan E, Frankenburg F, Brenkel M, Shulman K. The test of time: a history of clock drawing. *Int J Geriatr Psychiatry.* 2018;33:e22–30.
- Shulman KI. Clock-drawing: is it the ideal cognitive screening test? *Int J Geriatr Psychiatry.* 2000;15:548–61.
- Cavalcanti de Noronha I, dos Santos Barreto S, Zazo Ortiz K. The influence of education on performance of adults on the Clock Drawing Test. *Dement Neuropsychol.* 2018;12:61–7.
- Spenciere B, Alves H, Charchat-Fichman H. Scoring systems for the Clock Drawing Test: a historical review. *Dement Neuropsychol.* 2017;11:6–14.
- Park J, Jeong E, Seomun G. The clock drawing test: a systematic review and meta-analysis of diagnostic accuracy. *J Adv Nurs.* 2018;74:2742–54.
- Borson S, Scanlan J, Brush M, Vitaliano P, Dokmak A. The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry.* 2000;15:1021–7.
- Fage BA, Chan CC, Gill SS, Noel-Storr AH, Herrmann N, Smailagic N, et al. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a community setting. *Cochrane Database Syst Rev.* 2015:CD010860.
- Seitz DP, Chan CC, Newton HT, Gill SS, Herrmann N, Smailagic N, et al. Mini-Cog for the diagnosis of Alzheimer's disease dementia and other dementias within a primary care setting. *Cochrane Database Syst Rev.* 2018;2:CD011415.
- Milian M, Leiherr AM, Straten G, Muller S, Leyhe T, Eschweiler GW. The Mini-Cog versus the Mini-Mental State Examination

- and the Clock Drawing Test in daily clinical practice: screening value in a German Memory Clinic. *Int Psychogeriatr.* 2012;24:766–74.
16. McCarten JR, Anderson P, Kuskowski MA, McPherson SE, Borson S. Screening for cognitive impairment in an elderly veteran population: acceptability and results using different versions of the Mini-Cog. *J Am Geriatr Soc.* 2011;59:309–13.
 17. Brodaty H, Low LF, Gibson L, Burns K. What is the best dementia screening instrument for general practitioners to use? *Am J Geriatr Psychiatry.* 2006;14:391–400.
 18. Milne A, Culverwell A, Guss R, Tuppen J, Whelton R. Screening for dementia in primary care: a review of the use, efficacy and quality of measures. *Int Psychogeriatr.* 2008;20:911–26.
 19. Kamenski G, Dorner T, Lawrence K, Psota G, Rieder A, Schwarz F, et al. Detection of dementia in primary care: comparison of the original and a modified Mini-Cog Assessment with the Mini-Mental State Examination. *Ment Health Fam Med.* 2009;6:209–17.
 20. Holsinger T, Plassman BL, Stechuchak KM, Burke JR, Coffman CJ, Williams JW Jr. Screening for cognitive impairment: comparing the performance of four instruments in primary care. *J Am Geriatr Soc.* 2012;60:1027–36.
 21. Tsoi KK, Chan JY, Hirai HW, Wong SY, Kwok TC. Cognitive tests to detect dementia: a systematic review and meta-analysis. *JAMA Intern Med.* 2015;175:1450–8.
 22. Lin JS, O'Connor E, Rossom RC, Perdue LA, Eckstrom E. Screening for cognitive impairment in older adults: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med.* 2013;159:601–12.
 23. Dechent Rivera C, Bozani Leal A, Jorín Aliste M, Fuentes Gugliemetti P. Utilidad clínica del Memory Impairment Scree (MIS): análisis del instrumento como prueba de tamizaje de memoria. *Rev Chil Neuropsicol.* 2013;8:46–9.
 24. Wong-Achi X, Egas G, Cabrera D. Cognitive Impairment in Rural Elderly Population in Ecuador. *J Neurosci Rural Pract.* 2017;8:S20–2.
 25. Carnero-Pardo C, Cruz-Orduna I, Espejo-Martinez B, Martos-Aparicio C, Lopez-Alcalde S, Olazaran J. Utility of the mini-cog for detection of cognitive impairment in primary care: data from two spanish studies. *Int J Alzheimers Dis.* 2013;2013:285462.
 26. Carnero Pardo C, Saez-Zea C, Montiel Navarro L, Del Sazo P, Feria Vilar I, Perez Navarro MJ, et al. Utilidad diagnóstica del Test de las Fotos (Fototest) en deterioro cognitivo y demencia. *Neurologia.* 2007;22:860–9.
 27. Carnero Pardo C, Carrera Munoz I, Triguero Cueva L, Lopez Alcalde S, Vilchez Carrillo R. Valores normativos del Fototest en pacientes neurológicos sin deterioro cognitivo. *Neurologia.* 2018, <http://dx.doi.org/10.1016/j.nrl.2018.03.001>.
 28. Morris JC, Mohs RC, Rogers H, Fillenbaum G, Heyman A. Consortium to establish a registry for Alzheimer's disease (CERAD) clinical and neuropsychological assessment of Alzheimer's disease. *Psychopharmacol Bull.* 1988;24:641–52.
 29. Calero MD, Arnedo ML, Navarro E, Ruiz-Pedrosa M, Carnero C. Usefulness of a 15-item version of the Boston Naming Test in neuropsychological assessment of low-educational elders with dementia. *J Gerontol B Psychol Sci Soc Sci.* 2002;57:P187–91.
 30. Carnero Pardo C, Lendínez-González A. Utilidad del test de fluencia verbal semántica en el diagnóstico de demencia. *Rev Neurol.* 1999;29:709–14.
 31. Perez-Marmol JM, Lopez-Alcalde S, Carnero-Pardo C, Canadas-De la Fuente GA, Peralta-Ramirez MI, Garcia-Rios MC. Creation and design of a test for the Evaluation of Upper Limb Apraxia (EULA) based on a cognitive model: a pilot study. *Rev Neurol.* 2015;60:66–74.
 32. Carnero-Pardo C, Montoro-Rios MT. Evaluación preliminar de un nuevo test de cribado de demencia (Eurotest). *Rev Neurol.* 2004;38:201–9.
 - [33]. Real Academia Española. Banco de datos (Corpus de referencia del español actual, versión anotada) [online] [Accessed 22/10/18]. Available from: <http://www.rae.es>.
 34. del Ser Quijano T, Sanchez Sanchez F, Garcia de Yebenes MJ, Otero Puime A, Zunzunegui MV, Munoz DG. Versión española del Test de los 7 Minutos. Datos normativos de una muestra poblacional de ancianos de más de 70 años. *Neurologia.* 2004;19:344–58.
 35. Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* 2011;7:270–9.
 36. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR Jr, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* 2011;7:263–9.
 37. Trachtenberg DI, Trojanowski JQ. Dementia: a word to be forgotten. *Arch Neurol.* 2008;65:593–5.
 38. Hanley JA, McNeil BJ. A method of comparing the areas under receiver operating characteristic curves derived from the same cases. *Radiology.* 1983;148:839–43.
 39. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics.* 1988;44:837–45.
 - [40]. MedCalc Statistical Software versión 18.9.1 [computer program]. Ostend, Belgium: Published: MedCalc Software bvba; 2018 <http://www.medcalc.org>
 41. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, et al. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. *BMJ.* 2003;326:41–4.
 42. Lee KS, Kim EA, Hong CH, Lee DW, Oh BH, Cheong HK. Clock drawing test in mild cognitive impairment: quantitative analysis of four scoring methods and qualitative analysis. *Dement Geriatr Cogn Disord.* 2008;26:483–9.
 43. Duro D, Freitas S, Tabuas-Pereira M, Santiago B, Botelho MA, Santana I. Discriminative capacity and construct validity of the Clock Drawing Test in Mild Cognitive Impairment and Alzheimer's disease. *Clin Neuropsychol.* 2018:1–17.
 44. Ehreke L, Luppá M, König HH, Riedel-Heller SG. Is the Clock Drawing Test a screening tool for the diagnosis of mild cognitive impairment? A systematic review. *Int Psychogeriatr.* 2010;22:56–63.
 45. Cacho J, Benito-Leon J, Garcia-Garcia R, Fernandez-Calvo B, Vicente-Villardón JL, Mitchell AJ. Does the combination of the MMSE and clock drawing test (mini-clock) improve the detection of mild Alzheimer's disease and mild cognitive impairment? *J Alzheimers Dis.* 2010;22:889–96.
 46. Sunderland T, Hill JL, Mellow AM, Lawlor BA, Gundersheimer J, Newhouse PA, et al. Clock drawing in Alzheimer's disease. A novel measure of dementia severity. *J Am Geriatr Soc.* 1989;37:725–9.
 47. Borson S, Scanlan JM, Watanabe J, Tu SP, Lessig M. Simplifying detection of cognitive impairment: comparison of the Mini-Cog and Mini-Mental State Examination in a multiethnic sample. *J Am Geriatr Soc.* 2005;53:871–4.

48. Lourenco RA, Filho Ribeiro ST. The accuracy of the Mini-Cog in screening low-educated elderly for dementia. *J Am Geriatr Soc.* 2006;54:376–7, author reply 7-8.
49. Ribeiro Filho ST, Lourenco RA. The performance of the Mini-Cog in a sample of low educational level elderly. *Dement Neuropsychol.* 2009;3:81–7.
50. Sinclair AJ, Gadsby R, Hillson R, Forbes A, Bayer AJ. Brief report: use of the Mini-Cog as a screening tool for cognitive impairment in diabetes in primary care. *Diabetes Res Clin Pract.* 2013;100:e23–5.