



REVISTA PAULISTA DE PEDIATRIA

www.rpped.com.br



ORIGINAL ARTICLE

Nutritional status, nutritional self-perception, and use of licit drugs in adolescents



Denise Máximo Garcia, Eduardo Mekitarian Filho*, Alfredo Elias Gilio, João Paulo Becker Lotufo, Denise Swei Lo

Universidade de São Paulo (USP), São Paulo, SP, Brazil

Received 11 September 2014; accepted 26 November 2014
Available online 29 June 2015

KEYWORDS

Body weight;
Alcohol drinking;
Tobacco use

Abstract

Objective: To associate the nutritional status and the self-perception of nutritional status with the use of licit drugs among adolescents.

Methods: A cross-sectional study was conducted in which 210 adolescents answered a questionnaire on alcohol and tobacco experimentation and self-perceptions about their nutritional status. The correspondence between the adolescents' perception of their own nutritional status and actual nutritional status was analyzed, as well as associations between nutritional status, self-perception of nutritional status, gender, age, and presence of smokers at home with alcohol and tobacco use. The variables were analyzed separately in a bivariate analysis and, subsequently, a multivariate analysis determined the factors associated with drug use.

Results: The study included 210 adolescents with a median age of 148 months; 56.6% were females. Of the total sample, 6.6% have tried cigarettes, and 20% have tried alcohol; 32.3% had BMI Z-Score ≥ 1 , 12.85% had BMI Z-Score ≥ 2 , and 50.7% had a correct perception of his/her weight. After a multivariate analysis, only the self-perception about weight statistically influenced experimentation of tobacco, and patients who identified themselves as having very high weight were more likely to experiment tobacco (*odds ratio* (OR) 13.57; confidence interval (95% CI) 2.05–89.8; $p=0.007$); regarding alcohol use, adolescents who identified themselves as having high weight were 2.4 times more likely to experiment with alcohol than adolescents that identified themselves as having normal weight (95% CI 1.08–5.32, $p=0.031$).

Conclusions: Adolescents with self-perception of excess weight may constitute a risk group for alcohol and tobacco use.

© 2015 Sociedade de Pediatria de São Paulo. Published by Elsevier Editora Ltda. All rights reserved.

DOI of original article: <http://dx.doi.org/10.1016/j.rpped.2014.11.015>

* Corresponding author.

E-mail: emf2002@uol.com.br (E. Mekitarian Filho).

PALAVRAS-CHAVE

Peso corporal;
Alcoolismo;
Hábito de fumar

Estado nutricional, autopercepção do estado nutricional e experimentação de drogas lícitas em adolescentes**Resumo**

Objetivo: Associar o estado nutricional e a autopercepção do estado nutricional com a experimentação de drogas lícitas em adolescentes.

Métodos: Estudo transversal no qual 210 adolescentes responderam a um questionário sobre experimentação de álcool e tabaco e sobre suas autopercepções nutricionais. Foi analisada a concordância entre autopercepção nutricional do adolescente e seu estado nutricional, bem como associações entre o estado nutricional, a autopercepção nutricional, sexo, idade e presença de tabagistas em casa com a experimentação de álcool e tabaco. As variáveis foram analisadas separadamente em análise bivariada e, a seguir, análise múltipla determinou fatores associados à experimentação.

Resultados: Participaram do estudo 210 adolescentes com mediana de idade de 148 meses, 56,6% do sexo feminino. Do total da amostra, 6,6% já experimentaram cigarro e 20% já experimentaram álcool; 32,3% tinham Z-escore $IMC \geq 1$ e 12,85% tinham Z-escore $IMC \geq 2$ e 50,7% acertaram suas classificações nutricionais. Após análise multivariada, apenas a autoimagem sobre o peso influenciou estatisticamente na experimentação de fumo. Pacientes que se identificavam com peso muito alto apresentaram maior chance de experimentação de fumo (*odds ratio* (OR) 13,57; intervalo de confiança (95% IC) 2,05–89,8; $p=0,007$); em relação ao uso de álcool, adolescentes que se identificavam com peso alto apresentaram chance de uso de álcool 2,40 vezes maior do que crianças que se viam com peso normal (IC 95% 1,08–5,32; $p=0,031$).

Conclusões: Adolescentes com autopercepção de excesso de peso podem constituir um grupo de risco para a experimentação de álcool e tabaco.

© 2015 Sociedade de Pediatria de São Paulo. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

Introduction

There has been a significant increase in the number of overweight children and adolescents in the last two decades worldwide.^{1–3} In Brazil, when comparing data between 1975 and 2010 in the age group between 10 and 19 years, excess weight increased from 3.7% to 21.7% in boys and 7.6% to 19% in girls. According to the Brazilian Institute of Geography and Statistics (Instituto Brasileiro de Geografia e Estatística – IBGE) data, 34.8% of boys and 32% of girls were overweight and 16.6% of boys and 11.8% of girls were obese in 2010.⁴

Obesity has become the most prevalent chronic non-communicable disease in children and adolescents, with major future effects on health.⁵ Excess weight in this population is associated with the occurrence of type 2 diabetes mellitus and cardiovascular disease, increased prevalence of obesity and increased morbidity and mortality in adulthood, as well as increased risk of psychosocial problems and psychological abuse at school.^{6–10}

Adolescence should be seen as a phase of growth and development, characterized by transformation. It is a stage of human life characterized by biological, psycho-emotional and socio-cultural changes that constitute an important period to implement new practices, behaviors and gain autonomy.¹¹ The discrepancy between these changes may contribute to the adolescents' dissatisfaction with their bodies and lead to risky behaviors and vulnerability. Previous studies have suggested that adolescents generally perceive

their nutritional status inappropriately^{12,13}; most often, this population tends to underestimate their own nutritional status, especially those who are overweight. However, in recent years, there has been a change in the perception of adolescent nutritional status, more prone to overestimation and the habit of dieting, especially girls, reflecting the consequences of the ideal thin body glorified by society.^{12,13}

The normal adolescence syndrome consists of important features such as the search for one's identity, group trends, mood swings, the evolution of sexuality and separation from parents, among others.¹¹ These characteristics may interfere with the perception that adolescents have about their own nutritional status and thus, with their degree of satisfaction with their own body image.¹⁴

The adolescent dissatisfied with his/her body image can resort to risky behaviors, such as very restrictive diets, practice of unsafe sex and use of licit and illicit drugs in order to change one's body or to be more accepted by their peers^{15,16} Every year, millions of adolescents around the world lose their lives, most due to automobile accidents, suicide and homicide,¹⁷ often associated with the consumption of alcohol or other drugs.

It is known that early use of licit drugs constitutes an immediate hazard to the adolescent's health.¹⁸ Some studies have suggested that when experimenting with illicit drugs in adolescence is associated with other risk factors, such as family history of substance abuse, practice of other risk behaviors, substance abuse among peers, lack of interest in school activities, among others, it can be the first step

toward the abuse of these drugs or experimenting with illicit drugs in the future.^{18,19}

In Brazil, there have been few studies to verify the association between nutritional status and the adolescent's self-perception of his/her nutritional status and experimentation of licit drugs, and few studies have associated the adolescent's perception of the nutritional status with nutritional diagnostic criteria. Our goal is to verify the association between excess weight or self-perception of excess weight and experimentation with alcohol and tobacco in adolescence.

Method

This is a cross-sectional study carried out from February to March 2010 in the emergency department of the University Hospital at the University of Sao Paulo – USP (HU – USP).

The nutritional status of 210 adolescents aged 10–14 years, whose legal guardian had sought hospital care for the adolescent, was evaluated. A convenience sample was used, with the inclusion of one adolescent per hour, from 1 PM to 7 PM, from Monday to Friday, during the study period.

The emergency department where the study was carried out belongs to a secondary hospital, of low complexity, with rare cases of patients presenting with chronic underlying diseases. The study excluded adolescents who, at the time of the treatment, had diarrhea and/or vomiting complaints, patients with chronic diseases (including asthma), changes in the general status on admission and adolescents that during or shortly after treatment, showed worsening in the general status or were referred for emergency care observation. The most frequent complaints were colds and flu, hoarseness, sore throat, ear pain, conjunctivitis, urinary complaints, menstrual cramps, headache, acute skin changes and mild trauma.

The adolescents were included in the study after the free and informed consent form was signed by a parent or legal guardian, and after the adolescent agreed to participate in the study and answer the questionnaire. All participants had the weight measured using a Toledo electronic scale, model 2096 PP/2, series 10113407, 2007 (Toledo®, São Paulo, Brazil), with a precision of 50g and capacity of 200kg, calibrated by a specialized company according to the criteria of the National Institute of Metrology, Quality and Technology (INMETRO).

Height was measured using a fixed Tonelli aluminum stadiometer, model E 150 A (Tonelli®, Santa Catarina, Brazil), with a precision of 1mm and measurement capacity of 400–2200mm. The measurements were performed in triplicate and the mean value was used to calculate the body mass index (BMI), using the formula: weight in kg/height.²

The adolescents answered a questionnaire that consisted of the following questions: (1) Do you attend school? (2) What grade are you in? (3) Have you ever smoked? (4) Do you smoke regularly? If yes, how many cigarettes do you smoke a day? (5) Do you live with someone who smokes? (6) Have you ever drunk alcohol? (7) Do you drink alcohol regu-

larly? If yes, how often do you drink alcohol? and (8) What do you think of your weight? Five answers were offered for this last question, which included the following alternatives: (1) very low; (2) low; (3) it is OK as it is; (4) high and (5) very high. The answer "I do not know" was made available to all questions, but disregarded in the analysis.

The questionnaire was completed by the adolescents in a private room, so that responses were not influenced by the presence of the parent/guardian. The questionnaires were answered in the presence of the researcher for any clarification of doubts; however, without interfering with the responses.

The World Health Organization (WHO) criteria were used for the nutritional diagnosis. The adolescents' responses about their nutritional status were evaluated together with the WHO diagnostic criteria, using BMI Z-score according to gender. The study found that the adolescent got a correct classification of his/her nutritional status in the following situations: BMI Z-score <−3 and the answer about his/her own weight was "Very Low"; BMI Z-score between −3 and −2 and the answer was "Low"; BMI Z-score between −2 and +1 and the answer was "It's OK as it is"; BMI Z-score between +1 and +2 and the answer was "High"; and BMI-Z score >+2 and the answer was "Very High".

The answers obtained from the questionnaires were first described and categorized according to the adolescent's gender. The responses on the nutritional status were compared with the objective nutritional classification to assess the adolescents' self-perception, initially in the whole sample, and then, according to gender. We also analyzed the association between the adolescent's nutritional status and alcohol or tobacco experimentation, and between the adolescent's nutritional self-classification and experimentation of the same licit drugs.

The quantitative characteristics were described according to the consumption of alcohol and tobacco with the use of summary measures (mean, standard deviation, median, minimum and maximum values) and compared between groups using Student's *t*-tests. The assessed qualitative characteristics were described according to the consumption of alcohol and tobacco with the use of absolute and relative frequencies and significant associations were verified using the chi-square test or Fisher's exact test. The *odds ratio* (OR) of each variable of interest with the use of alcohol and tobacco was estimated with the respective 95% confidence intervals, using simple logistic regression.

Multiple logistic regression models were estimated for consumption of alcohol and tobacco, selecting the variables that showed significance levels <0.20 ($p < 0.20$) in the bivariate tests. The tests were performed with a 5% significance level. This study was approved by the Research Ethics Committee of Hospital Universitário da USP (registration CEP/HU-USP 868/09).

Results

A total of 210 adolescents aged 10–14 years, with a median age of 148 months, were included in the study. Of these,

Table 1 Adolescents' own opinions on their nutritional status and their objective nutritional classification according to body mass index Z-score (Z).

Adolescents' self-classification	Nutritional Classification – n (%)					Total
	Severe thinness (Z<-3)	Thinness (-3<Z<-2)	Normal weight (-2<Z<+1)	Overweight (+1<Z<+2)	Obesity (Z>+2)	
Very skinny	0	2 (18.2)	6 (54.5)	2 (18.2)	1 (9.1)	11
Skinny	0	1 (4.8)	18 (85.7)	1 (4.8)	1 (4.8)	21
It is good as it is	0	2 (1.8)	83 (76.1)	21 (19.3)	3 (2.8)	109
Fat	0	0	23 (46.9)	12 (24.5)	14 (28.6)	49
Very fat	0	0	2 (15.4)	4 (30.8)	7 (53.8)	13
Total	0	5 (2.5)	132 (65)	40 (19.7)	26 (12.8)	203

56.6% were females and all attended school, with 79.2% of adolescent males and 62.3% of females being enrolled between the 5th and 7th grades of Elementary School.

Regarding the experimentation of licit drugs, 6.6% had tried cigarettes, 0.95% smoked regularly (at least one cigarette a day) and 44.7% lived with at least one smoker in the house. As for alcohol, 20% had tried it, and 0.48% drank regularly (one adolescent reported drinking alcohol once a week).

Regarding the adolescents' nutritional classification, 32.3% had BMI Z-score ≥ 1 , of which 50% were boys and 12.85% had BMI Z-score ≥ 2 , of which 51.8% of them were

boys. Of the 210 adolescents who participated in the study, seven were unable to classify their nutritional status, choosing the answer "I do not know"; these responses were not considered in the analysis, thus totaling 203 adolescents. Of these, 50.7% correctly classified their nutritional status. Of those that incorrectly classified their nutritional status, 69% underestimated, whereas 31% overestimated their own nutritional status.

Among the adolescents that classified themselves as having high weight, 24.5% actually had BMI Z-score between +1 and +2. Of those that classified themselves as having very high weight, 53.8% were obese (BMI Z-score $>+2$). Among

Table 2 Description of the assessed characteristics according to smoking experimentation.

Variable	Have you ever smoked?		Total (n=210)	OR	95%CI		p-value
	No (n=196)	Yes (n=14)			Inferior	Superior	
Sex, n (%)							0.602
Female	112 (57.1)	7 (50)	119 (56.7)	1.00			
Male	84 (42.9)	7 (50)	91 (43.3)	1.33	0.45	3.95	
Smoker at home, n (%)							0.656
No	110 (56.1)	7 (50)	117 (55.7)	1.00			
Yes	86 (43.9)	7 (50)	93 (44.3)	1.28	0.43	3.79	
Opinion about their own weight, n (%)			Total (n=203)				0.004
Very skinny	10 (5.3)	1 (7.1)	11 (5.4)	3.53	0.34	37.21	
Skinny	21 (11.1)	0 (0)	21 (10.3)	-			
Normal	106 (56.1)	3 (21.4)	109 (53.7)	1.00			
Fat	43 (22.8)	6 (42.9)	49 (24.1)	4.93	1.18	20.61	
Very fat	9 (4.8)	4 (28.6)	13 (6.4)	15.70	3.03	81.31	
Age							0.841
Mean (SD)	148.2 (17.4)	0.4 (1.3)	1.2 (1.6)	0.4 (1.3)	1.00		
Median (min; max)	148 (116; 180)	146 (123; 180)	148 (116; 180)	1.00	0.97	1.04	
BMI Z score							0.021
Mean (SD)	0.4 (1.3)	1.2 (1.6)	0.4 (1.3)	1.00			
Median (min; max)	0.4 (-2.8; 4.4)	1.3 (-2.5; 3.4)	0.4 (-2.8; 4.4)	1.64	1.07	2.52	

OR, odds ratio; CI, confidence interval; SD, standard deviation.

the adolescents with excess weight (Z-score >+1) and who incorrectly classified their nutritional status, 65.1% underestimated it, as shown in Table 1.

Among girls, 49.2% correctly classified their nutritional status; of those that incorrectly classified it, 60% underestimated and 40% overestimated their nutritional status. Among boys, 52.9% correctly classified their nutritional status; among those that incorrectly classified their nutritional status, 82.5% underestimated and 17.5% overestimated it. Of the girls with excess weight, 27.2% correctly classified their nutritional status, and of those that incorrectly classified it, 65.15% underestimated it. Of all the boys with excess weight, 30.3% correctly classified their nutritional status, and of those that incorrectly classified it, 69.7% underestimated it.

As for the bivariate analysis for tobacco use, Table 2 shows that, alone, the adolescents' self-perception of their own weight and BMI Z-score influenced tobacco experimentation ($p=0.004$ and $p=0.021$, respectively). In the multivariate regression, as shown in Table 3, it is observed that only the self-image on weight was associated with smoking experimentation, and children who saw themselves as having very high weight were more likely to experiment tobacco use (OR: 13.57; 95% CI: 2.05–89.8, $p=0.007$).

Regarding the experimentation of alcohol, it was verified that, according to Table 4, in the bivariate analysis, only the self-perception of their own weight influenced the use of alcohol by adolescents ($p=0.042$). The multivariate analysis (Table 5) shows that only the self-perception of their own

Table 3 Multiple logistic regression model for the response variable smoking experimentation by adolescents.

Variable	OR	95%CI		p-value
		Inferior	Superior	
Perception of their own weight				
Very skinny	3.69	0.35	39.54	0.280
Skinny	^a			0.998
Normal	1.00			
Fat	4.53	0.98	20.94	0.053
Very fat	13.57	2.05	89.80	0.007
BMI Z score	1.09	0.63	1.60	0.763

^a Not possible to estimate.

OR, *odds ratio*; CI, confidence interval.

weight together influenced the experimentation of alcohol by the adolescents and those who classified themselves as having high weight had a 2.40-fold higher chance of alcohol consumption than those who classified themselves as having normal weight.

Discussion

Till date, there are no known data on the association between body weight perception by adolescents and the risk of experimenting with licit drugs. Our study showed that the

Table 4 Description of the assessed characteristics evaluated according to alcohol experimentation.

Variable	Have you ever drunk alcohol?		Total (n=210)	OR	95%CI		p-value
	No (n=169)	Yes (n=41)			Inferior	Superior	
Gender, n (%)							0.256
Female	99 (58.6)	20 (48.8)	119 (56.7)	1.00			
Male	70 (41.4)	21 (51.2)	91 (43.3)	1.49	0.75	2.95	
Smoker at home, n (%)							0.145
No	90 (53.3)	27 (65.9)	117 (55.7)	1.00			
Yes	79 (46.7)	14 (34.1)	93 (44.3)	0.59	0.29	1.21	
Perception of their own weight, n (%)			Total (n=203)				0.042
Very low	10 (6.1)	1 (2.5)	11 (5.4)	0.47	0.06	3.92	
Low	20 (12.3)	1 (2.5)	21 (10.3)	0.24	0.03	1.87	
Normal	90 (55.2)	19 (47.5)	109 (53.7)	1.00			
High	33 (20.8)	16 (40)	49 (24.1)	2.30	1.06	4.99	
Very high	10 (6.1)	3 (7.5)	13 (6.4)	1.42	0.36	5.66	
Age							0.071
Mean (SD)	147.2 (17.2)	152.6 (18.3)	148.2 (17.5)	1.00	1.00		
Median (min; max)	147 (116; 180)	156 (119; 179)	148 (116; 180)	1.02	1.00	1.04	
BMI Z-score							0.482
Mean (SD)	0.4 (1.3)	0.3 (1.2)	0.4 (1.3)	1.00			
Median (min; max)	0.5 (-2.8; 4.4)	0.3 (-2.5; 2.5)	0.4 (-2.8; 4.4)	0.91	0.70	1.19	

OR, *odds ratio*; CI, confidence interval; SD, standard deviation.

Table 5 Results of the multiple logistic regression model for the response variable alcohol experimentation by adolescents.

Variable	OR	95%CI		p-value
		Inferior	Superior	
Age	1.01	0.99	1.03	0.261
Perception of their own weight, n (%)				0.102
Very skinny	0.55	0.07	4.65	0.584
Skinny	0.28	0.04	2.25	0.231
Normal	1.00			
Fat	2.40	1.08	5.32	0.031
Very fat	1.33	0.33	5.43	0.692
Smoker at home	0.58	0.27	1.23	0.152

OR, odds ratio; CI, confidence interval.

self-assessment of weight in cases of overweight and obesity may be associated with increased chance of consumption of these drugs during adolescence.

The high prevalence of overweight (32.3%) and obesity (12.9%) found in this study reinforces the increasing global trend of these nutritional disorders in children and adolescents.¹⁻⁴ In our study, the prevalence of excess weight was higher in boys, and this is a trend that is observed in the younger age groups.⁴ Excess weight in adolescents is, by itself, a major cardiovascular risk factor and, when associated with alcohol use and smoking, can lead to important and early effects on adult health.^{5,20}

The overall agreement between the adolescent's self-perception and their nutritional classification given by BMI Z-score was 50.7%. This value differs from previous studies in Brazil, which suggest that 65–70% of adolescences correctly evaluate their nutritional status.^{21,22} This difference is due to two main reasons. First, our sample encompasses an age group that includes early adolescence (10–14 years), while another Brazilian study analyzed the late adolescence age range (14–19 years), with the latter being more stable from a physiological point of view and better able to achieve correct self-perception of one's body.²² Second, the population evaluated in this study attended public schools, while the population from the other study carried out in Salvador and that evaluated the same age group attended private schools, which may suggest that socioeconomic differences interfere with the adolescent's nutritional status self-perception; however, additional data on such differences are necessary to support this conclusion.²²

Among the adolescents that incorrectly assessed their nutritional status, most underestimated the weight. This was observed in both genders, including adolescents with excess weight; however, boys underestimated their own nutritional status more often than girls (82.5% vs. 60%). This may reflect a cultural characteristic of praising "strong" boys in all age groups, especially among adolescents, who often seek a more muscular body.²³ Previous studies have shown that since infancy, boys have their

weights more often underestimated by their mothers than girls.²⁴ As our study only evaluated early adolescence, many answers given by the adolescents may reflect maternal opinions.

In the present study, many adolescents lived with at least one smoker at home; however, this factor was not associated with experimentation of licit drugs, a finding that differs from findings of previous studies suggesting that parental example increases the risk of experimentation of these drugs by adolescents.¹⁸ In addition, in our study, the frequency of smoking experimentation was low (6.6%) compared with previous national data.²⁵ A national survey carried out in 2004 in 12 Brazilian cities, with adolescents aged 13–15 years, showed that cigarette experimentation frequency is high, reaching 50% in Northeastern states.²⁵ The results obtained in this study may reflect the global awareness effort about the dangers of smoking to health.²⁶

Approximately 20% of the adolescents analyzed here had already tried alcohol, a lower frequency than that found in previous population studies, in which the percentage of alcohol experimentation among adolescents in São Paulo was higher than 70%.²⁷ The explanation for this finding is the median age of our sample (12 years), with the most common age for first alcohol experimentation among adolescents being close to 12 years (20%) and increasing between 13 and 14 years (30–40%).²⁸

Previous studies have shown similar data regarding the positive association between excess weight and tobacco experimentation.^{29,30} In these studies, it is suggested that excess food consumption can be conceptualized as an addictive behavior. Much of the evidence comes from neurobiological similarities between drug and food consumption. Moreover, an inverse association has been observed between alcohol consumption and body mass index, an association that can be influenced by psychosocial factors such as family history of alcoholism.²⁸

A review study on excess weight and risk behavior in adolescence showed that the former is not necessarily a risk factor for the use of licit or illicit drugs; however, psychosocial factors involving overweight and obesity may increase adolescents' vulnerability to this type of behavior.²⁹ The present study showed a strong association between self-classification of having high or very high weight and the previous experimentation with tobacco and alcohol, regardless of the actual nutritional status of adolescents. A factor that may contribute to this association is the possible feeling of exclusion of adolescents who consider themselves obese in relation to their peers, as they are often the target of aggression and psychological abuse at school¹⁰ and, consequently, may resort to risky behaviors, such as the use of licit drugs in order to feel accepted by their groups.^{22,23,29}

The present study has limitations that should be mentioned. The first is the selection of a convenience sample, which does not allow the extrapolation of the results to the general population. The second limitation involves the analysis of the nutritional status perception. As this is a subjective variable, the use of the questionnaire with closed alternatives may limit the adolescent's responses. Tools such as body silhouette scales and questionnaires to assess the self-perceived body image could improve the adolescents' response specificity in relation to what they think about

their own body. The third limitation is the lack of outpatient follow-up.

The emergency care approach may have interfered with the adolescents' responses regarding the experimentation of licit drugs, due to a possible lack of trust regarding the confidentiality of answers or the stress caused by the event that led the adolescent to the emergency department. It is worth mentioning that there may be an overlap between the environmental and psychological factors that lead an adolescent to obesity and also to the use of licit and illicit drugs, which may have contributed further to the findings of the study. Finally, the questionnaire applied to the adolescents is designed only for the present study and has not been validated and its variability between examiners has not been assessed. A more reliable questionnaire tool can yield more reliable information on the variables associated with experimentation of licit drugs in adolescence.

In summary, our findings on the association between excess weight, self-perception of nutritional status and experimentation of licit drugs deserve special attention regarding the promotion of adolescent health, not only because the use of licit drugs is an immediate hazard to adolescents' health, but because it can also be related to the abuse of these drugs in the future and the experimentation of illicit drugs.^{19,27,30}

Funding

This study did not receive funding.

Conflicts of interest

The authors declare no conflicts of interest.

References

- World Health Organization. Obesity: preventing and managing the global epidemic [Report of a WHO Consultation on Obesity, 1997]. Geneva: WHO; 1998.
- World Health Organization. The World Health report 2002: reducing risks, promoting healthy life. Geneva: WHO; 2002.
- Wang Y, Monteiro C, Popkin BM. Trends of obesity and underweight in older children and adolescents in the United States, Brazil, China, and Russia. *Am J Clin Nutr.* 2002;75:971-7.
- Brasil - Instituto Brasileiro de Geografia e Pesquisa; Ministério do Planejamento, Orçamento e Gestão. Pesquisa do Orçamento Familiar - 2008-2009. Available at: www.ibge.gov.br [cited 10.01.14].
- Freedman DS, Khan LK, Dietz WH, Srinivasan SR, Berenson GS. Relationship of childhood obesity to coronary heart disease risk factors in adulthood: the Bogalusa Heart Study. *Pediatrics.* 2001;108:712-8.
- Fagot-Campagna A, Saaddine JB, Flegal KM, Beckles GL. Diabetes, impaired fasting glucose, and elevated HbA_{1c} in U.S. adolescents: the Third National Health and Nutrition Examination Survey. *Diabetes Care.* 2001;24:834-7.
- Sinha R, Fisch G, Teague B, Tamborlane WV, Banyas B, Allen K, et al. Prevalence of impaired glucose tolerance among children and adolescents with marked obesity. *N Engl J Med.* 2002;346:802-10.
- Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH. Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med.* 1992;327:1350-5.
- Pearce MJ, Boergers J, Prinstein MJ. Adolescent obesity, overt and relational peer victimization, and romantic relationships. *Obes Res.* 2002;10:386-93.
- Janssen I, Craig WM, Boyce WF, Pickett W. Associations between overweight and obesity with bullying behaviors in school-aged children. *Pediatrics.* 2004;113:1187-94.
- Saito MI. Adolescência, cultura, vulnerabilidade e risco [editorial], 22. São Paulo: *Pediatria*; 2000. p. 217-9.
- Quick V, Nansel TR, Liu D, Lipsky LM, Due P, Iannotti TJ. Body size perception and weight control in youth: 9-year international trends from 24 countries. *Int J Obes (Lond).* 2014;38:988-94.
- Jasik CB. Body image and health: eating disorders and obesity. *Prim Care.* 2014;41:519-37.
- Wang Y, Liang H, Chen X. Measured body mass index, body weight perception, dissatisfaction and control practices in urban, low-income African American adolescents. *BMC Public Health.* 2009;9:183.
- Johnson F, Wardle J. Dietary restraint, body dissatisfaction, and psychological distress: a prospective analysis. *J Abnorm Psychol.* 2005;114:119-25.
- Tedesco S. Adolescência e drogas: algumas indicações éticas e políticas. In: Mello A, Castro AL, Geiger M, editors. *Conversando sobre adolescência e contemporaneidade.* Porto Alegre: Libretos; 2004. p. 106-11.
- World Health Organization. Adolescents: health risks and solutions. Available at: <http://www.who.int/mediacentre/factsheets/fs345/en> [accessed 23.11.14].
- Latimer W, Zur J. Epidemiologic trends of adolescent use of alcohol, tobacco, and other drugs. *Child Adolesc Psychiatr Clin N Am.* 2010;19:451-64.
- Everett SA, Giovino GA, Warren CW, Crossett L, Kann L. Other substance use among high school students who use tobacco. *J Adolesc Health.* 1998;23:289-96.
- Li S, Yun M, Fernandez C, Xu J, Srinivasan SR, Chen W, et al. Cigarette smoking exacerbates the adverse effects of age and metabolic syndrome on subclinical atherosclerosis: the Bogalusa Heart Study. *PLOS ONE.* 2014;9:e96368.
- Boa-Sorte N, Neri LA, Leite ME, Brito SM, Meirelles AR, Luduvise FB, et al. Maternal perceptions and self-perception of the nutritional status of children and adolescents from private schools. *J Pediatr (Rio J).* 2007;83:349-56.
- Branco LM, Hilário MO, Cintra IP. Perception and satisfaction with body image in adolescents and correlations with nutrition status. *Rev Psiq Clin.* 2006;33:292-6.
- Iriart JA, Andrade TM. Body-building, steroid use, and risk perception among young body-builders from a low-income neighborhood in the city of Salvador, Bahia State, Brasil. *Cad Saude Publica.* 2002;18:1379-87.
- Francescato C, Santos NS, Coutinho VF, Costa RF. Mothers' perceptions about the nutritional status of their overweight children: a systematic review. *J Pediatr (Rio J).* 2014;90:332-43.
- Instituto Nacional de Câncer [página na Internet]. Vigéscola: vigilância de tabagismo em escolares: dados e fatos de 12 capitais brasileiras. Available at: http://bvsms.saude.gov.br/bvs/publicacoes/vigescola_vol1.pdf [cited 11.08.14].
- Reichert J, Araújo AJ, Gonçalves CM, Godoy I, Chatkin JM, Sales MP, et al. Smoking cessation guidelines - 2008. *J Bras Psiquiatr.* 2008;34:845-80.
- Malta DC, Mascarenhas MD, Porto DL, Duarte EA, Sardinha LM, Barreto SM, et al. Prevalence of alcohol and drug consumption among adolescents: data analysis of the National Survey of School Health. *Rev Bras Epidemiol.* 2011;14:136-46.

28. Gearhardt AN, Corbin WR. Body mass index and alcohol consumption: family history of alcoholism as a moderator. *Psychol Addict Behav.* 2009;23:216–25.
29. Rosa MF, Gonçalves S, Antunes H. Risk-taking behaviors and overweight in adolescence. A review of the literature. *Acta Pediatr Port.* 2012;43:128–34.
30. Everett SA, Malarcher AM, Sharp DJ, Husten CG, Giovino GA. Relationship between cigarette, smokeless tobacco, and cigar use, and other health risk behaviors among U.S. high school students. *J Sch Health.* 2000;70:234–40.