



Crowding correction through extraction of a lower incisor

Corrección del apiñamiento, con extracción del incisivo inferior

Mauricio Servin Muñoz,* Beatriz Gurrola Martínez,§ Adán Casasa Araujo||

ABSTRACT

Tooth-bone discrepancy, or crowding in the lower arch, is traditionally corrected by removing first or second premolars; the extraction of a lower incisor is an uncommon approach but presents another treatment alternative. Treatment goal: to present the case report of a patient who was given this alternative treatment. The patient had 15 years 4 months of age. The intraoral analysis revealed a normodivergent skeletal class I with bilateral molar class I, bilateral canine class I, decreased overbite, and increased overjet. She had lower moderate and mild upper crowding as well as a habit of digital suction. Treatment time was one year and four months.

Key words: Extraction, lower incisor, class 1, normodivergent.

Palabras clave: Extracción, incisivo inferior, clase I, normodivergente.

RESUMEN

La discrepancia de longitud, o el apiñamiento en la arcada inferior, tradicionalmente se corrige por medio de la extracción de primeros o segundos premolares; la extracción de un incisivo inferior es un enfoque poco común pero es otra alternativa sobre todo en el arco mandibular. **Objetivo:** Presentar el tratamiento de una paciente de ortodoncia con estas características, a la cual se le da este tratamiento, de 15 años cuatro meses de edad. Al análisis intraoral, es clase I esquelética, normodivergente, con clase I molar bilateral, clase I canina bilateral, con el *overbite* disminuido, y *overjet* aumentado, tiene apiñamiento moderado inferior, y leve superior, presenta hábito de succión digital. El tiempo que duró el tratamiento fue de un año cuatro meses.

INTRODUCTION

Jackson describes a case in which two lower incisors were extracted at different times to solve mandibular crowding.¹ Since then, a series of clinical cases have been treated with this option.^{2,3} Authors such as: Canut, Bahreman, have listed specific criteria for the removal of a mandibular incisor such as: permanent dentition, minimum growth potential, class I molar relationship, a harmonious soft tissue profile, minimal to moderate overbite, little or no crowding in the upper arch, a Bolton discrepancy.^{2,4} A good diagnosis is highly recommended with this kind of treatment.^{5,6} On the other hand, the removal of a mandibular incisor has several advantages over premolar extractions. First, treatment time may be reduced especially if the crowding is limited only to the anterior segment.⁵ Second, it offers a stable result in the anterior region because the expansion is not necessary and canine width is not altered.³ Finally, since it requires little retraction in comparison with the therapy of premolars removal, the anteroposterior position of the lower incisors is not changed which maintains a harmonious profile.⁴ This treatment also offers us a series of disadvantages: if there is a Bolton discrepancy, space closure will result in a

greater overjet, in addition to altering the interproximal papillae, which can cause black triangles.⁷

CASE REPORT

During clinical evaluation, the patient reported as reason for consultation «I have crooked teeth». In the initial radiographic studies, the lateral head film showed normodivergency and a mild proclination of the upper and lower incisors (*Figure 1*). In the panoramic radiograph no pathological data were

* Second year Resident of the Masters Program in Orthodontics and Maxillofacial Orthopedics.

§ Professor at the CESO and full-time professor in the Odontology Faculty of the Faculty of Higher Studies Zaragoza UNAM.

|| Director CESO.

Center for Higher Studies in Orthodontics (CESO).

© 2017 Universidad Nacional Autónoma de México, [Facultad de Odontología]. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

This article can be read in its full version in the following page:
<http://www.medigraphic.com/ortodoncia>

noted; the patient had 28 permanent teeth, formation of the lower third molars and agenesis of the upper (Figure 2).

Initial records

In the facial analysis of the photographs of the patient, the profile was assessed, and it was found to be adequate. The dental midline was diverted 0.5 mm to the right in relation to the facial; facial fifths and thirds were proportional. All these were meaningful data in decision-making for the treatment plan (Figure 3), facial midline (Figure 4), and aesthetic line of Ricketts, E line.

When performing the intraoral analysis the objectives posed to cover were observed: elimination of upper mild and lower moderate anterior crowding, to correct the lack of coincidence between the upper and lower dental midlines. The patient was in canine and molar class I, which would be maintained. Another



Figure 1. Initial lateral head film.



Figure 2. Initial panoramic radiograph.



Figure 3. Facial midline.




Figure 4. E line.

aspect to highlight was the 6 mm overjet caused by the habit and the upper and lower triangular arch form intraoral front (Figure 5), in the intraoral photographs, right, and left side (Figures 6 and 7).

Treatment plan

To eradicate the habit of digital suction, a palatal grid was placed for two months. Afterwards, the appliance was withdrawn and Roth prescription brackets were placed giving instructions for the removal of the lower right lateral incisor.

Alignment and leveling was begun on both the upper and lower arches, with stripping sessions in the upper arch and closure of the extraction space in the lower arch. After the alignment and leveling (Figure 8),

lower occlusal photograph indicating the tooth to be extracted, in the lower occlusal image of progress after space closure (*Figure 9*).

Final records

The objectives were met: the E line was improved, bilateral molar and canine class I were maintained, a better arch form on both the upper and lower arch was achieved and therefore, a normal overjet and overbite as well. Good intercuspation was obtained, as it may



Figure 5. Intraoral frontal photograph.



Figure 6. Initial right lateral photograph.



Figure 7. Initial left side.



Figure 8. Initial lower occlusal photograph.



Figure 9. Treatment progress photograph.

be observed in the intraoral photographs: final front view (*Figures 10 to 12*), in the right and left side.

In the final lower occlusal photograph, the fixed retainer from canine to canine may be observed (*Figure 13*).

RESULTS

As it may be observed in this case the set objectives were achieved. The patient ended with excellent aesthetics and a harmonious smile. Total treatment time was one year and four months. A comparison between the initial and final smile is shown in *figure 14*. Although the extraction of a lower incisor is not a very common treatment, it may provide an excellent result if a good diagnosis is done.

DISCUSSION

Canut² points out the aspects to be considered for this type of treatment, which were taken as



Figure 10. Final intraoral front photograph.



Figure 11. Final right lateral photograph.



Figure 12. Final left lateral photograph.

reference for the procedure: X-rays, models and intra and extra oral photographs, were obtained in order to follow those guidelines. Uribe⁸ suggested that the extraction of a mandibular incisor is the most appropriate treatment in patients who do not require modifications in the anteroposterior buccal segments, as was our patient, and for whom this option was ideal since one of the goals was to maintain that anteroposterior relationship. On the other hand Faerovig⁷ points out that a great disadvantage with this treatment lies in the



Figure 13. Final lower occlusal photograph.



Figure 14. Initial and final smile comparison.

interproximal papillae, which may be flattened and may lead to the development of «black triangles». This was the reason why this case was handled with extreme care, in order to obtain optimal aesthetics.

CONCLUSIONS

The extraction of a mandibular incisor is an effective treatment in patients who meet the requirements established by the literature and who are aware of it in order to obtain excellent results, since a wrong diagnosis can lead to unwanted side effects and will directly affect the aesthetics of the patient.

REFERENCES

1. Jackson VH. *Orthodontia and orthopedic of the face*. Philadelphia: J.B. Lippincott Co.; 1904.
2. Canut JA. Mandibular incisor extraction: indications and long-term evaluation. *Eur J Orthod*. 1996; 18 (5): 485-489.
3. Riedel RA, Little RM, Bui TD. Mandibular incisor extraction--postretention evaluation of stability and relapse. *Angle Orthod*. 1992; 62 (2): 103-116.
4. Bahreman AA. Lower incisor extraction in orthodontic treatment. *Am J Orthod*. 1977; 72 (5): 560-567.
5. Kokich VG, Shapiro PA. Lower incisor extraction in orthodontic treatment. Four clinical reports. *Angle Orthod*. 1984; 54 (2): 139-153.
6. Valinoti JR. Mandibular incisor extraction therapy. *Am J Orthod Dentofacial Orthop*. 1994; 105 (2): 107-116.
7. Faerovig E, Zachrisson BU. Effects of mandibular incisor extraction on anterior occlusion in adults with Class III malocclusion and reduced overbite. *Am J Orthod Dentofacial Orthop*. 1999; 115 (2): 113-124.
8. Uribe F, Nanda R. Considerations in mandibular incisor extraction cases. *J Clin Orthod*. 2009; 43 (1): 45-51.

Mailing address:

Dra. Beatriz Gurrola Martínez

E-mail: beatgurrola@gmail.com