

ever, both types of *Coxsackieviruses* were predominant in patients between 6 and 24 months old (31.3 per cent).

The predominance of a given EV in the different age groups varies depending on the circulation of such virus. Thus, in the period that goes from 2010 to 2013, in our country, most neonatal infections were due to *Coxsackievirus* type A (83 per cent),⁴ but today it seems that the predominant viruses are different types of *Echoviruses*.

Also, the EV-D68 is the most highly detected virus in patients under 6 months old (54.5 per cent), although very homogeneous distributions have been confirmed in the remaining ages. However, something consistent with other studies⁷ is the fact that this virus has been the predominant virus in patients >5 years old (63.6 per cent). Unlike other studies,⁸ when it comes to EV-A71, 62.5 per cent of the patients were <1 year old.

We have been able to confirm that EVs preferably affect patients under 15 years old because in this same period we confirmed three (3) cases in adult patients only. This is why we should take their participation into account not only in the ARIs of these patients, but in most fever syndromes without a focus. The distribution of the different types of viruses shows a certain preference for certain age segments, specially *Echoviruses*.

References

- Pallansch MA, Oberste MS, Whitton JL. Enteroviruses: Polioviruses, Coxsackieviruses, Echoviruses and newer Enteroviruses. In: Knipe DM, Howley PM, editors. *Fields virology*, 6th ed. Philadelphia: Wolter Kluwer; 2013. p. 490–530.
- Dunn JJ. Enteroviruses and parvoviruses. *Microbiol Spectr*. 2016; <http://dx.doi.org/10.1128/microbiolspec.DMIH2-0006-2015>
- de Crom SC, Rossen JW, van Furth AM, Obihara CC. Enterovirus and parvovirus infection in children: a brief overview. *Eur J Pediatr*. 2016;175:1023–9.
- Cabrero M, Díaz-Cerio M, Muñoz-Almagro C, Rabella N, Tarragó D, Romero MP, et al. Molecular epidemiology of enterovirus and parvovirus infections according to patient age over a 4-year period in Spain. *J Med Virol*. 2016; [doi:10.1002/jmv.24658](https://doi.org/10.1002/jmv.24658).
- Molet L, Saloum K, Marque-Juillet S, Garbarg-Chenon A, Henquell C, Schuffenecker I, et al. Enterovirus infections in hospitals of Ile de France region over 2013. *J Clin Virol*. 2016;74:37–42.
- Reina J, Cabrero M, Aliaga F. Infecciones respiratorias agudas comunitarias causadas por enterovirus en la población pediátrica. *Enferm Infecc Microbiol Clin*. 2016; <http://dx.doi.org/10.1016/j.eimc.2016.05.007>
- Xiang Z, Wang J. Enterovirus D68 and human respiratory infections. *Semin Respir Crit Care Med*. 2016;37:578–85.
- Chang PC, Chen SC, Chen KT. The current status of the disease caused by enterovirus 71 infections: epidemiology, pathogenesis, molecular epidemiology and vaccine development. *Int J Environ Res Public Health*. 2016;13; [doi:10.3390/ijerph13090890](https://doi.org/10.3390/ijerph13090890).

Jordi Reina^{a,*}, María Cabrero^b

^a Unidad de Virología, Servicio de Microbiología, Hospital Universitario Son Espases, Palma de Mallorca, Balearic Islands, Spain

^b Centro de Referencia de Enterovirus, Centro Nacional de Microbiología, Majadahonda, Madrid, Spain

* Corresponding author.

E-mail address: jorge.reina@ssib.es (J. Reina).

2529-993X/

© 2017 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.

Infective endocarditis due to *Lactobacillus rhamnosus*: Risks of probiotic consumption in a patient with structural heart disease[☆]



Endocarditis infecciosa por *Lactobacillus rhamnosus*: riesgos del consumo de probióticos en un paciente con cardiopatía estructural

Lactobacillus includes one heterogeneous group of gram-positive, microaerophilic, non-sporulating, catalase-negative bacilli.¹ They are part of the common bacterial flora within the oral cavity, the GI tract, and the genitourinary apparatus.² Its isolation in clinical samples usually goes unnoticed due to its special nutritional requirements; long periods of incubation; and inadequate identifications through conventional systems.³ In most cases, they are usually regarded as contaminants; however, their participation in serious infections such as bacteremia, and endocarditis, especially in hosts with certain comorbidities has often been reported.^{3,4}

We hereby present the case of a twenty-eight (28) year old male who was referred to our hospital for the surgical assessment of an aortic valve endocarditis. His personal history showed the presence of one bicuspid aortic valve. He presented to his hospital emergency room a 10-day history of fever and dyspnea. Upon his arrival, hemocultures were collected and one transthoracic echocardiogram was performed that showed the thickening of his bicuspid aortic valve, and a 5 × 7 mm image of mobile nodules confirmed the presence

of vegetation and severe aortic failure. Empirical antibiotic therapy with ampicillin, cloxacillin, and ceftriaxone was started. Since the patient's heart failure was getting worse and his fever persisted even after 72 h of antibiotic treatment, the patient was referred to our center. After hospital admission, three (3) new sets of hemocultures were collected and the patient's valve was replaced by one biological device. All hemocultures obtained in our center tested positive 48–72 h after incubation even though the patient was receiving antibiotic treatment.

The Gram staining revealed the presence of one gram-positive, non-sporulating bacillus (Fig. 1) that was isolated in both blood agar and chocolate agar after 48 h of incubation under aerobic and anaerobic conditions. Alpha-hemolytic and catalase-negative colonies were seen in pure cultures. Using the MALDI-TOF MS analysis (Bruker Daltonics, Bremen, Germany) the *Lactobacillus rhamnosus* bacteria was isolated (score: 2.18). Eventually, the 16S ARNr gene was sequenced showing 99 per cent homology with *L. rhamnosus* (GenBank: CP016823.1). The surgical material of the heart valve showed presence of *L. rhamnosus* bacteria too. Also, the reference hospital confirmed growth of *L. rhamnosus* in the hemocultures.

One antibiogram was conducted using the E-test method, and the minimal inhibitory concentration (MIC) was interpreted based on the recommendations established by EUCAST. The strain was sensitive to penicillin (MIC 0.25 µg/ml); amoxicillin-clavulanic acid (MIC 0.75 µg/ml); imipenem (MIC 0.75 µg/ml); and clindamycin (MIC 0.047 µg/ml). Once the microbiological results were available, a course of treatment with ampicillin (2 g/4 h), and gentamicin (240 mg/24 h) was started and it went on for another six (6) weeks. The patient was interrogated in search for any predisposing factors of lactobacilli endocarditis; the patient denied having any recent dental procedures done, but confirmed the daily use of probiotic products enriched with *Lactobacillus*.

DOI of original article: <http://dx.doi.org/10.1016/j.eimc.2017.02.004>

[☆] Please cite this article as: Recio R, Chaves F, Reyes CA, Orellana MÁ. Endocarditis infecciosa por *Lactobacillus rhamnosus*: riesgos del consumo de probióticos en un paciente con cardiopatía estructural. *Enferm Infecc Microbiol Clin*. 2017;35:609–610.




Fig. 1. Gram staining of one hemoculture (1000×). The presence of gram-positive, non-sporulating bacilli with a tendency to make short chains is consistent with the *Lactobacillus rhamnosus* bacteria.

L. rhamnosus is a rare cause of endocarditis and it has been reported in the medical literature in 18 cases only.⁵ The predisposing factors documented in the development of this entity are dental procedures, periodontitis, immunosuppression, heart valve disease, and the use of probiotics.^{5–7} In our patient, the combination of an underlying heart condition and the use of probiotics were the two (2) factors that most likely contributed to the patient's endocarditis due to *L. rhamnosus*. This microorganism is able to bind itself to different components of the extracellular matrix, such as collagen and fibrinogen, which allows its colonization and survival in valve surfaces.⁸

We should bear in mind that the *L. rhamnosus* bacteria may be erroneously identified using conventional methods.³ Nevertheless, the MALDI-TOF MS system allows us to conduct a quick and reliable

identification of these microorganisms, which helps us minimize the need to use molecular techniques for their characterization.⁹ Most strains of *L. rhamnosus* are sensitive to penicillin and aminoglycosides. However, we may confront clinical manifestations of persistent bacteremia as it happened in our case due to the ability of this microorganism to reduce the pH of its environment through the production of lactic acid, which may complicate the way antibiotics behave.⁶

In sum, although it is a rare condition, *lactobacillus* endocarditis may occur, above all, in patients who have undergone dental procedures or have been treated with probiotics, especially if they show certain comorbidities such as immunosuppression, or structural heart disease. This is why we should not underestimate the importance of isolating it with hemocultures.

References

- Hall V, Copsy S. Propionibacterium, Lactobacillus, Actinomyces, and other non-spore-forming anaerobic gram-positive rods. In: Jorgensen J, Pfaller M, Carroll K, Funke G, Landry M, Richter S, et al., editors. *Manual of clinical microbiology*. 11th ed. Washington, DC: ASM Press; 2015. p. 920–39.
- Slover CM, Dazynger L. Lactobacillus: a review. *Clin Microbiol Newslett.* 2008;30:23–7.
- Salminen MK, Rautelin H, Tynkkynen S, Poussa T, Saxelin M, Valtonen V, et al. Lactobacillus bacteremia, species identification and antimicrobial susceptibility of 85 blood isolates. *Clin Infect Dis.* 2006;42:35–44.
- Salvana EM, Frank M. Lactobacillus endocarditis: case report and review of cases reported since 1992. *J Infect.* 2006;53:5–10.
- Gouriet F, Million M, Henri M, Fournier PE, Raoult D. Lactobacillus rhamnosus bacteremia: an emerging clinical entity. *Eur J Clin Microbiol Infect Dis.* 2012;31:2469–80.
- Cannon P, Lee TA, Bolanos JT, Danzinger LH. Pathogenic relevance of *Lactobacillus*: a retrospective review of over 200 cases. *Eur J Clin Microbiol Infect Dis.* 2005;24:31–40.
- Presterl E, Kneifel W, Mayer HK, Zehetgruber M, Makristathis A, Graninger W. Endocarditis by *Lactobacillus rhamnosus* due to yogurt ingestion. *Scand J Infect Dis.* 2001;33:710–4.
- Vankerckhoven V, Moreillon P, Piu S, Giddey M, Huys G, Vancanneyt M, et al. Infectivity of *Lactobacillus rhamnosus* and *Lactobacillus paracasei* isolates in a rat model of experimental endocarditis. *J Med Microbiol.* 2007;56:1017–24.
- Bizzini A, Jaton K, Romo D, Bille J, Prod'hom G, Greub G. Matrix-assisted laser desorption ionization-time of flight mass spectrometry as an alternative to 16S rRNA gene sequencing for identification of difficult-to-identify bacterial strains. *J Clin Microbiol.* 2011;49:693–6.

Raúl Recio*, Fernando Chaves, Carmen Alhena Reyes, María Ángeles Orellana

Servicio de Microbiología, Hospital Universitario 12 de Octubre, Madrid, Spain

* Corresponding author.

E-mail address: raul.recio@salud.madrid.org (R. Recio).

2529-993X/

© 2017 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.