



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Brief report

Characterization of NDM-1- and CMH-3-producing *Enterobacter cloacae* complex ST932 in a patient transferred from Ukraine to Spain



Inmaculada López-Hernández^{a,b,*}, Aurora García Barrionuevo^c, Paula Díaz de Alba^a, Encarnación Clavijo^c, Alvaro Pascual^{a,b,d}

^a Laboratorio PIRASOA, Unidad Clínica de Enfermedades Infecciosas, Microbiología y Medicina Preventiva, Hospital Universitario Virgen Macarena, Seville, Spain

^b Instituto de Biomedicina de Sevilla (IBiS), Seville, Spain

^c Servicio de Microbiología, Hospital Virgen de la Victoria, Málaga, Spain

^d Departamento de Microbiología, Universidad de Sevilla, Seville, Spain

ARTICLE INFO

Article history:

Received 7 June 2019

Accepted 11 October 2019

Available online 6 December 2019

Keywords:

E. cloacae complex

NDM-1

CMH-3

Spain

ABSTRACT

Introduction: To characterize a carbapenem-resistant *Enterobacter cloacae* complex isolate recovered from a patient from Ukraine.

Methods: The isolate was sent to a regional reference laboratory for molecular characterization by whole genome sequencing. Susceptibility assays, carbapenemase identification, imipenem hydrolysis and clonality were performed.

Results: The isolate showed resistance or reduced susceptibility to all β -lactam agents tested. Genome analysis led to the identification of an NDM-1-producing *E. cloacae* complex strain that was assigned to a new multilocus sequence type, ST932. The *bla*_{NDM-1} enzyme was located in a conjugative IncX3 plasmid of ca. 50 kb. In addition, *bla*_{CMH-3}, a recently described AmpC β -lactamase sequence, which has not previously been reported in Europe, was also detected and its genetic environment was studied.

Conclusion: To our knowledge, this is the first reported case in Europe of an *E. cloacae* complex strain that produces both *bla*_{NDM-1} and *bla*_{CMH-3}.

© 2019 Published by Elsevier España, S.L.U.

Caracterización de *Enterobacter cloacae* complex ST932 productor de NDM-1 y CMH-3 en un paciente trasladado desde Ucrania a España

RESUMEN

Palabras clave:

E. cloacae complex

NDM-1

CMH-3

España

Introducción: Caracterizar una cepa de *Enterobacter cloacae* complex resistente a los carbapenémicos detectado en un paciente procedente de Ucrania.

Métodos: Se envió la cepa a un laboratorio regional de referencia para proceder a su caracterización molecular mediante secuenciación del genoma completo. Se realizaron estudios de susceptibilidad, identificación de carbapenemasas, hidrólisis del imipenem y de clonalidad.

Resultados: El aislado mostró resistencia o sensibilidad reducida a todos los betalactámicos estudiados. El análisis molecular permitió la identificación de una cepa de *E. cloacae* complex productor de NDM-1 que se asignó a un nuevo secuenciotipo, ST932. Se localizó la enzima *bla*_{NDM-1} en plásmido conjugativo IncX3 de 50 kb. Además, se detectó la enzima *bla*_{CMH-3}, una nueva betalactamasa de tipo AmpC descrita recientemente pero no detectada con anterioridad en Europa, llevándose a cabo el estudio de su entorno genético.

Conclusión: Según nuestros conocimientos, esta es la primera descripción una cepa de *E. cloacae* complex productora de *bla*_{NDM-1} junto con *bla*_{CMH-3} en Europa.

© 2019 Publicado por Elsevier España, S.L.U.

* Corresponding author.

E-mail address: inlopezh@us.es (I. López-Hernández).

Introduction

In recent years, *Enterobacter cloacae* complex has emerged as a common nosocomial pathogen with several outbreaks being reported.¹ *E. cloacae* complex naturally exhibits resistance (first and second generation cephalosporins, aminopenicillins) mediated by a chromosomal AmpC. In addition, plasmid-mediated AmpC genes have also been reported in *Enterobacter* spp.¹ The overproduction of AmpC confers resistance to most β -lactam antibiotics, except carbapenems and fourth-generation cephalosporins. Recently, novel plasmid-mediated AmpC-type- β -lactamases (CMH) have been described in Taiwan and India.^{2,3}

Carbapenem resistance in *Enterobacter* spp. has been frequently associated to the production of AmpC enzymes and/or extended-spectrum betalactamases coupled with efflux pump and porin loss.¹ However, worldwide emergence and spread of carbapenemases, such as NDM, in mobile genetic elements are becoming a cause of concern.⁴ In Spain, NDM-1-producing *Enterobacter* spp. isolates have been detected but this is not the predominant carbapenemase in this genus.^{5–8}

Thus, the aim of this study was the molecular characterization of an *E. cloacae* complex clinical isolate carrying *bla*_{NDM-1} and *bla*_{CMH-3}.

Material and methods

In 2016, two carbapenem-resistant *E. cloacae* complex strains were recovered from a patient who had a traffic accident in Ukraine and was transferred, two months later, to the Hospital Virgen de la Victoria (Málaga, Spain). The isolates were recovered from a blood culture (Ec2016198) and a rectal swab (Ec2016212) and were identified as *E. cloacae* complex by MALDI-TOF (Bruker Daltonics).

Antimicrobial susceptibility testing was performed by microdilution (Microscan, Beckman Coulter) and interpreted according to EUCAST (EUCAST, 2018).⁹ β Carba test (BioRad) was used to assess imipenem hydrolysis. The study of inhibitors (phenyl-boronic, dipicolinic acid, cloxacillin) was performed using disc diffusion (Rosco Diagnostica) according to EUCAST recommendations.¹⁰

The presence of carbapenemases (NDM/VIM/KPC/IMP/OXA-48 groups); extended-spectrum beta-lactamases (CTX-M/SHV/TEM); and plasmid-mediated AmpCs (ACC/CMY/FOX/DHA) were evaluated by PCR and DNA sequencing.^{11–13}

Both *E. cloacae* complex isolates were compared by PFGE-*Xba*I.

Ec2016212 was selected for WGS (Illumina MiSeq Inc) and de novo assembly was performed with CLC Genomics Workbench (Qiagen). Antimicrobial resistance and plasmid characterization were determined using ResFinder and PlasmidFinder databases.

For bacterial conjugation *E. coli* J53 was used as recipient. For plasmid electroporation, the plasmid DNA of Ec2016212 was used to electroporate into *E. coli* DH10B. Several colonies of transconjugants and electroporants were tested by microdilution and plasmid transfer by PCR.

The genetic context of the *bla*_{NDM-1} gene was analyzed by SI-PFGE and Southern Blot using probes specific to *bla*_{NDM-1} and IncX3. The Whole Genome project has been deposited at GenBank under the following BioProject No. PRJNA529367 (*E. cloacae* isolate Ec2016212).

Results

Both *E. cloacae* complex isolates were resistant or showed reduced susceptibility to β -lactam agents (Table 1). The β Carba test was positive and a positive synergy meropenem/dipicolinic acid was observed. PCR detection of ESBL, AmpC and carbapenemase genes indicated presence of only *bla*_{NDM}.

PFGE-*Xba*I showed identical patterns in both *E. cloacae* complex isolates.

The assembled data from Ec2016212 provided an allelic profile that was assigned to a new MLST type, ST932. The antimicrobial associated resistance genes identified were *bla*_{NDM-1}; *bla*_{CMH-3}; *aph*(3')-Ia, *aadA16*, *strA* and *strB*; *aac*(6')Ib-cr; *mph*(A), *tet*(A), *sul1* and *dfpA27*. Plasmid Finder identified IncX3 and IncR replicons. S1-PFGE analysis revealed three plasmids, one of ~50-kb and two small plasmids <7 kb. The transformants and transconjugants only contained the 50-kb plasmid, showing signal with NDM and IncX3 probes, suggesting that *bla*_{NDM-1} was in that plasmid.

Conjugation assays confirmed that *bla*_{NDM-1} could be transferred between *E. cloacae* complex and *E. coli* with a transfer frequency of 7×10^{-7} per donor. All transconjugants and electroporants showed similar phenotype regarding β -lactams and the presence of the *bla*_{NDM} gene was confirmed by PCR (Table 1).

*bla*_{CMH-3} could not be transferred between *E. cloacae* complex and *E. coli* and PCR assays did not detect *bla*_{CMH-3} in any transconjugant or electroporant.

*bla*_{NDM} gene was surrounded by a truncated Tn125 (Δ Tn125) structure (Fig. 1). The analysis revealed that upstream of *bla*_{NDM-1} appeared the insertion sequence IS_{Aba}125 disrupted by the IS5 element. Downstream the *bla*_{NDM-1} were located *ble*_{MBL}, *trpF* and *dsbC* genes and further downstream was an IS26 element. The region flanking *bla*_{NDM-1} was like that described for other *bla*_{NDM} genes.¹⁴

The *bla*_{CMH-3} genetic background is shown in Fig. 1. A protein BLAST analysis performed between the enzyme detected in Ec2016212 and the known variants of CMH showed 100% identity to CMH-3, 99% to CMH-1 and 98% to CMH-2.

Discussion

The present study investigated the molecular characterization of an NDM-1 producing *E. cloacae* complex isolate obtained from a patient transferred from Ukraine to Spain. The strain belonged to a new clone ST932. In Spain, NDM is not the predominant carbapenemase in *Enterobacter* spp.^{7,8} A recent work showed that *bla*_{NDM} producers *Enterobacter* spp. were predominant in the Balkans, India and Vietnam.⁴ In accordance with this, our isolate recovered after the transfer from another hospital located in one of these geographic regions suggest that the patient was colonized during his hospitalization in Ukraine. NDM-1 in Enterobacterales is usually located in a plasmid genetic platform and it has been reported that *bla*_{NDM-1} is carried by various types of plasmids such as IncF, IncA/C, IncL/M and IncX3.¹⁵ The analysis of Ec2016212 showed that *bla*_{NDM-1} was carried by an IncX3 plasmid, which was readily transferrable to recipient *E. coli* J53 by conjugation suggesting the possibility of dissemination. Moreover, a recent work has confirmed the in vivo horizontal transfer of *bla*_{NDM} carrying plasmids in Enterobacterales.¹⁵


NDM-1 harbouring strains from different geographic area have shown similar genetic environments.^{14,15} In Ec2016212 the *bla*_{NDM-1} genetic surrounding compared with GenBank sequences revealed that the immediate NDM genetic environment of the IncX3 plasmid was highly similar to others IncX3.¹⁴ The presence of this highly conserved region detected in other NDM harbouring bacteria also supports the horizontal transmission.

The other interesting feature of the strain was the presence of CMH-3, which belongs to CMH enzymes, a new group of AmpC-type β -lactamases. To date, three variants have been described (CMH-1, 2 and 3). In 2015, CMH-1 was reported in an *E. cloacae* isolate from Taiwan, it was successfully conjugated and showed a 99% similarity to a chromosome-borne AmpC in *E. cloacae* ATCC13047.³ Unfortunately, the biochemical characterization was not studied. In 2017 the variant CMH-2 was described as a plasmid-encoded enzyme in

Table 1MICs (mg/L) for the NDM-carrying *E. cloacae* complex clinical isolate Ec2016212, *E. coli* J53, *E. coli* DH10B, transconjugants and transformants.

	Donor strain	Recipient strain	Transconjugants		Recipient strain	Transformants	
	Ec2016212	J53	TC2	TC5	DH10B	TF1	TF3
Amoxicillin-CLA	>64/2	0.5/2	>64/2	>64/2	2/2	>64/2	>64/2
Piperacillin-TZ	128/4	0.125/4	32/4	16/4	0.5/4	32/4	32/4
Cefoxitin	>32	2	>32	>32	4	>32	>32
Ceftriaxone	>32	0.125	>32	>32	≤0.015	>32	>32
Ceftazidime	>32	0.125	>32	>32	0.125	>32	>32
Cefepime	4	0.03	8	4	0.03	4	2
Ertapenem	4	0.015	2	1	0.015	2	2
Meropenem	4	0.03	2	2	0.03	1	1
Imipenem	4	0.25	2	1	0.125	1	0.5
Ciprofloxacin	>4	0.008	0.008	0.004	≤0.002	≤0.002	≤0.002
Amikacin	2	0.5	1	0.5	1	1	2
Gentamicin	0.5	0.25	0.25	0.125	0.25	0.25	0.25
Trimethoprim-sulfamethoxazole	>32	0.06	0.06	0.06	0.125	0.125	0.125
Colistin	0.25	0.5	0.5	0.5	0.25	0.25	0.25

Amoxicillin-CLA: Amoxicillin-clavulanate; Piperacillin-TZ: Piperacillin-tazobactam.

**Fig. 1.** Schematic representation of the DNA sequences surrounding the *bla*_{NDM-1} genes (A) and *bla*_{CMH-3} (B) in *E. cloacae* complex Ec2016212.

clinical isolates of *Klebsiella pneumoniae* collected in India.² CMH-2 showed a 98.6% identity to CMH-1 and provided non inducible resistance to cefotaxime, ceftriaxone and ceftazidime. Transformants and transconjugants were obtained confirming the presence of CMH-2 in a plasmid (IncK replicon).

Regarding CMH-3, the only available information is the nucleotide sequence submitted in 2016 to Genbank (accession number KX192165) from USA without information provided about the genetic location. In Ec2016212 we were unable to determine definitively whether the gene was located on a plasmid or encoded on a chromosome, but the analysis of Illumina contigs that included the enzyme in showed that the enzyme was probably chromosome encoded. No transformants or transconjugants could be obtained in repeated experiments supporting the non-transferability of *bla*_{CMH-3}. To our knowledge this is the first report of the presence of a CMH-type enzyme in a clinical isolate collected in Europe. In Ec2016212, apart from CMH-3, we detected another chromosomal AmpC showing <80% of homology to CMH-3, but 99% to AmpC detected in the chromosome of other *E. cloacae* complex isolates. Further studies focused on how the presence of both AmpC enzymes in terms of regulation and/or expression could influence the susceptibility would be challenging. Plasmid-mediated AmpC originated from chromosome of *E. cloacae* complex have been described.¹ In our isolate, knowing that variants CMH-1 and CMH-2 have been found in plasmids it is not unreasonable hypothesize that CMH-3 (if chromosomal location) could be transferred to a plasmid and disseminated in the community.

In summary, the characterization of this *E. cloacae* complex isolate harbouring NDM in an IncX3 plasmid alerts about the spreading of the enzyme in our geographic area. The detection of CMH-3 in a more than likely chromosomal location and its potential to relocation in a plasmid must be considered. According to this, further assays including kinetics and binding affinities would be interesting to define the role of CMH-3 in the susceptibility to beta-lactams.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Ethical approval

Not required.

Competing interests

None declared.

Acknowledgments

This work was supported by Plan Nacional de I+D+i 13-2016 and Instituto de Salud Carlos III, Subdirección General de Redes y Centros de Investigación Cooperativa, Ministerio de Ciencia, Innovación y Universidades, Spanish Network for Research in

Infectious Diseases (REIPI RD16/0016/0001) cofinanced by European Development Regional Fund “A way to achieve Europe”, Operative program Intelligent Growth 2014–2020.

References

1. Davin-Regli A, Lavigne JP, Pagès JM. *Enterobacter* spp.: update on taxonomy, clinical aspects, and emerging antimicrobial resistance. Clin Microbiol Rev. 2019;32:e00002–19, <http://dx.doi.org/10.1128/CMR.00002-19>
2. Ingti B, Laskar MA, Choudhury S, Maurya AP, Paul D, Talukdar AD, et al. Molecular and in silico analysis of a new plasmid-mediated AmpC β -lactamase (CMH-2) in clinical isolates of *Klebsiella pneumoniae*. Infect Genet Evol. 2017;48:34–9, <http://dx.doi.org/10.1016/j.meegid.2016.12.009>
3. Ku Y, Lee M, Chuang Y, Yu WL. Detection of plasmid-mediated β -lactamase genes and emergence of a novel AmpC (CMH-1) in *Enterobacter cloacae* at a medical center in Southern Taiwan. J Clin Med. 2019;8, <http://dx.doi.org/10.3390/jcm8010008> E8.
4. Peirano G, Matsumura Y, Adams MD, Bradford P, Motyl M, Chen L, et al. Genomic epidemiology of global carbapenemase-producing *Enterobacter* spp., 2008–2014. Emerg Infect Dis. 2018;24:1010–9, <http://dx.doi.org/10.3201/eid2406.171648>
5. Villa J, Carretero O, Viedma E, Lora-Tamayo J, Mingorance J, Chaves F. Emergence of NDM-7 producing multidrug-resistant *Enterobacter hormaechei* sequence type ST78 in Spain, a high-risk international clone. Int J Antimicrob Agents. 2018, <http://dx.doi.org/10.1016/j.ijantimicag.2018.11.009>, pii: S0924-8579(18)30334-0 (Epub ahead of print).
6. Oteo J, Ortega A, Sola-Campoy PJ, Bautista V, Fernández-Romero, Sáez D. P2343 Spread of NDM-producing *Enterobacter cloacae* complex in Spain: whole-genome sequence analysis of the population structure, resistance genes and plasmids. In: Poster presented at: 28th European Congress of Clinical Microbiology and Infectious Diseases. 2018.
7. García-Castillo M, García-Fernández S, Gómez-Gil R, Pitart C, Oviaño M, Gracia-Ahufinger I, et al. Activity of ceftazidime–avibactam against carbapenemase-producing Enterobacteriaceae from urine specimens obtained during the infection-carbapenem resistance evaluation surveillance trial (iCREST) in Spain. Int J Antimicrob Agents. 2018;51:511–5, <http://dx.doi.org/10.1016/j.ijantimicag.2018.01.011>
8. Oteo J, Ortega A, Bartolomé R, Bou G, Conejo C, Fernández-Martínez M, et al. Prospective multicenter study of carbapenemase-producing Enterobacteriaceae from 83 hospitals in Spain reveals high in vitro susceptibility to colistin and meropenem. Antimicrob Agents Chemother. 2015;59:3406–12, <http://dx.doi.org/10.1128/AAC.00086-15>
9. European Committee on Antimicrobial Susceptibility Testing (EUCAST). Breakpoint tables for interpretation of MICs and zone diameters, version 8.0, valid from 2018-01-01; 2018.
10. European Committee on Antimicrobial Susceptibility Testing (EUCAST). Guidelines for detection of resistance mechanisms and specific resistances of clinical and/or epidemiological importance, version 2.0; July 2017.
11. Miró E, Agüero J, Larrosa MN, Fernández A, Conejo MC, Bou G, et al. Prevalence and molecular epidemiology of acquired AmpC β -lactamases and carbapenemases in Enterobacteriaceae isolates from 35 hospitals in Spain. Eur J Clin Microbiol Infect Dis. 2013;32:253–9, <http://dx.doi.org/10.1007/s10096-012-1737-0>
12. Poirel L, Walsh TR, Cuvillier V, Nordmann P. Multiplex PCR for detection of acquired carbapenemase genes. Diagn Microbiol Infect Dis. 2011;70:119–23, <http://dx.doi.org/10.1016/j.diagmicrobio.2010.12.002>
13. Rodríguez-Baño J, López-Cerero L, Navarro MD, Díaz de Alba P, Pascual A. Faecal carriage of extended-spectrum β -lactamase-producing *Escherichia coli*: prevalence, risk factors and molecular epidemiology. J Antimicrob Chemother. 2008;62:1142–9, <http://dx.doi.org/10.1093/jac/dkn293>
14. Qu H, Wang X, Ni Y, Liu J, Tan R, Huang J, et al. NDM-1-producing Enterobacteriaceae in a teaching hospital in Shanghai China: IncX3-type plasmids may contribute to the dissemination of blaNDM-1. Int J Infect Dis. 2015;34:8–13, <http://dx.doi.org/10.1016/j.ijid.2015.02.020>
15. Paskova V, Medvecký M, Skalova A, Chudejova K, Bitar I, Jakubu V, et al. Characterization of NDM-encoding plasmids From Enterobacteriaceae recovered from Czech hospitals. Front Microbiol. 2018;9:1549, <http://dx.doi.org/10.3389/fmicb.2018.01549>