



# Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



## Brief report

### Detection of a *cfr*-positive MRSA CC398 strain in a pig farmer in Spain

Laura Ruiz-Ripa<sup>a</sup>, Alba Bellés<sup>b</sup>, Mercè García<sup>b</sup>, Carmen Torres<sup>a,\*</sup>

<sup>a</sup> Área de Bioquímica y Biología Molecular, Universidad de La Rioja, Logroño, Spain

<sup>b</sup> Sección de Microbiología, Hospital Universitari Arnau de Vilanova, Lleida, Spain

#### ARTICLE INFO

##### Article history:

Received 15 January 2020

Accepted 13 March 2020

##### Keywords:

Methicillin-resistant *Staphylococcus aureus*

CC398

*cfr*

linezolid

#### ABSTRACT

**Objective:** To characterize one linezolid- and methicillin-resistant *Staphylococcus aureus* (MRSA) isolate recovered from a nasal sample of a pig farmer patient.

**Methods:** The detection of linezolid resistance mechanisms was performed by PCR and sequencing. The antimicrobial resistance and virulence profile was investigated, and the molecular typing was performed by molecular techniques. The transference of *cfr* gene was assessed by conjugation experiments and its genetic environment was investigated by specific PCRs.

**Results:** The linezolid-resistant MRSA isolate was typed as t011-ST398/CC398-SCCmecV-*agrI* and carried the *cfr* gene. The isolate was multidrug-resistant but lacked the virulence genes studied. The *cfr* gene was co-located with the *fexA* gene on a Tn558 variant and was successfully transferred by conjugation.

**Conclusion:** We report the first description of LA-MRSA-CC398 carrying the *cfr* gene in Spain. This finding highlights the importance of surveillance programmes to determine the presence and spread of the *cfr* gene in the livestock and clinical settings.

© 2020 Elsevier España, S.L.U. and Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. All rights reserved.

### Detección de una cepa SARM del linaje CC398 y portadora del gen *cfr* en un granjero de cerdos en España

#### RESUMEN

**Objetivo:** Caracterizar una cepa *Staphylococcus aureus* resistente a meticilina (SARM) y a linezolid aislada de una muestra nasal de un paciente granjero de cerdos.

**Métodos:** Mediante PCR y secuenciación se investigaron los mecanismos de resistencia a linezolid. Se determinó el perfil de resistencia a antimicrobianos y el perfil de virulencia, y se llevó a cabo el tipado molecular mediante diferentes técnicas moleculares. Se estudió la transferibilidad del gen *cfr* por conjugación y su entorno genético mediante PCR específicas.

**Resultados:** El aislado fue tipado como t011-ST398/CC398-SCCmecV-*agrI* y portaba el gen *cfr*. Presentó un fenotipo/genotipo de multiresistencia, pero carecía de los genes de virulencia estudiados. Se detectó el gen *cfr* junto con *fexA* en una variante del Tn558 y se transfirió mediante conjugación.

**Conclusión:** Se describe el primer aislado SARM-CC398 *cfr*-positivo en España. Esto destaca la importancia de implementar programas de vigilancia para determinar su presencia y dispersión en el ámbito clínico y ganadero.

© 2020 Elsevier España, S.L.U. y Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Todos los derechos reservados.

##### Palabras clave:

*Staphylococcus aureus* resistente a

meticilina

CC398

*cfr*

Linezolid

#### Introduction

*Staphylococcus aureus* clonal complex CC398 is the most common genetic lineage of livestock-associated (LA) methicillin-resistant *S. aureus* (MRSA) in Europe. Pigs are the main host of

\* Corresponding author.

E-mail address: carmen.torres@unirioja.es (C. Torres).

MRSA-CC398, however, its detection in humans is increasing in recent years either as colonizer or as causative agent of infection.<sup>1</sup>

Besides methicillin resistance, MRSA-CC398 usually exhibits a multiresistance phenotype which limits the therapeutic options. In this context, linezolid is an important antimicrobial agent for the treatment of serious infections caused by Gram-positive bacteria, including MRSA. Point mutations in the domain V of the 23S rRNA, as well as amino acid changes in the ribosomal proteins L3, L4, and L22 are the most common mechanisms of linezolid resistance in staphylococci.<sup>2</sup> Moreover, linezolid resistance mediated by acquired resistance genes is of concern due to they are often located on mobile genetic elements, which can move between bacteria by horizontal gene transfer. To date, three transferable linezolid resistance genes have been described in staphylococcal species: *cfr*, *optrA*, and *poxtA*.<sup>3–5</sup> Furthermore, these genes also confer reduced susceptibility to other antimicrobial agents, such as phenicols.<sup>2</sup>

In this study, we report, as far as we know, the first description of MRSA-CC398 harbouring the *cfr* gene in Spain, recovered from a nasal sample of a patient with professional contact with pigs.

## Methods

A pig farmer was hospitalized in the Intensive Care Unit (ICU) of the Hospital Universitari Arnau de Vilanova (Lleida, Spain) due to a non-infectious disease. Active surveillance for multiresistant bacteria was performed upon ICU admission, and MRSA were isolated from nasal, pharyngeal, and bronchoalveolar lavage samples, that were considered as colonization.

The susceptibility to penicillin, oxacillin, erythromycin, clindamycin, gentamicin, tobramycin, amikacin, minocycline, tetracycline, ciprofloxacin, levofloxacin, vancomycin, teicoplanin, fosfomicin, mupirocin, rifampicin, fusidic acid, daptomycin, and trimethoprim-sulfamethoxazole was determined by commercialized broth microdilution method (MicroScan, Beckman Coulter, Brea, USA) and to ceftiofloxacin by disc diffusion.<sup>6</sup> In addition, the minimum inhibitory concentration (MIC) to linezolid was determined by E-test<sup>®</sup> (bioMérieux, Durham, USA), and to chloramphenicol<sup>6</sup> and florfenicol<sup>7</sup> by broth microdilution using *S. aureus* ATCC<sup>®</sup> 29213 as quality control. All MRSA isolates recovered showed identical antimicrobial resistance phenotype and the isolate recovered from the nasal sample, named X1761, was further characterized.

The presence of the linezolid resistance genes *cfr*, *cfr(B)*, *optrA*, and *poxtA* was investigated by PCR (Supplementary Table S1). Moreover, mutations and amino acid changes in the deduced sequence of 23S rRNA and in the genes encoding the ribosomal proteins L3 (*rplC*), L4 (*rplD*), and L22 (*rplV*) were determined by PCR and amplicon sequencing (Supplementary Table S1). The obtained sequences were compared with those of linezolid-susceptible *S. aureus* subsp. *aureus* ATCC<sup>®</sup> 25923 (GenBank accession number CP009361) using the EMBOSS Needle software for nucleotide or amino acid (BLOSUM 62 cost matrix) alignments.

Based on the antimicrobial resistance phenotype, the presence of the following antimicrobial resistance genes was studied by PCR: *mecA*, *mecB*, *mecC*, *blaZ*, *lnu(A/A')*, *lnu(B)*, *lsa(B)*, *vga(A)*, *tet(K)*, *tet(L)*, *tet(M)*, *ant(4')-Ia*, *aac(6')-Ie-aph(2'')-Ia*, *fexA*, *fexB*, *cat<sub>PC194</sub>*, *cat<sub>PC221</sub>*, and *cat<sub>PC223</sub>* (Supplementary Table S1). Moreover, amino acid changes in the genes encoding the GyrA and GrlA proteins were investigated by PCR and amplicon sequencing (Supplementary Table S1) and compared with those of the fluoroquinolone-susceptible strain *S. aureus* subsp. *aureus* ATCC<sup>®</sup> 25923 (GenBank accession number CP009361). The presence of the virulence factors Panton-Valentine leukocidin (*lukS/F-PV*), toxic shock syndrome toxin-1 (*tst*), and exfoliative toxins (*eta*, *etb*, and *etd*) was determined by PCR (Supplementary Table S1).

The MRSA isolate was subjected to *spa*-typing by PCR and amplicon sequencing, and the obtained sequence was analyzed using

Ridom Staph-Type software (Ridom GmbH, Münster, Germany). Moreover, the isolate was characterized by *agr*-typing, multilocus sequence typing, and staphylococcal cassette chromosome *mec* (SCC*mec*) typing (Supplementary Table S1).

The transferability of the *cfr* gene was evaluated by the filter-mating method. A rifampicin-resistant mutant of *S. aureus* ATCC<sup>®</sup> 29213 was used as recipient strain, and the selection of the transconjugants was performed on BHI agar containing linezolid (4 mg/L) and rifampicin (50 mg/L). The transconjugants were identified by *spa*-typing. The antimicrobial resistance phenotype and genotype was determined by broth microdilution and by PCR, respectively (Supplementary Table S1).

The genetic environment of the *cfr* gene was determined by PCR mapping and amplicon sequencing. The primers were designed based on the structure of the *cfr*-carrying plasmids pSCFS3 and pSCFS7 (GenBank accession number AM086211 and FR675942) (Supplementary Table S2).

## Results and discussion

The MRSA isolate was resistant to thirteen of the twenty-three antimicrobial agents tested, including linezolid (MIC 8 mg/L), but remained susceptible to vancomycin and daptomycin. It carried the *mecA*, *blaZ*, *ant(4')-Ia*, *tet(K)*, *tet(M)*, *cfr*, and *fexA* resistance genes and showed the amino acid changes S80F and S84L in GrlA and GyrA proteins, respectively, related to fluoroquinolone resistance (Table 1). The *cfr* gene, which encodes a rRNA methyltransferase, mediates combined resistance to phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramins A antimicrobial agents. This multiresistance gene was originally detected in the plasmid pSCFS1 from *S. sciuri*,<sup>3</sup> and, thereafter, in different Gram-positive and Gram-negative bacteria of diverse origins. Although the use of linezolid is not approved in food-producing animals, the *cfr* gene is spread among staphylococcal isolates from pigs in different countries such as China,<sup>8</sup> which might be due to the co-selection of resistances. The linezolid-resistant MRSA isolate of this study also displayed the amino acid change V118A in the deduced sequence of *rplD* gene. There is no data about the relation of this change with linezolid susceptibility, but it seems to be associated with decreased susceptibility to the pleuromutilin lefamulin in *S. aureus* isolates.<sup>9</sup>

The X1761 isolate was typed as t011-ST398/CC398-SCC*mecV*-*agrI*. MRSA-CC398 isolates carrying the *cfr* gene have been described sporadically in different countries recovered primarily from livestock, but also from humans.<sup>10,11</sup> However, as far as we know, this is the first detection of MRSA CC398-*cfr* in Spain. A recent study conducted among Spanish hospitals, demonstrated the correlation between the pig density of the region and the presence of MRSA-CC398 in human infections at hospital level.<sup>1</sup> The hospital in which this strain was obtained is located in the Spanish region with the highest pig density in the year 2019 (370 pigs/km<sup>2</sup>); moreover, the patient was a pig farm worker, which is known to be a risk factor for human colonization with MRSA-CC398.<sup>12</sup>

The isolate did not harbour any of the virulence determinants studied, which is commonly observed among LA-MRSA CC398 isolates in comparison with human-adapted lineages.<sup>13</sup> The isolate lacked the IEC genes, which suggests an animal origin and, therefore, an animal to human transmission.

The *cfr* gene was successfully transferred by conjugation into *S. aureus* ATCC<sup>®</sup> 29213 together with the *fexA* gene. The transconjugants showed reduced susceptibility to clindamycin, chloramphenicol, florfenicol, and linezolid (Table 2), which is in accordance with the antimicrobial resistance genotype detected. The PCR mapping allowed us to identify the *cfr* and *fexA* genes as part of the Tn558 variant described on the plasmid pSCFS3 of *S. aureus* (GenBank accession number AM086211),<sup>14</sup> where the *cfr*

**Table 1**

Molecular typing, and antimicrobial resistance phenotype and genotype of the linezolid-resistant MRSA X1761 isolate.

Isolate	Molecular typing				Antimicrobial resistance	
	<i>spa</i> -type	MLST/CC	<i>agr</i>	SCC <i>mec</i>	Phenotype <sup>a</sup>	Genotype
X1761	t011	ST398/CC398	I	V	FOX-OXA-PEN-CLI-AMK-TOB-TET-MIN-CIP <sup>b</sup> -LEV-CHL-FFN-LZD	<i>mecA</i> , <i>blaZ</i> , <i>ant(4')</i> -Ia, <i>tet(K)</i> , <i>tet(M)</i> , <i>cfr</i> , <i>fexA</i>

<sup>a</sup> FOX, cefoxitin; OXA, oxacillin; PEN, penicillin; CLI, clindamycin; AMK, amikacin; TOB, tobramycin; TET, tetracycline; MIN, minocycline; CIP, ciprofloxacin; LEV, levofloxacin; CHL, chloramphenicol; FFN, florfenicol; LZD, linezolid.

<sup>b</sup> The isolate showed the amino acid changes S80F and S84L in GrlA and GyrA proteins, respectively.

**Table 2**Minimum inhibitory concentration (MIC) of the recipient *S. aureus* ATCC® 29213, the donor *S. aureus* X1761, and the transconjugant X1761-TC.

Isolate	MIC (mg/L)			
	Clindamycin	Chloramphenicol	Florfenicol	Linezolid
<i>S. aureus</i> ATCC® 29213	0.12	4	4	2
X1761	>128	128	≥256	8
X1761-TC	>128	128	128	8

gene is flanked by an IS21–558 element. The nucleotide alignment of the *cfr* genetic environment of the *S. aureus* X1761 isolate and the Tn558 variant of the plasmid pSCFS3 revealed the presence of six nucleotide exchanges in the non-coding region upstream the IS21–558 element. Although the *cfr* gene has been reported integrated in the chromosomal DNA, it has been more frequently detected on plasmids and, in many cases, with other resistance genes, including *fexA*.<sup>15</sup> Kehrenberg and Schwarz<sup>14</sup> described that the partial deletion of the transposase genes *tnpA* and *tnpB* in the Tn558 variant, due to the integration of the IS21–558 element, resulted in a transposition-deficient structure. Hence, the transference of the *cfr* gene by conjugation indicates its possible location on a plasmid, what would facilitate its dissemination.

In conclusion, the presence of transferable linezolid resistance genes in MRSA recovered from humans represents a public health concern, since it is considered a key antimicrobial agent in human medicine. MRSA-CC398 is a genetic lineage of the human-animal interface that frequently colonize pigs and is disseminated to in-contact humans. Consequently, the detection of a *cfr*-positive MRSA CC398 strain colonizing the nose of a pig farmer patient suggests that this multidrug-resistant strain might be circulating within the pig farms, what should be monitored in the future.

## Funding

This work was partially supported by project SAF2016-76571-R from the Agencia Estatal de Investigación (AEI) of Spain and the Fondo Europeo de Desarrollo Regional (FEDER) of EU. Laura Ruiz-Ripa has a pre-doctoral fellowship from the Universidad de La Rioja (Spain).

## Conflict of interest

The authors declare no conflict of interest.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.eimc.2020.03.006](https://doi.org/10.1016/j.eimc.2020.03.006).

## References

- Ceballos S, Spiroz C, Ruiz-Ripa L, Reynaga E, Azcona-Gutiérrez JM, Rezusta A, et al. Epidemiology of MRSA CC398 in hospitals located in Spanish regions with

different pig-farming densities: a multicentre study. *J. Antimicrob. Chemother.* 2019;74:2157–61, [http://dx.doi.org/10.1093/jac/dkz180](https://doi.org/10.1093/jac/dkz180).

- Mendes RE, Deshpande LM, Jones RN. Linezolid update: stable in vitro activity following more than a decade of clinical use and summary of associated resistance mechanisms. *Drug Resist. Updat.* 2014;17:1–12, [http://dx.doi.org/10.1016/j.drug.2014.04.002](https://doi.org/10.1016/j.drug.2014.04.002).
- Schwarz S, Werckenthin C, Kehrenberg C. Identification of a plasmid-borne chloramphenicol-florfenicol resistance gene in *Staphylococcus sciuri*. *Antimicrob. Agents Chemother.* 2000;44:2530–3, [http://dx.doi.org/10.1128/aac.44.9.2530-2533.2000](https://doi.org/10.1128/aac.44.9.2530-2533.2000).
- Wang Y, Lv Y, Cai J, Schwarz S, Cui L, Hu Z, et al. A novel gene, *optrA*, that confers transferable resistance to oxazolidinones and phenicols and its presence in *Enterococcus faecalis* and *Enterococcus faecium* of human and animal origin. *J. Antimicrob. Chemother.* 2015;70:2182–90, [http://dx.doi.org/10.1093/jac/dkv116](https://doi.org/10.1093/jac/dkv116).
- Antonelli A, D'Andrea MM, Brenciani A, Galeotti CL, Morroni G, Pollini S, et al. Characterization of *poxA*, a novel phenicol-oxazolidinone-tetracycline resistance gene from a MRSA of clinical origin. *J. Antimicrob. Chemother.* 2018;73:1763–9, [http://dx.doi.org/10.1093/jac/dky088](https://doi.org/10.1093/jac/dky088).
- The European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MICs and zone diameters. Version 9.0, 2019. <http://www.eucast.org>.
- Clinical and Laboratory Standards Institute (CLSI), 2018. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals: 4th edition. CLSI supplement VET08. Wayne, PA, USA.
- Wang J, Lin DC, Guo XM, Wei HK, Liu XQ, Chen XJ, et al. Distribution of the multidrug resistance gene *cfr* in *Staphylococcus* isolates from pigs, workers, and the environment of a hog market and a slaughterhouse in Guangzhou, China. *Foodborne Pathog Dis.* 2015;12:598–605, [http://dx.doi.org/10.1089/fpd.2014.1891](https://doi.org/10.1089/fpd.2014.1891).
- Mendes RE, Paukner S, Doyle TB, Gelone SP, Flamm RK, Sader HS. Low prevalence of Gram-positive isolates showing elevated lefamulin MIC results during the SENTRY Surveillance Program for 2015–2016 and characterization of resistance mechanisms. *Antimicrob. Agents Chemother.* 2019;63:e02158–2218, [http://dx.doi.org/10.1128/AAC.02158-18](https://doi.org/10.1128/AAC.02158-18).
- Paridaens H, Coussement J, Argudin MA, Delaere B, Huang TD, Glupczynski Y, et al. Clinical case of *cfr*-positive MRSA CC398 in Belgium. *Eur. J. Clin. Microbiol. Infect. Dis.* 2017;36:1527–9, [http://dx.doi.org/10.1007/s10096-017-2953-4](https://doi.org/10.1007/s10096-017-2953-4).
- Peeters LE, Argudin MA, Azadikhah S, Butaye P. Antimicrobial resistance and population structure of *Staphylococcus aureus* recovered from pigs farms. *Vet. Microbiol.* 2015;180:151–6, [http://dx.doi.org/10.1016/j.vetmic.2015.08.018](https://doi.org/10.1016/j.vetmic.2015.08.018).
- Van Den Broek IV, Van Cleef BA, Haenen A, Broens EM, Van Der Wolf PJ, Van Den Broek MJ, et al. Methicillin-resistant *Staphylococcus aureus* in people living and working in pig farms. *Epidemiol. Infect.* 2009;137:700–8, [http://dx.doi.org/10.1017/S0950268808001507](https://doi.org/10.1017/S0950268808001507).
- Argudin MA, Tenhagen BA, Fetsch A, Sachsenröder J, Käsböhrer A, Schroeter A, et al. Virulence and resistance determinants of German *Staphylococcus aureus* ST398 isolates from nonhuman sources. *Appl. Environ. Microbiol.* 2011;77:3052–60.
- Kehrenberg C1, Schwarz S. Distribution of florfenicol resistance genes *fexA* and *cfr* among chloramphenicol-resistant *Staphylococcus* isolates. *Antimicrob. Agents Chemother.* 2006;50:1156–63, [http://dx.doi.org/10.1128/AEM.02260-10](https://doi.org/10.1128/AEM.02260-10).
- Lozano C, Ruiz-García M, Gómez-Sanz E, López-García P, Royo-García G, Zarazaga M, et al. Characterization of a *cfr*-positive methicillin-resistant *Staphylococcus epidermidis* strain of the lineage ST22 implicated in a life-threatening human infection. *Diagn. Microbiol. Infect. Dis.* 2012;73:380–2, [http://dx.doi.org/10.1016/j.diagmicrobio.2012.04.013](https://doi.org/10.1016/j.diagmicrobio.2012.04.013).