



## Case Report

# Obstructive sleep apnea syndrome in patients attending a psychiatry outpatient service: A case series<sup>☆</sup>



Nathalie Tamayo Martínez\*, Diego Rosselli Cock

Departamento de Epidemiología Clínica y Bioestadística, Pontificia Universidad Javeriana, Bogotá, Colombia

### ARTICLE INFO

#### Article history:

Received 4 January 2016

Accepted 15 August 2016

Available online 18 October 2017

#### Keywords:

Obstructive sleep apnea syndrome

Depressive disorder

Anxiety disorders

Schizophrenia

Bipolar disorder

Observational study

### ABSTRACT

**Background:** Obstructive sleep apnea syndrome (OSAS) is a condition associated with multiple negative outcomes. People with mental illness might be at increased risk of having it, given that medication given has adverse effects on weight and there are alterations in sleep associated with them; however, there are few studies in this population.

**Objective:** Describe the patients and the results of polysomnography ordered based on clinical symptoms in a psychiatric outpatient clinic between 2012 and 2014.

**Methods:** A case series in which medical records were evaluated.

**Results:** 58 patients who underwent polysomnography, 89% of them had OSAS, 16% were obese and 19% were treated with benzodiazepines.

**Conclusions:** This is a condition that must be considered during the clinical evaluation of patients with mental illness, since its presence should make clinicians think about drug treatment and follow up.

© 2016 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

## Síndrome de apnea obstructiva del sueño en personas atendidas en consulta externa de psiquiatría: serie de casos

### RESUMEN

El síndrome de apnea obstructiva del sueño (SAOS) se asocia a múltiples desenlaces negativos. Se ha propuesto que las personas con enfermedad mental están en mayor riesgo, en parte por sobrepeso y por las alteraciones del sueño asociadas con algunos medicamentos. Sin embargo, son pocos los estudios en esta población.

#### Palabras clave:

Síndrome de apnea obstructiva del sueño

sueño

Trastorno depresivo

DOI of original article: <http://dx.doi.org/10.1016/j.rcp.2016.08.007>.

\* Please cite this article as: Tamayo Martínez N, Rosselli Cock D. Síndrome de apnea obstructiva del sueño en personas atendidas en consulta externa de psiquiatría: serie de casos. Rev Colomb Psiquiatr. 2017;46:243-246.

\* Corresponding author.

E-mail address: [nathalietamayo@gmail.com](mailto:nathalietamayo@gmail.com) (N. Tamayo Martínez).

<http://dx.doi.org/10.1016/j.rcp.2017.09.006>

2530-3120/© 2016 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Trastornos de ansiedad  
Esquizofrenia  
Trastorno afectivo bipolar  
Reporte de casos

Objetivo: Describir a la población y el resultado de las polisomnografías solicitadas ante sospecha clínica en pacientes de consulta externa de una clínica psiquiátrica.

Métodos: Estudio descriptivo de una muestra de pacientes consecutivos atendidos entre 2012 y 2014.

Resultados: De los 58 pacientes de los que se solicitó polisomnografía, 52 (89%) presentaban SAOS. De estos, el 16% cursaba con obesidad y el 19% tomaba benzodiacepinas.

Conclusiones: Esta es una enfermedad que se debe tener en cuenta durante la evaluación clínica de los pacientes con enfermedad mental, dado que su presencia implica precaución al plantear el tratamiento farmacológico y hacer el seguimiento.

© 2016 Asociación Colombiana de Psiquiatría. Publicado por Elsevier España, S.L.U.  
Todos los derechos reservados.

## Introduction

Obstructive sleep apnoea syndrome (OSAS) is classified as a “breathing-related sleep disorder” and is characterised by repetitive closure of the upper airway during sleep, with decreased arterial oxygen saturation. The standard diagnostic method is polysomnography, and severity is classified according to the number of apnoea and hypopnoea events per hour. This disorder is mild if there are between 5 and 15 events plus some symptoms (such as excessive sleepiness, unrefreshing sleep, fatigue, insomnia, difficulty breathing at night, choking sensation during the night, breathing interruptions or loud snoring); moderate if there are more than 15 but less than 30, and severe when there are more than 30 events/h.<sup>1</sup>

OSAS is a highly prevalent disorder that is estimated to affect between 4% and 8% of adults, and has significant negative repercussions on quality of life and considerable associated morbidity and mortality.<sup>2-4</sup> Although OSAS was defined in the 1970s, its negative impact on mortality and quality of life was only documented some 20 years ago. OSAS patients are at greater risk of hypertension, insulin resistance, diabetes mellitus, coronary events, cerebrovascular events, cognitive alterations, perioperative complications, in-hospital mortality and all-cause mortality, as well as occupational and traffic accidents.<sup>5-7</sup> This is why it is so important to make the initial diagnosis and start treatment. Risk factors for OSAS include obesity, male gender, aged over 50, family history, nasal obstruction, use of alcohol or sedatives, cigarette smoking, gastro-oesophageal reflux, hypothyroidism and acromegaly, among others.<sup>8,9</sup>

Few studies have been published on the association between OSAS and the development, prognosis and impact of mental illness. It has been suggested that individuals with mental illness may be at increased risk of suffering from OSAS due to weight gain caused by most antipsychotics and mood stabilisers, the presence of respiratory events related to the use of sedative drugs, and a less healthy lifestyle.<sup>10-14</sup> A search of PubMed using the MeSH terms “Depressive Disorder, Major” and “Sleep Apnoea, Obstructive” retrieved 18 articles, one of which found that, according to a study of a database of over 2 million obese individuals, OSAS patients are at increased risk of depression (odds ratio [OR]=1.85; confidence interval [CI], 1.80-1.88;  $p < 0.001$ ).<sup>15</sup> Another study evaluated 53 people with coronary heart disease, and found a similar prevalence of depression in patients with and without OSAS.<sup>16</sup> However,

the small sample size prevented the authors from drawing definitive conclusions. A more recent systematic review found that up to 69% of individuals with bipolar affective disorder presented OSAS, albeit in studies with a high risk of selection bias.<sup>17</sup> A systematic review of the literature on people with major mental illness found a prevalence of 36.3% for OSAS in people with major depressive disorder, 24.5% in patients with bipolar affective disorder and 15.4% in patients with schizophrenia. However, a review of the articles included showed that some had requested polysomnography due to suspicion of OSAS, such as medication-induced symptoms of daytime sleepiness, and this may have distorted the true prevalence.<sup>18</sup> Furthermore, a study in older people found that apnoea-hypopnoea is associated with poor cognitive functioning and increased risk of dementia.<sup>19,20</sup>

The aim of this study is to describe the general characteristics and polysomnographic parameters of a sample of psychiatric patients seen on an outpatient basis in a psychiatric clinic.

## Material and methods

This is a descriptive study of a series of consecutive patients undergoing polysomnography for clinical suspicion of OSAS in a psychiatric clinic. The outpatient department of the clinic attends approximately 5000 adult patients per month, whose health insurance mainly covers individuals living in the city of Bogotá. Data from the medical history of patients seen in a psychiatry outpatient clinic between 2012 and 2014 were collected. The information obtained from the medical records was organised and summarised in a Microsoft® Excel® 2013 spreadsheet, which also calculated the summary measures of the findings using measures of central tendency and dispersion. The study was approved by the institution's ethics committee and is classified as risk-free research, since data were taken from the medical history and no intervention or data collection was performed.<sup>21</sup>

## Results

**Table 1** shows the general characteristics of the study sample formed of 58 adults, mostly female, half of whom were overweight. The median Epworth Sleepiness Scale score was 7 (0-24) points, and 52 patients (89%) met OSAS diagnostic

**Table 1 – General characteristics of patients (n = 58).**

Women	44 (76)
Age (years)	61 (37-80) [53-65]
Body mass index	
<25	19 (33)
25.0-29.9	26 (45)
30.0-34.9	9 (15)
>35.0	4 (7)
Apnoeas/h	32 ± 22
OSAS (apnoea-hypopnoea index/h)	
No apnoea	6 (10)
5-14 events plus daytime symptoms	10 (17)
15-30 events	14 (24)
More than 30 events	28 (48)
Diagnoses	
Depressive and anxiety disorders	35 (58)
Bipolar affective disorder	8 (13)
Schizophrenia or schizoaffective disorder	6 (10)
Sleep disorders	4 (7)
Cognitive impairment and Parkinson's disease	4 (7)
Other	3 (5)
Medication received	
Selective serotonin reuptake inhibitors	25 (43)
Trazodone	16 (28)
Benzodiazepines	11 (19)
Clozapine	10 (17)
Mood stabilisers	9 (16)
Typical antipsychotics	3 (5)
Atypical antipsychotics	2 (3)
Dual-action antidepressants	2 (3)
Non-benzodiazepine hypnotics	2 (3)
Tricyclic antidepressants	1 (2)

The values are expressed as n (%) or median (range) [interquartile range].

criteria. The patients were diagnosed with a variety of disorders, some patients with more than one diagnosis; 16 (28%) received 1 medication, 25 (43%) received 2, 12 (20%) received none, and the other 5 (9%) received 3 or more. The antidepressants prescribed were, in order of frequency: fluoxetine, sertraline, escitalopram, duloxetine, mirtazapine, amitriptyline and trazodone; the benzodiazepines lorazepam, clonazepam and alprazolam, the mood stabilisers lithium carbonate and valproic acid, the typical antipsychotics haloperidol, levomepromazine and piperazine, the atypical antipsychotics risperidone, aripiprazole and clozapine and the non-benzodiazepine hypnotic zolpidem.

## Discussion

This study is a preliminary evaluation of the importance of recognising OSAS in patients with mental illness in Colombia. Due to the sampling strategy used, we were unable to calculate the prevalence of this condition, since polysomnography was ordered for the clinical suspicion of sleep disorder, and therefore our sample is not representative of the population

with mental illness. Despite these limitations, it is interesting to observe that 89% of patients undergoing polysomnography had an apnoea-hypopnoea index of at least 5 events/h associated with daytime symptoms.<sup>1</sup> A small proportion (16%) were obese or taking benzodiazepines (19%). These results should be interpreted with caution, since the polysomnography was ordered on the basis of a clinical suspicion of OSAS, and the population studied consisted of overweight elderly women, which is a risk factor for this disorder.<sup>6</sup>

Few studies have evaluated the prevalence of OSAS in patients with mental illness, and those that have been published have small samples and are based, as in this case, on polysomnography ordered due to clinical suspicion, and for that reason are not representative of individuals with mental illness. One study,<sup>22</sup> for example, found an OSAS prevalence of 39% in a population of 51 patients with major depressive disorder, while another,<sup>23</sup> reported prevalence of 59% in individuals with major depressive disorder or bipolar affective disorder. The only study comparing patients with various diagnoses<sup>10</sup> found a prevalence of OSAS of 46% in men and 58% in women with schizophrenia, 20% of men and 3% of women with major depressive disorder, and 27% of men and 15% of women with bipolar affective disorder referred to polysomnography on suspicion of OSAS. Finally, another study<sup>24</sup> found that 21% of patients with bipolar affective disorder were referred to polysomnography on clinical suspicion.

## Conclusions

OSAS must be taken into account during the clinical evaluation of patients with mental illness. In the presence of this disorder, caution must be taken when prescribing pharmacological treatment and undertaking follow-up. Few studies of this disorder in patients with mental illness have been published, but the findings suggest that the prevalence is higher than that reported in the general population.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

**Confidentiality of data.** The authors declare that they have followed the protocols of their work center on the publication of patient data.

**Right to privacy and informed consent.** The authors have obtained the written informed consent of the patients or subjects mentioned in the article. The corresponding author is in possession of this document.

## Funding

This study did not receive any funding.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## REFERENCES

1. Epstein LJ, Kristo D, Strollo PJ, Friedman N, Malhotra A, Patil SP, et al. Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med.* 2009;5:263-76.
2. Kryger MH, Roth T, Dement WC. Principles and practice of sleep medicine: expert consult premium edition. Philadelphia: Elsevier Health Sciences; 2010.
3. Kushida CA, editor. Obstructive sleep apnea: pathophysiology, comorbidities and consequences. 1.<sup>a</sup> ed. CRC Press; 2007.
4. Ryan S, Taylor CT, McNicholas WT. Predictors of elevated nuclear factor-kappaB-dependent genes in obstructive sleep apnea syndrome. *Am J Respir Crit Care Med.* 2006;174:824-30.
5. Ejaz SM, Khawaja IS, Bhatia S, Hurwitz TD. Obstructive sleep apnea and depression: a review. *Innov Clin Neurosci.* 2011;8:17-25.
6. Vozoris NT. Sleep apnea-plus: prevalence, risk factors, and association with cardiovascular diseases using United States population-level data. *Sleep Med.* 2012;13:637-44.
7. Young T, Palta M, Dempsey J, Peppard PE, Nieto FJ, Hla KM. Burden of sleep apnea: rationale, design, and major findings of the Wisconsin Sleep Cohort study. *WMJ Off Publ State Med Soc Wis.* 2009;108:246-9.
8. Al Lawati NM, Patel SR, Ayas NT. Epidemiology, risk factors, and consequences of obstructive sleep apnea and short sleep duration. *Prog Cardiovasc Dis.* 2009;51:285-93.
9. Shepherd KL, James AL, Musk AW, Hunter ML, Hillman DR, Eastwood PR. Gastro-oesophageal reflux symptoms are related to the presence and severity of obstructive sleep apnoea. *J Sleep Res.* 2011;20 Pt 2:241-9.
10. Winkelman JW. Schizophrenia, obesity, and obstructive sleep apnea. *J Clin Psychiatry.* 2001;62:8-11.
11. Alam A, Chengappa KNR, Ghinassi F. Screening for obstructive sleep apnea among individuals with severe mental illness at a primary care clinic. *Gen Hosp Psychiatry.* 2012;34:660-4.
12. Soreca I, Levenson J, Lotz M, Frank E, Kupfer DJ. Sleep apnea risk and clinical correlates in patients with bipolar disorder. *Bipolar Disord.* 2012;14:672-6.
13. Forbus L, Kelly UA. Screening for obstructive sleep apnea in veterans seeking treatment of posttraumatic stress disorder. *ANS Adv Nurs Sci.* 2015;38:298-305.
14. Annamalai A, Palmese LB, Chwastiak LA, Srihari VH, Tek C. High rates of obstructive sleep apnea symptoms among patients with schizophrenia. *Psychosomatics.* 2015;56:59-66.
15. Babson KA, Del Re AC, Bonn-Miller MO, Woodward SH. The comorbidity of sleep apnea and mood, anxiety, and substance use disorders among obese military veterans within the Veterans Health Administration. *J Clin Sleep Med.* 2013;9:1253-8.
16. Carney RM, Freedland KE, Duntley SP, Rich MW. Obstructive sleep apnea and major depressive disorder in cardiovascular disease. *Int J Cardiol.* 2011;149:283-4.
17. Gupta MA, Simpson FC. Obstructive sleep apnea and psychiatric disorders: a systematic review. *J Clin Sleep Med.* 2015;11:165-75.
18. Stubbs B, Vancampfort D, Veronese N, Solmi M, Gaughran F, Manu P, et al. The prevalence and predictors of obstructive sleep apnea in major depressive disorder, bipolar disorder and schizophrenia: a systematic review and meta-analysis. *J Affect Disord.* 2016;197:259-67.
19. Ramos AR, Tarraf W, Rundek T, Redline S, Wohlgemuth WK, Loredo JS, et al. Obstructive sleep apnea and neurocognitive function in a Hispanic/Latino population. *Neurology.* 2015;84:391-8.
20. Chang W-P, Liu M-E, Chang W-C, Yang AC, Ku Y-C, Pai J-T, et al. Sleep apnea and the risk of dementia: a population-based 5-year follow-up study in Taiwan. *PLOS ONE.* 2013;8:e78655.
21. Resolución 8430. Normas científicas, técnicas y administrativas para la investigación en salud. Bogotá: Ministerio de Salud, República de Colombia; 1993.
22. Ong JC, Gress JL, San Pedro-Salcedo MG, Manber R. Frequency and predictors of obstructive sleep apnea among individuals with major depressive disorder and insomnia. *J Psychosom Res.* 2009;67:135-41.
23. Hattori M, Kitajima T, Mekata T, Kanamori A, Imamura M, Sakakibara H, et al. Risk factors for obstructive sleep apnea syndrome screening in mood disorder patients. *Psychiatry Clin Neurosci.* 2009;63:385-91.
24. Kelly T, Douglas L, Denmark L, Brasuell G, Lieberman DZ. The high prevalence of obstructive sleep apnea among patients with bipolar disorders. *J Affect Disord.* 2013;151:54-8.